

Response of the Intertropical Convergence Zone to Antarctic Ice Sheet melt

P. Bakker¹ * and M. Prange¹

P. Bakker, p.bakker@vu.nl

¹MARUM-Center for Marine

Environmental Sciences, University of
Bremen, Bremen, Germany

*Now at Faculty of Earth and Life
Sciences, Vrije Universiteit Amsterdam, The
Netherlands

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the Version of Record. Please cite this article as doi: 10.1029/2018GL078659

©2018 American Geophysical Union. All Rights Reserved.

Past cooling events in the Northern Hemisphere have been shown to impact the location of the intertropical convergence zone (ITCZ) and thereby induce a southward shift of tropical precipitation. Here we use high-resolution coupled ocean-atmosphere simulations to show that reasonable past melt rates of the Antarctic Ice Sheet can similarly have led to shifts of the ITCZ, albeit in opposite direction, through large-scale surface air temperature changes over the Southern Ocean. Through sensitivity experiments employing slightly negative to large positive meltwater fluxes we deduce that meridional shifts of the Hadley cell and therewith the ITCZ are, to a first order, a linear response to Southern Hemisphere high-latitude surface air temperature changes and Antarctic Ice Sheet melt rates. This highlights the possibility to use past episodes of anomalous melt rates to better constrain a possible future response of low latitude precipitation to continued global warming and a shrinking Antarctic Ice Sheet.

Keypoints:

- Realistic past and future melt rates of the Antarctic Ice Sheet have impact on Southern Ocean
- Southern Ocean surface air temperature changes lead to ITCZ shifts

Plain Language Summary

Changes in high latitude climate can impact the tropical regions through so-called atmospheric and oceanic teleconnections. Research has mostly focused on past southward shifts in the band of heavy tropical precipitation, called the intertropical convergence zone (ITCZ), linked to large-scale cooling in the Northern Hemisphere resulting from large-scale continental ice sheet buildup or a slowdown of the large-scale Atlantic meridional ocean circulation. Here we use high resolution climate simulations to show that melting of the Antarctic Ice Sheet can similarly lead to northward shifts of the ITCZ and the displacement of the accompanying rain belt. Future melt rates of the Antarctic Ice Sheet are highly uncertain, but our work shows that it might have a non-negligible impact on the tropical climate. Moreover, we find that because of the apparent linearity of the system under consideration, studying episodes of past changes in the size of the Antarctic Ice Sheet can help us constrain the possible changes in the low latitude hydroclimate.

1. Introduction

The intertropical convergence zone (ITCZ), a band of intense precipitation and cloud cover encircling the Earth near the equator associated with the ascending branch of the Hadley cell, has been shown early on to be intimately linked with the interhemispheric temperature contrast [Manabe and Stouffer, 1980]. Broccoli *et al.* [2006] found that a Northern Hemisphere (NH) extratropical cooling causes an increase in the total atmospheric energy flux from the tropics to the extratropics of the cooling hemisphere while the opposite happens in the Southern Hemisphere (SH). The resulting low latitude imbalance in the poleward energy fluxes leads to a reorganization of the mean meridional

tropical circulation (Hadley cell), effectively transporting more energy from the SH low latitudes across the equator to the NH low latitudes [Broccoli *et al.*, 2006]. As a result, the ITCZ moves away from the cooling hemisphere [e.g. Schneider *et al.*, 2014].

Because of the dependency on the interhemispheric temperature contrast, ITCZ displacements on timescales ranging from seasonal, decadal to glacial-interglacial have been linked to temperature changes at mid-to-high latitudes [Donohoe *et al.*, 2013], including the build-up of continental ice sheets and North Atlantic cooling as a result of disruptions of the Atlantic Meridional Overturning Circulation [AMOC; Chiang *et al.*, 2003; Chiang and Bitz, 2005; Zhang and Delworth, 2005; Mulitza *et al.*, 2017]. Most studies thus far have investigated the response of the ITCZ to NH cooling, but also the SH mid-to-high latitudes can potentially exhibit strong cooling, for instance as a result of enhanced melt-water input into the Southern Ocean from the Antarctic Ice Sheet (AIS). A large-scale cooling of the Southern Ocean as a response to AIS melt has been shown with a range of different climate models, e.g. intermediate complexity climate models [Swingedouw *et al.*, 2009; Bakker *et al.*, 2017], high resolution ocean only models [Stammer, 2008; Morrison, 2015] and coupled ocean-atmosphere general circulation models [Ma and Wu, 2011; Ma *et al.*, 2013]. Because of the lack or highly simplified nature of atmospheric dynamics, the first two types of models do not allow to investigate the impact of Southern Ocean cooling on the ITCZ. The studies using coupled general circulation models did indeed find an impact of Southern Ocean cooling on low latitude climates [Ma and Wu, 2011] as well as changes in the characteristics of ENSO [Ma *et al.*, 2013], but did not analyze the impact on the general atmospheric circulation in the tropics.

Recent measurements show that the AIS has lost a considerable amount of ice over the last decades [*The IMBIE team*, 2018]. Periods of enhanced meltwater input into the Southern Ocean from the AIS are also thought to have occurred in the past and are projected to occur in the future under the influence of continued global warming. One of the largest pulses of freshwater into the Southern Ocean is postulated to have occurred during the last deglaciation in the so-called meltwater pulse 1A event [*Clark et al.*, 1996]. The magnitude of this event [*Golledge et al.*, 2014], although highly uncertain, is similar to the AIS melt rates that are projected for the end of the century under continued future global warming [*deConto and Pollard*, 2016], up to several tenths of a Sverdrup ($1\text{Sv} = 10^6 \text{m}^3\text{s}^{-1}$). Less dramatic, but more recent and reoccurring episodes of AIS growth and melt were recently suggested by *Bakker et al.* [2017] as part of a multi-centennial variability in the coupled AIS-climate system.

Here we extend on previous work [e.g. *Ma et al.*, 2013] by employing a state-of-the-science coupled climate model at relatively high spatial resolution and test a range of meltwater fluxes that are realistic in light of estimated past and future changes in the AIS. We focus specifically on the simulated changes at low latitudes resulting from the meltwater induced changes in SH high-latitude surface conditions and the relationship between the magnitude of the meltwater-induced temperature anomalies and the resulting climatic change in the low latitudes. The latter allows us to deduce whether or not the results are, in the context of this climate model, more generally relevant for the study of past and future climate change regardless of the exact magnitude or even the sign of the imposed melt rates.

2. Methods

The simulations were performed using the Community Earth System Model version 1.2 (CESM1.2), a global climate model that includes interactive atmosphere (CAM4), ocean (POP2), land (CLM4.0; including carbon-nitrogen dynamics), and sea-ice (CICE4) components. For the atmosphere (running with a finite volume dynamical core) and land a horizontal resolution of $0.9^\circ \times 1.25^\circ$ was used with the former having 26 vertical levels. The ocean and sea-ice components use a displaced dipole grid with a nominal horizontal resolution of 1° . The ocean grid has 60 levels.

The AIS contribution to meltwater pulse 1A [Clark *et al.*, 1996] is highly uncertain, but in the most thorough attempt thus far to quantify this flux, a mean value of 0.034 Sv was found, with maximum values up to 0.11 Sv for a period of 350 years [Golledge *et al.*, 2014].

The Holocene AIS variability suggested by Bakker *et al.* [2017] is 0.048 Sv (1σ). Finally, the future response of the AIS if global warming is to continue in the next centuries again varies widely, ranging from ~ 0.2 -0.5 Sv for the year 2100 in, respectively the so-called Representative Concentration Pathway (RCP) scenario 4.5 and RCP8.5 [Meinshausen *et al.*, 2011], and ranging from ~ 0.2 -0.25 Sv for the year 2500 in again RCP4.5 and RCP8.5, respectively [deConto and Pollard, 2016]. Taken together, credible past and future rates of AIS meltwater into the Southern Ocean seem to range from slightly negative values, i.e. periods of minor AIS growth [Bakker *et al.*, 2017], to positive values of several tenths of Sverdrups. To assess the impact of such AIS meltwater rates we performed four 200 year long experiments in which different magnitudes of freshwater forcing (FWF), namely -48 mSv, 48 mSv, 100 mSv and 200 mSv, were continuously added to the surface of the

Southern Ocean south of 60°S (referred to as PIm48, PI48, PI100 and PI200, respectively).

The negative -48 mSv meltwater flux implies that freshwater is removed from the internally calculated freshwater flux that enters the ocean, comprising of precipitation, continental runoff and sea-ice melt. AIS freshwater input is not compensated for elsewhere. All experiments start from the same spinup pre-industrial state and in addition a 200 year long control simulation was performed in which no additional freshwater was added to the Southern Ocean (referred to as 'PI'). For the analyses averages over the last 20 years of each simulation are taken.

To investigate the robustness of the relationship between the magnitude of the imposed AIS meltwater forcing and the resulting climatic impact, we calculate for every grid cell the correlation between the imposed AIS meltwater forcing and the climatic variable under consideration (e.g. annual mean temperature anomalies), only showing the result if the linear correlation coefficient exceeds the critical value (for a two-sided t-test with five data pairs and $\alpha=0.1$) of 0.805 and presenting the relationship as the slope of the linear regression. In the remainder of the manuscript the term 'robust' is thus used in the context of a significant linear correlation between the imposed FWF and the resulting climatic impact, indicating that the effect is to a first order linear over the range of applied positive and negative FWFs.

3. Results

The imposed changes in surface salinity in the Southern Ocean south of 60°S induce a strong response of the coupled ocean-atmosphere-sea ice system. A decrease of sea-surface salinity leads to a stabilization of the water column in the Southern Ocean, reduced vertical

mixing and convection (not shown) and a resulting cooling of the surface (Figure 1) and an increased sea ice cover (not shown), in line with previous studies [*Stammer, 2008; Morrison, 2015; Ma and Wu, 2011; Ma et al., 2013; Swingedouw et al., 2009; Bakker et al., 2017*]. For the negative FWF experiment PIm48, the increase in surface salinity leads to a surface warming. The largest cooling, up to 2 K per 100 mSv of AIS meltwater, takes place in the Southern Ocean, but a robust temperature response can be found almost globally, with the exception of the NH mid-latitudes (Figure 1). For precipitation the situation looks quite different. A robust annual mean drying of up to 100 mm year⁻¹ per 100 mSv of AIS meltwater is simulated over the Southern Ocean and the Antarctic continent (Figure 1). However, outside of the SH high latitudes, a robust precipitation response is largely confined to the low latitudes of the eastern/central Pacific and the Atlantic. In both regions a positive AIS meltwater forcing leads to a drying south of the equator and a precipitation increases to the north of the equator, most clearly so in the eastern Pacific. However, here the negative precipitation response to AIS melt is likely to be overestimated due to the simulation of an unrealistic double ITCZ. As such, the Southern Ocean cooling may partly remedy this well-known model artefact [cf. *Hwang and Frierson, 2013*]. Note that no robust precipitation response is simulated over the low latitude continents or over the Indian ocean except for the West African, Australian and Venezuelan monsoon regions, which are known to be sensitively affected by ITCZ shifts [*Mohtadi et al., 2016*]. In line with the findings of *Baker et al. [2018]* we find that, in contrast to the temperature anomalies (not shown), the precipitation anomalies in the low latitudes show a clear seasonal dependency (Supporting Information Figure

1). Nonetheless, the overall picture of meridional shifts of the low latitude precipitation maxima in the eastern Pacific and Atlantic persists.

This spatial pattern of low latitude precipitation anomalies is a strong indication of meridional shifts of the ITCZ, northward for a Southern Ocean cooling and southward for a Southern Ocean warming. To further investigate these changes in low latitude atmospheric circulation, we plot the meridional atmospheric streamfunction changes as a function of the applied meltwater forcing (Figure 2) and find that for a positive AIS melt rate the Hadley cell to the south of the equator strengthens while it weakens to the north of the equator, involving a northward shift of the mean ITCZ. We find that there is to the first order a linear negative relationship between the degree of surface air temperature change in the SH high-latitudes (south of 50°S) and the meridional location of the ITCZ (Figure 3; the ITCZ location is defined here as the latitude of zero meridional streamfunction at 500hPa in the low latitudes). Moreover, we find positive (negative) linear relationships between SH high-latitude surface air temperature changes and the change in strength of the northerly maximum (southerly minimum) of the Hadley cell (Figure 3). We focus here on the relationship between these metrics of the low latitude atmospheric circulation and SH high-latitude surface air temperatures because of the mechanistic link, however, the relationships are very similar if we look at the imposed FWFs because those are in turn closely related to SH high-latitude surface air temperatures (Supporting Information Figure 2).

In line with the findings of *Broccoli et al.* [2006] our results show that changing the pole-to-equator temperature gradient leads to a meridional reorientation of the Hadley

cell and the typical corresponding low latitude precipitation finger print. This meridional shift effectively changes the cross-equatorial energy transport, which is determined by the upper branch of the Hadley cell [e.g. *Schneider et al.*, 2014], towards the hemisphere that is being cooled, similar to what is found on a seasonal basis [*Donohoe et al.*, 2013]. In the simulations presented here we similarly find that a cooling (warming) of the SH mid-to-high latitudes strengthens (weakens) the southward atmospheric energy transport in the SH and weakens (strengthens) the northward energy transport in the NH (Figure 4). The net result is a change in the cross-equatorial energy transport towards the hemisphere that is cooled relative to the other hemisphere. Due to their mutual association with the Hadley cell this change in the cross-equatorial energy transport is directly related to a meridional displacement of the mean ITCZ [*Donohoe et al.*, 2013].

4. Discussion

The experiments presented here have been performed using pre-industrial boundary conditions. Possibly the response in the Southern Ocean to FWFs as well as the remote tropical response is depending on the climatic boundary conditions. For a first estimate of the applicability of these results in the broader context of past or future climate change, we performed additional experiments in which enhanced AIS meltwater input is combined with increasing greenhouse-gas concentrations. Following an RCP scenario in which the radiative forcing in the year 2100 stabilizes at $+4.5 \text{ Wm}^{-2}$ relative to the pre-industrial [RCP4.5; *Meinshausen et al.*, 2011], we analyze three simulations in which a constant AIS meltwater flux of 0 mSv, 100 mSv and 200 mSv, respectively, is applied (similar to the pre-industrial experiments). The results shown here are averaged over the years

2080-2100, thus making them somewhat different from the 200 year long pre-industrial experiments described before. Despite these differences in the experimental set-up, and, more importantly, despite the fact that these simulations combine the effects of enhanced AIS meltwater input and global warming, we see that the dependency (linear trend) of the low latitude atmospheric circulation (minimum and maximum of the Hadley cell and the location of the ITCZ) on the imposed Southern Ocean cooling is quite similar to the pre-industrial results (Supporting Information Figure 3). Notwithstanding the many differences between the pre-industrial and global warming simulations and the smaller number of global warming experiments analyzed here, the teleconnections that convey the SH high latitude temperature changes to the tropical region appear stable. Our results further suggest that the effect of Antarctic meltwater on the location of the ITCZ may be as strong as (or even stronger than) the direct radiative effect in global warming scenarios.

Several mechanisms are proposed in the literature to act as teleconnections between the SH high latitudes and the tropics. They include fast oceanic waves [*Ivchenko et al.*, 2004, 2006; *Richardson et al.*, 2005], fast atmospheric teleconnections [*Blaker et al.*, 2006] and a surface coupled wind-evaporation-SST feedback [*Ma and Wu*, 2011]. In any case, changes in tropical SST gradient are intimately linked to ITCZ displacements [*Lindzen and Nigam*, 1987; *Donohoe et al.*, 2013]. More recently it was proposed that in coupled atmosphere-ocean models changes in atmospheric cross-equatorial heat transport and corresponding ITCZ shifts are limited because the majority of the change in cross-equatorial heat transport is taken up by the ocean [*Kay et al.*, 2016; *Tomas et al.*, 2016; *Yoshimori*

et al., 2018; *Hawcroft et al.*, 2017] in an anomalous Ekman-driven subtropical-tropical overturning cell [*Green and Marshall*, 2017]. Even though our experiments show a similar response in the shallow wind-driven cell (Supporting Information Figure 4), we still find that the changes in atmospheric cross-equatorial heat transport are larger than the oceanic changes (Figure 4 and Supporting Information Figure 5). A comparison to the study by *Kay et al.* [2016], who used almost the same climate model, reveals that in our experiments the anomalous wind-driven overturning cell that transports heat across the equator [*Green and Marshall*, 2017] is shallower and hence less efficient. Moreover, in contrast to *Kay et al.* [2016], we find changes in the thermohaline-driven mid-depth and deep-ocean meridional overturning circulation (MOC) below the wind-driven cell (Supporting Information Figure 4), which modify the cross-equatorial oceanic heat transport in opposite direction to the atmospheric energy transport. We argue that the different oceanic responses can be attributed to the different ways that high latitude SH temperature changes are generated. In the work by *Kay et al.* [2016] modification of cloud physics resulted in reduced absorbed shortwave radiation leading to both surface and subsurface cooling in the SH extratropics. By contrast, imposing an AIS meltwater forcing leads to a freshening of the ocean surface which decreases the air-sea heat exchange and results in a surface cooling accompanied by a subsurface warming. This change in vertical density gradient in turn impacts deep vertical mixing, convection and hence the MOC. Subsurface warming and salinity changes possibly interact with the wind-driven changes in the shallow subtropical-tropical cell [*Green and Marshall*, 2017], explaining the much smaller changes in cross-equatorial oceanic heat transport compared to previous studies

[*Kay et al.*, 2016; *Tomas et al.*, 2016; *Yoshimori et al.*, 2018; *Hawcroft et al.*, 2017]. The response in atmospheric and oceanic cross-equatorial heat transport to different types of perturbations of the meridional surface temperature gradient is an interesting avenue for future research.

Our results suggest that the low latitude atmospheric response to a Southern Ocean temperature change is to a first order linear. Nonetheless, in line with previous work and connected to the discussion in the previous paragraph, our results also indicate that the changes in the deep ocean thermohaline circulation are more complex and dependent on the magnitude of the imposed meltwater forcing. A number of studies have shown that changing the surface salinity of the Southern Ocean impacts the formation of Antarctic Bottom Water, North Atlantic Deep Water and therewith the AMOC as a whole [*Rooth*, 1982; *Stocker et al.*, 1992; *Seidov et al.*, 2001; *Brix and Gerdes*, 2003; *Swingedouw et al.*, 2009; *Bakker et al.*, 2017]. *Swingedouw et al.* [2009] showed that the oceanic response to imposing a freshening of the Southern Ocean surface is strongly dependent on the magnitude and duration of the forcing, with different mechanisms acting at different time-scales and in opposite directions. Indeed, we find a non-linear relationship between the applied meltwater forcing and the simulated changes in the global MOC. The strength of the deep MOC cell rapidly decreases when applying increasingly large meltwater fluxes, but a saturation effect is found for very large meltwater input when the magnitude of the deep MOC cell approaches zero. On the contrary, the strength of the upper thermohaline-driven MOC cell hardly changes for small meltwater fluxes of up to 48 mSv, but starts to weaken more significantly when larger fluxes are applied (-6% and -14% for PI100mSv and

PI200mSv, respectively). This complexity in the relationship between the applied meltwater forcing and the resulting changes in the MOC partly explains the lack of a robust response in oceanic heat transport and in NH extratropical atmospheric heat transport (Figure 4 and Supporting Information Figure 5).

The impact of AIS melting on the climate of the low latitudes presented in this study first of all depends on the simulated changes in the surface characteristics of the Southern Ocean for a given FWF. The ocean model is not eddy-resolving with a horizontal resolution of $\sim 1^\circ$, nor does it resolve some of the details of the boundary currents encircling the Antarctic continent, potentially impacting the validity of the presented results. However, high resolution ocean-only simulations show that the decadal time-evolution of the spreading of the freshwater from the Antarctic continent into the Southern Ocean depends on model resolution, but the long-term equilibrium response is less affected [*Stammer, 2008; Morrison, 2015*].

5. Conclusions

A number of studies examined the impact of North Atlantic freshening and resulting cooling on the tropical atmospheric circulation and the location of the ITCZ. Here, we have used a high resolution coupled Earth system model to show that changes in Southern Ocean surface temperatures as a result of reasonable past and future AIS melt rates can also lead to substantial climatic changes in the tropics. This shows that the position of the ITCZ cannot only be changed by large-scale cooling of the NH, but similarly by temperature changes in the SH extratropics. Moreover, the induced tropical changes scale to a first order linearly with the imposed meltwater fluxes and associated SH high

latitude surface air temperature changes. This also suggests that the multi-centennial Holocene variations in AIS meltwater fluxes described by *Bakker et al.* [2017], despite being relatively small, could have an imprint in low latitude climate variability by moving the position of the ITCZ back and forth, changes that can potentially be recorded by precipitation-sensitive low latitude climate proxies [e.g. *Collins et al.*, 2013]. The first order linear relationship between the imposed AIS melt rates and the tropical climatic changes and the finding of similar results if AIS melting is imposed on top of increasing greenhouse-gas concentrations, provides confidence that investigating past episodes of climate change could help to constrain model-based projections on the role of AIS melt in driving future changes in Southern Ocean temperatures as well as in the tropical climate.

Acknowledgments. This study is a contribution to the PalMod project funded by the German Federal Ministry of Education and Science (BMBF). The climate model simulations were carried out on the supercomputer of the Norddeutscher Verbund fuer Hoch- und Hochleistungsrechnen (HLRN). For all of the described experiments, a selected number of simulated climate variables is available as netcdf-file at PANGAEA (<https://doi.pangaea.de/10.1594/PANGAEA.891414>).

References

Baker, H. S., Mbengue, C., and Woollings, T. (2018). Seasonal sensitivity of the Hadley cell and cross-hemispheric responses to diabatic heating in an idealized GCM. *Geophysical Research Letters* Online 22-02-2018

- Bakker, P., Clark, P. U., Golledge, N. R., Schmittner, A., and Weber, M. E. (2017). Centennial-scale Holocene climate variations amplified by Antarctic Ice Sheet discharge. *Nature* 541(7635), 72-76.
- Blaker, A. T., Sinha, B., Ivchenko, V. O., Wells, N. C., and Zalesny, V. B. (2006). Identifying the roles of the ocean and atmosphere in creating a rapid equatorial response to a Southern Ocean anomaly. *Geophysical Research Letters* 33, L06720.
- Brix, H., and Gerdes, R. (2003). North Atlantic Deep Water and Antarctic Bottom Water: Their interaction and influence on the variability of the global ocean circulation. *Journal of Geophysical Research* 8(2), 3022.
- Broccoli, A. J., Dahl, K. A., and Stouffer, R. J. (2006). Response of the ITCZ to Northern Hemisphere cooling. *Geophysical Research Letters* 33, L01702.
- Chiang, J. C. H., Biasutti, M. and Battisti, D. S. (2003). Sensitivity of the Atlantic intertropical convergence zone to last glacial maximum boundary conditions. *Paleoceanography* 18(4), 1094.
- Chiang, J. C. H., and Bitz, C. M. (2005). Influence of high latitude ice cover on the marine Intertropical Convergence Zone. *Climate Dynamics* 25(5), 477-496.
- Clark, P. U., Alley, R. B., Keigwin, L. D., Licciardi, J. M., Johnsen, S. J., and Wang, H. (1996). Origin of the first global meltwater pulse following the Last Glacial Maximum. *Paleoceanography and Paleoclimatology* 11(5), 563-577.
- Collins, J. A., Schefu, E., Mulitza, S., Prange, M., Werner, M., Tharammal, T., Paul, A., and Wefer, G. (2013). Estimating the hydrogen isotopic composition of past precipitation using leaf-waxes from western Africa. *Quaternary Science Review* 65, 88-101.

deConto, R., and Pollard, D. (2016). Contribution of Antarctica to past and future sea-level rise. *Nature* 531(7596), 591-597.

Donohoe, A., Marshall, J., Ferreira, D., and McGee, D. (2013). The relationship between ITCZ location and cross-equatorial atmospheric heat transport: From the seasonal cycle to the Last Glacial Maximum. *Journal of Climate* 26, 3597-3618.

Golledge, N. R., Menviel, L., Carter, L., Fogwill, C. J., England, M. H., Cortese, G., and Levy, R. H., (2014). Antarctic contribution to meltwater pulse 1A from reduced Southern Ocean overturning. *Nature Communications* 5, 1-10.

Green, B., and Marshall, J. (2017). Coupling of Trade Winds with Ocean Circulation Damps ITCZ Shifts. *Journal of Climate* 30, 4395-4411.

Hawcroft, M., Haywood, J. M., Collins, M., Jones, A., Jones, A. C., and Stephens, G. (2017). Southern Ocean albedo, inter-hemispheric energy transports and the double ITCZ: Global impacts of biases in a coupled model. *Climate Dynamics* 48 (7-8), 2279-2295.

Hwang, Y. T., and Frierson, D. M. W. (2013). Link between the double-Intertropical Convergence Zone problem and cloud biases over the Southern Ocean. *Proceedings of the National Academy of Sciences* 110(13), 4935-4940.

The IMBIE team (2018). Mass balance of the Antarctic Ice Sheet from 1992 to 2017. *Nature* 558 (7709), 219-222.

Ivchenko, V. O., Zalesny, V. B., and Drinkwater, M. R. (2004). Can the equatorial ocean quickly respond to Antarctic sea ice/salinity anomalies?. *Geophysical Research Letters* 31, L15310.

Ivchenko, V. O., Zalesny, V. B., Drinkwater, M. R. and Schröter, J. (2006). A quick response of the equatorial ocean to Antarctic sea ice/salinity anomalies. *Geophysical Research Letters* 111, C10018.

Kay, J. E., Wall, C., Yettella, V., Medeiros, B., Hannay, C., Caldwell, P., and Bitz, C. (2016). Global Climate Impacts of Fixing the Southern Ocean Shortwave Radiation Bias in the Community Earth System Model (CESM). *Journal of Climate* 29, 4617-4639.

Lindzen, R. S., and Nigam, S. (1987). On the Role of Sea Surface Temperature Gradients in Forcing Low-Level Winds and Convergence in the Tropics. *Journal of Atmospheric Sciences* 44 (17), 2418-2436.

Ma, H., Wu, L., and Li, Z. (2013). Impact of freshening over the southern ocean on ENSO. *Atmospheric Science Letters* 14, 28-33.

Ma, H., and Wu, L. (2011). Global teleconnections in response to freshening over the Antarctic Ocean. *Journal of Climate* 24, 1071-1088.

Manabe, S., and Stouffer, R. J. (1980). Sensitivity of a global climate model to an increase of CO₂ concentrations in the atmosphere. *Journal of Geophysical Research* 85, 5529-5554.

Meinshausen, M., Smith, S. J., Calvin, K., Daniel, J. S., Kainuma, M. L. T., Lamarque, J. F., Matsumoto, K., Montzka, S. A., Raper, S. C. B., Riahi, K., Thomson, A., Velders, G. J. M., and van Vuuren, D. P. P. (2011). The RCP greenhouse gas concentrations and their extensions from 1765 to 2300. *Climate Change* 109, 213-241.

Mohtadi, M., Prange, M., and Steinke, S. (2016). Palaeoclimatic insights into forcing and response of monsoon rainfall. *Nature* 533, 191-199.

Morrison, A. K. (2015). Response of Southern Ocean Convection and Abyssal Overturning to Surface Buoyancy Perturbations. *Journal of Climate* 28, 4263-4278.

Mulitza, S., Chiessi, C. M., Schefu, E., Lippold, J., Wichmann, D., Antz, B., Mackensen, A., Paul, A., Prange, M., Rehfeld, K., Werner, M., Bickert, T., Frank, N., Kuhnert, H., Lynch-Stieglitz, J., Portilho-Ramos, R. C., Sawakuchi, O. A., Schulz, M., Schwenk, T., Tiedemann, R., Vahlenkamp, M., and Zhang, Y. (2017). Synchronous and proportional deglacial changes in Atlantic meridional overturning and northeast Brazilian precipitation. *Paleoceanography* 32, 622-633.

Richardson, G., Wadley, M. R., Heywood, K. J., Stevens, D. P., and Banks, H. T. (2005). Short-term climate response to a freshwater pulse in the Southern Ocean. *Geophysical Research Letters* 31, L03702.

Rooth, C. (1982). Hydrology and ocean circulation. *Progress in Oceanography* 11, 131-149

Schneider, T., Bischoff, T. and Haug, G. (2014). Migration and dynamics of the intertropical convergence zone. *Nature* 513, 45-53.

Seidov, D., Haupt, B. J., Barron, E. J., and Maslin, M. (2001). Ocean BiPolar Seesaw and Climate: Southern Versus Northern Meltwater Impacts. *Past, present, and future*, 147-167.

Stammer, D. (2008). Response of the global ocean to Greenland and Antarctic ice melting. *Journal of Geophysical Research* 113, C06022.

Stocker, T. F., Wright, D. G., and Broecker, W. S. (1992). The influence of highlatitude surface forcing on the global thermohaline circulation. *Paleoceanography* 7(5), 529-541.

- Swingedouw, D., Fichefet, T., Goosse, H., and Loutre, M. F. (2009). Impact of transient freshwater releases in the Southern Ocean on the AMOC and climates. *Climate Dynamics* 33(2), 365-381.
- Tomas, R. A., Deser, C., and Lantao, S. (2016). The Role of Ocean Heat Transport in the Global Climate Response to Projected Arctic Sea Ice Loss. *Journal of Climate* 29, 6841-6859.
- Yoshimori, M., Abe-Ouchi, A., Tatebe, H., Nozawa, T., and Oka, A. (2018). The importance of ocean dynamical feedback for understanding the impact of mid-high latitude warming on tropical precipitation change. *Journal of Climate* 31 (6), 2417-2434.
- Zhang, R., and Delworth, T. L. (2005). Simulated tropical response to a substantial weakening of the Atlantic thermohaline circulation. *Journal of Climate* 18(12), 1853-1860.




Figure 1. Annual mean surface air temperature (left) and precipitation (right) changes as a function of the imposed FWF over all five experiments from linear regression analysis (K per 100 mSv and mm year⁻¹ per 100 mSv for the left and right panels, respectively). For both figures, results are only shown if the linear correlation is robust (i.e. significant at $\alpha=0.1$). For reference, the contours in the right-hand-side figure give the absolute precipitation values (mm year⁻¹) for the PI simulation. Note unevenly spaced colors in left-hand panel.




Figure 2. Atmospheric meridional streamfunction changes as a function of the imposed FWF over all five experiments from linear regression analysis (10^9 kg s^{-1} per 100 mSv). Results are only shown if the linear correlation is robust (i.e. significant at $\alpha=0.1$). For reference, the contours give the absolute atmospheric meridional streamfunction values (10^9 kg s^{-1}) for the PI simulation.




Figure 3. Low latitude atmospheric dynamics as a function of SH high-latitude surface air temperatures. In blue the changes in the minimum (pluses) and maximum (asterisks) meridional streamfunction (left-hand-side vertical axis; %) and in red the location of the ITCZ (right-hand-side vertical axis; °N) as a function of the SH high-latitude surface air temperatures (south of 50°S; °C). The latitude of zero meridional streamfunction at 500hPa is used as a measure of the position of the mean ITCZ.




Figure 4. Northward atmospheric heat transport anomalies. Shown are anomalies compared to the PI reference case (PW) for FWF magnitudes of -48 mSv, 48 mSv, 100 mSv and 200 mSv. Atmospheric heat transport is calculated by integrating the residual between the total top-of-the-atmosphere and surface energy fluxes.