A comprehensive quantification of global nitrous oxide sources and sinks

Accepted version

(Post-prints are subject to Springer Nature re-use terms)

Tian, H., Xu, R., Canadell, J. G., Thompson, R. L., Winiwarter, W.,
Suntharalingam, P., et al.

Published in: Nature

Reference: Tian, H., Xu, R., Canadell, J. G., Thompson, R. L., Winiwarter, W.,
Suntharalingam, P., et al. (2020). A comprehensive quantification of global nitrous oxide
sources and sinks. *Nature*, 586, 248-256. [doi:10.1038/s41586-020-2780-0](https://doi.org/10.1038/s41586-020-2780-0)

This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement No 647204

A comprehensive quantification of global nitrous oxide sources and sinks

Hanqin Tian¹, Rongting Xu¹, Josep G. Canadell², Rona L. Thompson³, Wilfried Winiwarter^{4,5}, Parvadha Suntharalingam⁶, Eric A. Davidson⁷, Philippe Ciais⁸, Robert B. Jackson^{9,10,11}, Greet Janssens-Maenhout^{12,13}, Michael J. Prather¹⁴, Pierre Regnier¹⁵, Naiqing Pan^{1,16}, Shufen Pan¹, Glen P. Peters¹⁷, Hao Shi¹, Francesco N. Tubiello¹⁸, Sönke Zaehle¹⁹, Feng Zhou²⁰, Almut Arneth²¹, Gianna Battaglia²², Sarah Berthet²³, Laurent Bopp²⁴, Alexander F. Bouwman^{25,26,27}, Erik T. Buitenhuis^{6,28}, Jinfeng Chang^{8,29}, Martyn P. Chipperfield^{30,31}, Shree R. S. Dangal³², Edward Dlugokencky³³, James W. Elkins³³, Bradley D. Eyre³⁴, Bojie Fu^{16,35}, Bradley Hall³³, Akihiko Ito³⁶, Fortunat Joos²², Paul B. Krummel³⁷, Angela Landolfi^{38,39}, Goulven G. Laruelle¹⁵, Ronny Lauerwald^{8,15,40}, Wei Li^{8,41}, Sebastian Lienert²², Taylor Maavara⁴², Michael MacLeod⁴³, Dylan B. Millet⁴⁴, Stefan Olin⁴⁵, Prabir K. Patra^{46,47}, Ronald G. Prinn⁴⁸, Peter A. Raymond⁴², Daniel J. Ruiz¹⁴, Guido R. van der Werf⁴⁹, Nicolas Vuichard⁸, Junjie Wang²⁷, Ray F. Weiss⁵⁰, Kelley C. Wells⁴⁴, Chris Wilson^{30,31}, Jia Yang⁵¹ & Yuanzhi Yao¹

¹International Center for Climate and Global Change Research, School of Forestry and Wildlife Sciences, Auburn University, Auburn, AL, USA

²Global Carbon Project, CSIRO Oceans and Atmosphere, Canberra, Australian Capital Territory, Australia

³Norsk Institutt for Luftforskning, NILU, Kjeller, Norway

⁴International Institute for Applied Systems Analysis, Laxenburg, Austria

⁵Institute of Environmental Engineering, University of Zielona Góra, Zielona Góra, Poland.

⁶School of Environmental Sciences, University of East Anglia, Norwich, UK

⁷Appalachian Laboratory, University of Maryland Center for Environmental Science, Frostburg, MD, USA

⁸Laboratoire des Sciences du Climat et de l'Environnement, LSCE, CEA CNRS, UVSQ UPSACLAY, Gif sur Yvette, France

⁹Department of Earth System Science, Stanford University, Stanford, CA, USA

¹⁰Woods Institute for the Environment, Stanford University, Stanford, CA, USA

¹¹Precourt Institute for Energy, Stanford University, Stanford, CA, USA

¹²European Commission, Joint Research Centre (JRC), Ispra, Italy

¹³Ghent University, Faculty of Engineering and Architecture, Ghent, Belgium

¹⁴Department of Earth System Science, University of California Irvine, Irvine, CA, USA

¹⁵Department Geoscience, Environment & Society, Université Libre de Bruxelles, Brussels, Belgium

¹⁶State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, China

¹⁷CICERO Center for International Climate Research, Oslo, Norway

¹⁸Statistics Division, Food and Agriculture Organization of the United Nations, Via Terme di Caracalla, Rome, Italy

¹⁹Max Planck Institute for Biogeochemistry, Jena, Germany

²⁰Sino-France Institute of Earth Systems Science, Laboratory for Earth Surface Processes, College of Urban and Environmental Sciences, Peking University, Beijing, China

²¹Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research/Atmospheric Environmental Research, Garmisch-Partenkirchen, Germany

- ²²Climate and Environmental Physics, Physics Institute and Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland
- ²³Centre National de Recherches Météorologiques (CNRM), Université de Toulouse, Météo-France, CNRS, Toulouse, France
- ²⁴LMD-IPSL, Ecole Normale Supérieure / PSL Université, CNRS; Ecole Polytechnique, Sorbonne Université, Paris, France
- ²⁵PBL Netherlands Environmental Assessment Agency, The Hague, The Netherlands
- ²⁶Department of Earth Sciences – Geochemistry, Faculty of Geosciences, Utrecht University, Utrecht, The Netherlands
- ²⁷Key Laboratory of Marine Chemistry Theory and Technology, Ministry of Education, Ocean University of China, Qingdao, China
- ²⁸Tyndall Centre for Climate Change Research, School of Environmental Sciences, University of East Anglia, Norwich, UK
- ²⁹College of Environmental and Resource Sciences, Zhejiang University, Hangzhou, China.
- ³⁰National Centre for Earth Observation, University of Leeds, Leeds, UK
- ³¹Institute for Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, Leeds, UK
- ³²Woods Hole Research Center, Falmouth, MA, USA
- ³³NOAA Global Monitoring Laboratory, Boulder, CO, USA
- ³⁴Centre for Coastal Biogeochemistry, School of Environment Science and Engineering, Southern Cross University, Lismore, New South Wales, Australia
- ³⁵Faculty of Geographical Science, Beijing Normal University, Beijing, China
- ³⁶Center for Global Environmental Research, National Institute for Environmental Studies, Tsukuba, Japan
- ³⁷Climate Science Centre, CSIRO Oceans and Atmosphere, Aspendale, Victoria, Australia
- ³⁸GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany
- ³⁹Istituto di Scienze Marine, Consiglio Nazionale delle Ricerche (CNR), Rome, Italy
- ⁴⁰Université Paris-Saclay, INRAE, AgroParisTech, UMR ECOSYS, Thiverval-Grignon, France
- ⁴¹Ministry of Education Key Laboratory for Earth System modeling, Department of Earth System Science, Tsinghua University, Beijing, China
- ⁴²Yale School of Forestry and Environmental Studies, New Haven, CT, USA
- ⁴³Land Economy, Environment & Society, Scotland's Rural College (SRUC), Edinburgh, UK
- ⁴⁴Department of Soil, Water, and Climate, University of Minnesota, St Paul, MN, USA
- ⁴⁵Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden
- ⁴⁶Research Institute for Global Change, JAMSTEC, Yokohama, Japan
- ⁴⁷Center for Environmental Remote Sensing, Chiba University, Chiba, Japan
- ⁴⁸Center for Global Change Science, Massachusetts Institute of Technology, Cambridge, MA, USA
- ⁴⁹Faculty of Science, Vrije Universiteit, Amsterdam, Netherlands.
- ⁵⁰Scripps Institution of Oceanography, University of California San Diego, La Jolla, USA
- ⁵¹Department of Forestry, Mississippi State University, Mississippi State, MS, USA

Nitrous oxide (N₂O), like carbon dioxide, is a long-lived greenhouse gas that accumulates in the atmosphere. The increase in atmospheric N₂O concentrations over the past 150 years has contributed to stratospheric ozone depletion¹ and climate change². Current national inventories do not provide a full picture of N₂O emissions owing to their omission of natural sources and the limitations in methodology for attributing anthropogenic sources. In order to understand the steadily increasing atmospheric burden (about 2 percent per decade) and develop effective mitigation strategies, it is essential to improve quantification and attribution of natural and anthropogenic contributions and their uncertainties. Here we present a global N₂O inventory that incorporates both natural and anthropogenic sources and accounts for the interaction between nitrogen additions and the biochemical processes that control N₂O emissions. We use bottom-up (inventory; statistical extrapolation of flux measurements; process-based land and ocean modelling) and top-down (atmospheric inversion) approaches to provide a comprehensive quantification of global N₂O sources and sinks resulting from 21 natural and human sectors between 1980 and 2016. Global N₂O emissions were 17.0 (minimum-maximum: 12.2–23.5) teragrams of nitrogen per year (bottom-up) and 16.9 (15.9–17.7) teragrams of nitrogen per year (top-down) between 2007 and 2016. Global human-induced emissions, which are dominated by nitrogen additions to croplands, increased by 30% over the past four decades to 7.3 (4.2–11.4) teragrams of nitrogen per year. This increase was mainly responsible for the growth in the atmospheric burden. Our findings point to growing N₂O emissions in emerging economies—particularly Brazil, China and India. Analysis of process-based model estimates reveals an emerging N₂O–climate feedback resulting from interactions between nitrogen additions and climate change. The recent growth in N₂O emissions exceeds some

of the highest projected emission scenarios^{3,4}, underscoring the urgency to mitigate N₂O emissions.

Nitrous oxide (N₂O) is a long-lived stratospheric ozone-depleting substance and greenhouse gas (GHG) with a current atmospheric lifetime of 116±9 years (ref. ¹). The concentration of atmospheric N₂O has increased by over 20% from 270 parts per billion (ppb) in 1750 to 331 ppb in 2018 (Extended Data Fig. 1), with the fastest growth observed in the past five decades^{5,6}. Two key biochemical processes, nitrification and denitrification, control N₂O production in both terrestrial and aquatic ecosystems, and are regulated by multiple environmental and biological factors, such as temperature, water, oxygen, acidity, substrate availability⁷, particularly nitrogen (N) fertilizer use and livestock manure management, and recycling⁸⁻¹⁰. In the coming decades, N₂O emissions are expected to continue increasing due to the growing demand for food, feed, fiber and energy, and a rising source from waste generation and industrial processes^{4,11,12}. Since 1990, anthropogenic N₂O emissions have been annually reported by Annex I Parties to the United Nations Framework Convention on Climate Change (UNFCCC). More recently, over 190 national signatories to the Paris Agreement are now required to report biannually their national GHG inventory with sufficient detail and transparency to track progress towards their Nationally Determined Contributions. Yet, these inventories do not provide a full picture of N₂O emissions due to their omission of natural sources, the limitations in methodology for attributing anthropogenic sources, and missing data for a number of key regions (e.g., South America, Africa)^{2,9,13}. Moreover, we need a complete account of all human activities that accelerate the global N cycle and that interact with the biochemical processes controlling the fluxes of N₂O in both terrestrial and aquatic ecosystems^{2,8}. Here we present a comprehensive, consistent analysis

and synthesis of the global N₂O budget across all sectors, including natural and anthropogenic sources and sinks, using both bottom-up (BU) and top-down (TD) methods and their cross-constraints. Our assessment enhances understanding of the global N cycle and will inform policy development for N₂O mitigation, ideally helping to curb warming to levels consistent with the long-term goal of the Paris Agreement.

A reconciling framework (described in Extended Data Fig. 2) was utilized to take full advantage of BU and TD approaches in estimating and constraining sources and sinks of N₂O. BU approaches include emission inventories, spatial extrapolation of field flux measurements, nutrient budget modeling, and process-based modeling for land and ocean fluxes. The TD approaches combine measurements of N₂O mole fractions with atmospheric transport models in statistical optimization frameworks (inversions) to constrain the sources. Here we constructed a total of 43 flux estimates including 30 with BU approaches, five with TD approaches, and eight other estimates with observation and modeling approaches (see Methods; Extended Data Fig. 2).

With this extensive data and BU/TD framework, we establish the most comprehensive global and regional N₂O budgets that include 18 sources and different versions of its chemical sink, which are further grouped into six categories (Fig. 1 and Table 1): 1) Natural sources (no anthropogenic effects) including a very small biogenic surface sink, 2) Perturbed fluxes from ecosystems induced by changes in climate, carbon dioxide (CO₂) and land cover, 3) Direct emissions of N additions in the agricultural sector (Agriculture), 4) Other direct anthropogenic sources, which include fossil fuel and industry, waste and waste water, and biomass burning, 5) Indirect emissions from ecosystems that are either downwind or downstream from the initial release of reactive N into the environment, which include N₂O release following transport and deposition of anthropogenic N via the atmosphere or water bodies as defined by the

Intergovernmental Panel on Climate Change (IPCC)¹⁴, and 6) The atmospheric chemical sink with one value derived from observations and the other (TD) from the inversion models. To quantify and attribute the regional N₂O budget, we further partition the Earth's ice-free land into ten regions (Fig. 2 and Supplementary Fig. 1). With the construction of these budgets, we explore the relative temporal and spatial importance of multiple sources and sinks driving the atmospheric burden of N₂O, their uncertainties, and interactions between anthropogenic forcing and natural fluxes of N₂O as an emerging climate feedback.

The Global N₂O Budget (2007–2016)

The BU and TD approaches give consistent estimates of global total N₂O emissions in the recent decade to well within their respective uncertainties, with values of 17.0 (min-max: 12.2–23.5) Tg N yr⁻¹ and 16.9 (15.9–17.7) Tg N yr⁻¹ for BU and TD sources, respectively. The global calculated atmospheric chemical sink (i.e., N₂O losses via photolysis and reaction with O(¹D) in the troposphere and stratosphere) is 13.5 (12.4–14.6) Tg N yr⁻¹. The imbalance of sources and sinks of N₂O derived from the averaged BU and TD estimates is 4.1 Tg N yr⁻¹. This imbalance agrees well with the observed 2007–2016 increase in atmospheric abundance of 3.8–4.8 Tg N yr⁻¹ (see Methods). Natural sources from soils and oceans contributed 57% of total emissions (mean: 9.7; min-max: 8.0–12.0 Tg N yr⁻¹) for the recent decade according to our BU estimate. We further estimate the natural soil flux at 5.6 (4.9–6.5) Tg N yr⁻¹ and the ocean flux at 3.4 (2.5–4.3) Tg N yr⁻¹ (see Methods).

Anthropogenic sources contributed on average 43% to the total N₂O emission (mean: 7.3; min-max: 4.2–11.4 Tg N yr⁻¹), in which direct and indirect emissions from N additions in agriculture and other sectors contributed ~52% and ~18%, respectively. Of the remaining

anthropogenic emissions, ~27% were from other direct anthropogenic sources including fossil fuel and industry (~13%), with ~3% from perturbed fluxes caused by climate/CO₂/land cover change.

Four Decades of the Global N₂O Budget

The atmospheric N₂O burden increased from 1462 Tg N in the 1980s to 1555 Tg N in the recent decade, with a possible uncertainty ± 20 Tg N. Our results (Table 1) demonstrate that global N₂O emissions have also significantly increased, primarily driven by anthropogenic sources, with natural sources relatively steady throughout the study period. Our BU and TD global N₂O emissions are comparable in magnitude during 1998–2016, but TD results imply a larger inter-annual variability (1.0 Tg N yr⁻¹; Extended Data Fig. 3a). BU and TD approaches diverge in the magnitude of land versus ocean emissions, although they are consistent with respect to trends. Specifically, the BU land estimate during 1998–2016 was on average 1.8 Tg N yr⁻¹ higher than the TD estimate, but showed a slightly slower increasing rate of 0.8 ± 0.2 Tg N yr⁻¹ per decade (95% confidence interval; $P < 0.05$) compared to 1.1 ± 0.6 Tg N yr⁻¹ per decade ($P < 0.05$) from TD (Extended Data Fig. 3b). Since 2005, the difference in the magnitude of emissions between the two approaches has become smaller due to a large TD-inferred emission increase, particularly in South America, Africa, and East Asia (Extended Data Fig. 3d, f, i). Oceanic N₂O emissions from BU [3.6 (2.7–4.5) Tg N yr⁻¹] indicate a slight decline at a rate of 0.06 Tg N yr⁻¹ per decade ($P < 0.05$), while the TD approach gave a higher but stable value of 5.1 (3.4–7.1) Tg N yr⁻¹ during 1998–2016 (Table 1).

Based on BU approaches, anthropogenic N₂O emissions increased from 5.6 (3.6–8.7) Tg N yr⁻¹ in the 1980s to 7.3 (4.2–11.4) Tg N yr⁻¹ in the recent decade at a rate of 0.6 ± 0.2 Tg N yr⁻¹ per

decade ($P < 0.05$). Up to 87% of this increase is from direct emission from agriculture (71%) and indirect emission from anthropogenic N additions into soils (16%). Direct soil emission from fertilizer applications is the major source for agricultural emission increases, followed by a small but significant increase in emissions from livestock manure and aquaculture. The model-based estimates of direct soil emissions¹⁵⁻¹⁷ exhibit a faster increase than the three inventories used in our study (see Methods; Extended Data Fig. 4a), which is largely attributed to the interactive effects between climate change and N additions as well as spatio-temporal variability in environmental factors such as rainfall and temperature that modulate the N₂O yield from nitrification and denitrification. This result is in line with the elevated emission factor (EF) deduced from the TD estimates, in which the inversion-based soil emissions increased at a faster rate than suggested by the IPCC Tier 1 EF¹⁴ (which assumes a linear response), especially after 2009 (ref. ¹⁸). The remaining causes of the increase are attributed to other direct anthropogenic sources (6%) and perturbed fluxes from climate/CO₂/land cover change (8%). The part of fossil fuel and industry emissions decreased rapidly over 1980–2000 largely due to the installation of emissions abatement equipment in industrial facilities producing nitric and adipic acid. However, after 2000 such emissions began to increase slowly due to rising fossil fuel combustion (Extended Data Fig. 5a-b).

Our analysis of process-based model estimates indicates that soil N₂O emissions accelerated substantially due to climate change since the early 1980s, which has offset the reduction due to elevated CO₂ concentration (Extended Data Fig. 6a). Elevated CO₂ enhances plant growth and thus increases N uptake, which in turn decreases soil N₂O emissions^{16,19}. Land conversion from tropical mature forests with higher N₂O emissions to pastures and other unfertilized agricultural lands has significantly reduced global natural N₂O emissions^{11,20,21}. This decrease, however, was

partly offset by an increase in soil N₂O emissions attributable to the temporary rise of emissions following deforestation (post-deforestation pulse effect) and background emissions from converted croplands or pastures²¹ (see Methods; Extended Data Fig. 7).

From the ensemble of process-based land model emissions^{15,16}, we estimate a global agricultural soil EF of 1.8% (1.3%–2.3%), which is significantly larger than the IPCC Tier-1 default for direct emission of 1%. This higher EF, derived from process-based models, suggests a strong interactive effect between N additions and other global environmental changes (Table 1, Perturbed fluxes from climate, atmospheric CO₂, and land cover change). Previous field experiments reported a better fit to local observations of soil N₂O emissions when assuming a non-linear response to fertilizer N inputs under varied climate and soil conditions^{17,22}. The non-linear response is likely also associated with long-term N accumulation in agricultural soils from N fertilizer use and in aquatic systems from N loads (the legacy effect)^{18,23}, which provides more substrate for microbial processes^{18,24}. The increasing N₂O emissions estimated by process-based models¹⁶ also suggest that recent climate change (particularly warming) may have boosted soil nitrification and denitrification processes, contributing to the growing trend in N₂O emissions together with rising N additions to agricultural soils^{16,25-27} (Extended Data Fig. 8).

Regional N₂O Budgets (2007–2016)

BU approaches give estimates of N₂O emissions in the five source categories, while TD approaches only provide total emissions (Fig. 2). BU and TD approaches indicate that Africa was the largest N₂O source in the last decade, followed by South America (Fig. 2). BU and TD approaches agree well in the magnitudes and trends of N₂O emissions from South Asia and Oceania (Extended Data Fig. 3j, l). For the remaining regions, BU and TD estimates are

comparable in their trends but diverge in their source strengths. Clearly, much more work on regional N₂O budgets is needed, particularly for South America and Africa where we see larger differences between BU and TD estimates and larger uncertainty in each approach. Advancing the understanding and model representation of key processes responsible for N₂O emissions from land and ocean are priorities for reducing uncertainties in BU estimates. Atmospheric observations in underrepresented regions of the world and better atmospheric transport models are essential for uncertainty reduction in TD estimates, while more accurate activity data and robust EFs are critical for GHG inventories (See Methods for additional discussion on uncertainty).

Based on the Global N₂O Model Intercomparison Project (NMIP) estimates¹⁶, natural soil emissions (to different extents) dominated in tropical and sub-tropical regions. Soil N₂O emissions in the tropics ($0.1 \pm 0.04 \text{ g N m}^{-2} \text{ yr}^{-1}$) are about 50% higher than the global average, since many lowland, highly-weathered tropical soils have excess N relative to phosphorus²⁰. Total anthropogenic emissions in the ten terrestrial regions were highest in East Asia (1.5; 0.8–2.6 Tg N yr⁻¹), followed by North America, Africa, and Europe. High direct agricultural N₂O emissions can be attributed to large-scale synthetic N fertilizer applications in East Asia, Europe, South Asia, and North America, which together consume over 80% of the world's synthetic N fertilizers²⁸. In contrast, direct agricultural emissions from Africa and South America are mainly induced by livestock manure that is deposited in pastures and rangelands^{28,29}. East Asia contributed 71%–79% of global aquaculture N₂O emissions; South Asia and Southeast Asia together contributed 10%–20% (refs. ^{30,31}). Indirect emissions play a moderate role in the total N₂O budget, with the highest emission in East Asia (0.3; 0.1–0.5 Tg N yr⁻¹). Other direct

anthropogenic sources together contribute N₂O emissions of approximately 0.2–0.4 Tg N yr⁻¹ in East Asia, Africa, North America, and Europe.

Both BU and TD estimates of ocean N₂O emissions for northern, tropical, and southern ocean regions (90°–30°N, 30°N–30°S, and 30°–90°S, respectively) reveal that the tropical oceans contribute over 50% to the global oceanic source. In particular, the upwelling regions of the equatorial Pacific, Indian and tropical Atlantic (Fig. 3) provide significant sources of N₂O³²⁻³⁴. BU estimates suggest the southern ocean is the second largest regional contributor with emissions about twice as high as from the northern oceans (53% tropical oceans, 31% southern oceans, 17% northern oceans), in line with their area, while the TD estimates suggest approximately equal contributions from the southern and northern oceans.

Four Decades of Anthropogenic N₂O Emissions

Trends in anthropogenic emissions varied among regions (Fig. 3). Fluxes from Europe and Russia decreased by a total of 0.6 (0.5–0.7) Tg N yr⁻¹ over the past 37 years (1980–2016). The decrease in Europe is associated with successful emissions abatement in industry as well as agricultural policies, while the decrease in Russia is associated with the collapse of the agricultural cooperative system after 1990. In contrast, fluxes from the remaining eight regions increased by a total of 2.9 (2.4–3.4) Tg N yr⁻¹ (Fig. 3), of which 34% came from East Asia, 18% from Africa, 18% from South Asia, 13% from South America, only 6% from North America, and with remaining increases due to other regions.

The relative importance of each anthropogenic source to the total emission increase differs among regions. East Asia, South Asia, Africa, and South America show larger increases in total agricultural N₂O emissions (direct and indirect) compared to the remaining six regions during

1980–2016 (Fig. 3). Southeast Asia, North America, and Middle East also show increasing direct N₂O emissions but to smaller extent. Rising indirect emissions in these four regions (East Asia, South Asia, Africa, and South America) on average constitute 20% of total agricultural N₂O emissions and are largely induced by the considerable increase in fertilizer N inputs to agricultural soils^{35,36}. The most rapid increase in emissions from other direct anthropogenic sources was found in East Asia, primarily owing to the fast-growing industrial emissions. Africa and South Asia show a fast emission increase due to emissions from fossil fuel and industry and waste and waste water.

Our findings point to growing N₂O emissions in emerging economies, particularly Brazil, China, and India. For example, we find here that the substantial increases in livestock manure left on pasture and in fertilizer use caused a ~120% increase in Brazilian agricultural N₂O emissions during 1980–2016 (Extended Data Fig. 9). In addition to fertilizer applications, global livestock manure production has been growing steadily, in line with increased livestock numbers^{15,28}. Rising demand for meat and dairy products has significantly increased global N₂O emissions from livestock manure production and management associated with the expansion of pastures and grazing land³⁷. Meanwhile, expansion of feed crop production to support the growth of livestock could further enhance global N₂O emissions^{37,38}. Likewise, increasing demand for fish has triggered a five-fold increase in global aquaculture production since the late 1980s³⁹, with demand projected to increase further⁴⁰, although this remains a small fraction (<1%) of total N₂O emissions.

The acceleration of global N₂O emissions resulting from anthropogenic sources is apparent in both BU and TD results and currently tracks the highest Representative Concentration Pathway (RCP8.5)⁴ in the fifth assessment report (AR5) of IPCC² and exceeds all the Shared

Socioeconomic Pathways (SSPs)³ in CMIP6 for the sixth assessment report (AR6) of IPCC (Fig. 4). Observed atmospheric N₂O concentrations are beginning to exceed predicted levels across all scenarios. Emissions need to be reduced to a level that is consistent with or below that in RCP2.6 or SSP1-2.6 in order to limit warming well below the 2° C target of the Paris Agreement. Failure to include N₂O within climate mitigation strategies will necessitate even greater abatement of CO₂ and CH₄. Although N₂O mitigation is difficult because N is the key-limiting nutrient in the agricultural production, this study demonstrates that effective mitigation actions have reduced emissions in some regions, such as Europe, through technological improvements in industry and improved N use efficiency in agriculture.

There are a number of mitigation options in the agriculture sector available for immediate deployment, including increased N use efficiency in (i) animal production through tuning of feed rations to reduce N excretion, and (ii) in crop production through precision delivery of N fertilizers, split applications and better timing to match N applications to crop demand, conservation tillage, prevention of waterlogging, and the use of nitrification inhibitors^{43,44}. Success stories include the stabilization or reduction of N₂O emissions through improving N use efficiency in the United States and Europe, while maintaining or even increasing crop yields^{44,45}. There is every reason to expect that additional implementation of more sustainable practices and emerging technologies will lead to further reductions in these regions. For example, N₂O emissions from European agricultural soils decreased by 21% between 1990 and 2010, a decline attributable to the implementation of the Nitrates Directive (an agricultural policy favoring optimization and reduction of fertilizer use as well as water protection legislation)⁴⁶. For regions where emissions are growing, an immediate opportunity lies in the reduction of excess fertilizer use along with the implementation of more sustainable agricultural practices that together have

been shown to increase crop yields, reduce N₂O emissions, increase water quality, and increase farm income⁴⁷. In addition, N₂O emissions can be efficiently abated in the chemical industry^{11,43,48,49}, as has been achieved successfully in nitric acid plants in the European Union where industrial N₂O emissions dropped from 11% to 3% of total emissions between 2007 and 2012 (ref. ⁴⁶). Additional available strategies to reduce N₂O emissions include promoting lower meat consumption in some parts of the world⁹ and reducing food waste¹¹.

We present the most comprehensive global N₂O budget to date, with a detailed sectorial and regional attribution of sources and sinks. Each of the past four decades had higher global N₂O emissions than the previous one, and in all, agricultural activities dominated the growth in emissions. Total industrial emissions have been quite stable with increased emissions from the fossil fuel sector offset to some extent by the decline in emissions in other industrial sectors as a result of successful abatement policies. We also highlight a number of complex interactions between N₂O fluxes and human-driven changes whose impact on the global atmospheric N₂O growth rate was previously unknown. Those interactions include the effects of climate change, increasing atmospheric CO₂, and deforestation. Cumulatively, these exert a relatively small effect on the overall N₂O growth, however, individual flux components, such as the growing positive climate-N₂O feedback, are significant. These fluxes are not currently included in the national GHG reporting. We further find that Brazil, China, and India dominate the regional contributions to the increase in global N₂O emissions over the most recent decade. Our extensive database and modelling capability fill current gaps in national and regional emissions inventories. Future research is needed to further constrain complex biogeochemical interactions between natural/anthropogenic fluxes and global environmental changes, which could lead to significant feedbacks in the future. Reducing excess N applications to croplands and adopting

precision fertilizer application methods provide the largest immediate opportunities for N₂O emissions abatement.

References

- 1 Prather, M. J. *et al.* Measuring and modeling the lifetime of nitrous oxide including its variability. *Journal of Geophysical Research: Atmospheres* **120**, 5693-5705 (2015).
- 2 Ciais, P. *et al.* in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* 465-570 (Cambridge University Press, 2014).
- 3 Gidden, M. J. *et al.* Global emissions pathways under different socioeconomic scenarios for use in CMIP6: a dataset of harmonized emissions trajectories through the end of the century. *Geoscientific Model Development* **12**, 1443-1475 (2019).
- 4 Davidson, E. A. Representative concentration pathways and mitigation scenarios for nitrous oxide. *Environmental Research Letters* **7**, 024005 (2012).
- 5 Hall, B., Dutton, G. & Elkins, J. The NOAA nitrous oxide standard scale for atmospheric observations. *Journal of Geophysical Research: Atmospheres* **112**, D09305 (2007).
- 6 Prinn, R. G. *et al.* History of chemically and radiatively important atmospheric gases from the Advanced Global Atmospheric Gases Experiment (AGAGE). *Earth System Science Data* **10**, 985-1018 (2018).
- 7 Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R. & Zechmeister-Boltenstern, S. Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Phil. Trans. R. Soc. B* **368**, 20130122 (2013).
- 8 Tian, H. *et al.* The terrestrial biosphere as a net source of greenhouse gases to the atmosphere. *Nature* **531**, 225-228 (2016).
- 9 UNEP. Drawing down N₂O to protect climate and the ozone layer. Report No. 9280733583, (United Nations Environment Programme (UNEP), 2013).
- 10 Park, S. *et al.* Trends and seasonal cycles in the isotopic composition of nitrous oxide since 1940. *Nature Geoscience* **5**, 261-265 (2012).
- 11 Davidson, E. A. & Kanter, D. Inventories and scenarios of nitrous oxide emissions. *Environmental Research Letters* **9**, 105012 (2014).
- 12 Reay, D. S. *et al.* Global agriculture and nitrous oxide emissions. *Nature Climate Change* **2**, 410-416 (2012).
- 13 Syakila, A. & Kroeze, C. The global nitrous oxide budget revisited. *Greenhouse Gas Measurement and Management* **1**, 17-26 (2011).
- 14 IPCC. 2006 IPCC Guidelines for National Greenhouse Gas Inventories., (Japan on behalf of the IPCC, Hayama, Japan, 2006).
- 15 Dangal, S. R. *et al.* Global nitrous oxide emissions from pasturelands and rangelands: Magnitude, spatio-temporal patterns and attribution. *Global Biogeochemical Cycles* **33**, 200-222 (2019).
- 16 Tian, H. Q. *et al.* Global soil nitrous oxide emissions since the preindustrial era estimated by an ensemble of terrestrial biosphere models: Magnitude, attribution, and uncertainty. *Global Change Biology* **25**, 640-659 (2019).

- 17 Wang, Q. *et al.* Data-driven estimates of global nitrous oxide emissions from croplands. *National Science Review* **7**, 441-452 (2020).
- 18 Thompson, R. L. *et al.* Acceleration of global N₂O emissions seen from two decades of atmospheric inversion. *Natural Climate Change* **9**, 993-998 (2019).
- 19 Zaehle, S., Ciais, P., Friend, A. D. & Prieur, V. Carbon benefits of anthropogenic reactive nitrogen offset by nitrous oxide emissions. *Nature Geoscience* **4**, 601-605 (2011).
- 20 Davidson, E. A. *et al.* Recuperation of nitrogen cycling in Amazonian forests following agricultural abandonment. *Nature* **447**, 995-998 (2007).
- 21 Verchot, L. V. *et al.* Land use change and biogeochemical controls of nitrogen oxide emissions from soils in eastern Amazonia. *Global Biogeochemical Cycles* **13**, 31-46 (1999).
- 22 Shcherbak, I., Millar, N. & Robertson, G. P. Global metaanalysis of the nonlinear response of soil nitrous oxide (N₂O) emissions to fertilizer nitrogen. *Proceedings of the National Academy of Sciences* **111**, 9199-9204 (2014).
- 23 Van Meter, K. J., Basu, N. B., Veenstra, J. J. & Burras, C. L. The nitrogen legacy: emerging evidence of nitrogen accumulation in anthropogenic landscapes. *Environmental Research Letters* **11**, 035014 (2016).
- 24 Firestone, M. K. & Davidson, E. A. Microbiological basis of NO and N₂O production and consumption in soil. *Exchange of trace gases between terrestrial ecosystems the atmosphere* **47**, 7-21 (1989).
- 25 Griffis, T. J. *et al.* Nitrous oxide emissions are enhanced in a warmer and wetter world. *Proceedings of the National Academy of Sciences* **114**, 12081-12085 (2017).
- 26 Pärn, J. *et al.* Nitrogen-rich organic soils under warm well-drained conditions are global nitrous oxide emission hotspots. *Nature Communications* **9**, 1135 (2018).
- 27 Smith, K. The potential for feedback effects induced by global warming on emissions of nitrous oxide by soils. *Global Change Biology* **3**, 327-338 (1997).
- 28 FAOSTAT. The Food and Agriculture Organization of the United Nations Statistics: Emissions-Agriculture, Emissions Land Use Trade (Crops and livestock products), Population, Agri-Environmental Indicators (Livestock Manure) (2019).
- 29 Xu, R. *et al.* Increased nitrogen enrichment and shifted patterns in the world's grassland: 1860–2016. *Earth System Science Data* **11**, 175-187 (2019).
- 30 Beusen, A. H., Bouwman, A. F., Van Beek, L. P., Mogollón, J. M. & Middelburg, J. J. Global riverine N and P transport to ocean increased during the 20th century despite increased retention along the aquatic continuum. *Biogeosciences* **13**, 2441-2451 (2016).
- 31 MacLeod, M., Hasan, M. R., Robb, D. H. F. & Mamun-Ur-Rashid, M. Quantifying and mitigating greenhouse gas emissions from global aquaculture. *FAO, Rome* (2019).
- 32 Buitenhuis, E. T., Suntharalingam, P. & Le Quéré, C. Constraints on global oceanic emissions of N₂O from observations and models. *Biogeosciences* **15**, 2161-2175 (2018).
- 33 Manizza, M., Keeling, R. F. & Nevison, C. D. On the processes controlling the seasonal cycles of the air–sea fluxes of O₂ and N₂O: A modelling study. *Tellus B: Chemical and Physical Meteorology* **64**, 18429 (2012).
- 34 Martinez-Rey, J., Bopp, L., Gehlen, M., Tagliabue, A. & Gruber, N. Projections of oceanic N₂O emissions in the 21st century using the IPSL Earth system model. *Biogeosciences* **12**, 4133-4148 (2015).
- 35 Maavara, T. *et al.* Nitrous oxide emissions from inland waters: Are IPCC estimates too high? *Global Change Biology* **25**, 473-488 (2019).

- 36 Yao, Y. *et al.* Increased global nitrous oxide emissions from streams and rivers in the Anthropocene. *Natural Climate Change* **10**, 138-142 (2020).
- 37 Gerber, P. J. *et al.* Tackling climate change through livestock: a global assessment of emissions and mitigation opportunities. *FAO* (2013).
- 38 Herrero, M. *et al.* Biomass use, production, feed efficiencies, and greenhouse gas emissions from global livestock systems. *Proceedings of the National Academy of Sciences* **110**, 20888-20893 (2013).
- 39 Yuan, J. *et al.* Rapid growth in greenhouse gas emissions from the adoption of industrial-scale aquaculture. *Nature Climate Change* **9**, 318-322 (2019).
- 40 Froehlich, H. E., Runge, C. A., Gentry, R. R., Gaines, S. D. & Halpern, B. S. Comparative terrestrial feed and land use of an aquaculture-dominant world. *Proceedings of the National Academy of Sciences* **115**, 5295-5300 (2018).
- 41 O'Neill, B. C. *et al.* The Scenario Model Intercomparison Project (ScenarioMIP) for CMIP6. *Geoscience Model Development* **9**, 3461-3482 (2016).
- 42 Gütschow, J. *et al.* The PRIMAP-hist national historical emissions time series. *Earth System Science Data* **8**, 571-603 (2016).
- 43 Winiwarter, W., Höglund-Isaksson, L., Klimont, Z., Schöpp, W. & Amann, M. Technical opportunities to reduce global anthropogenic emissions of nitrous oxide. *Environmental Research Letters* **13**, 014011 (2018).
- 44 Zhang, X. *et al.* Managing nitrogen for sustainable development. *Nature* **528**, 51-59 (2015).
- 45 Mueller, N. D. *et al.* Declining spatial efficiency of global cropland nitrogen allocation. *Global Biogeochemical Cycles* **31**, 245-257 (2017).
- 46 European Environment Agency. Annual European Union greenhouse gas inventory 1990-2017 and inventory report 2019. *Submission under the United Nations Framework Convention on Climate Change and the Kyoto Protocol, Copenhagen, DK* (2019).
- 47 Cui, Z. *et al.* Pursuing sustainable productivity with millions of smallholder farmers. *Nature* **555**, 363-366 (2018).
- 48 Kanter, D. *et al.* A post-Kyoto partner: considering the stratospheric ozone regime as a tool to manage nitrous oxide. *Proceedings of the National Academy of Sciences* **110**, 4451-4457 (2013).
- 49 Schneider, L., Lazarus, M. & Kollmuss, A. J. S. M. S. E. I. Industrial N₂O projects under the CDM: Adipic acid-A case of carbon leakage. *Stockholm Environment Institute* (2010).

496 **Table 1 The global N₂O budget in the 1980s, 1990s, 2000s, and 2007–2016.**

		the 1980s			the 1990s			the 2000s			2007-2016		
Anthropogenic sources		mean	min	max	mean	min	max	mean	min	max	mean	min	max
Direct emissions of N additions in the agricultural sector (Agriculture)	Direct soil emissions	1.5	0.9	2.6	1.7	1.1	3.1	2.0	1.3	3.4	2.3	1.4	3.8
	Manure left on pasture	0.9	0.7	1.0	1.0	0.7	1.1	1.1	0.8	1.2	1.2	0.9	1.3
	Manure management	0.3	0.2	0.4	0.3	0.2	0.4	0.3	0.2	0.5	0.3	0.2	0.5
	Aquaculture	0.01	0.00	0.03	0.03	0.01	0.1	0.1	0.02	0.2	0.1	0.02	0.2
	sub-total	2.6	1.8	4.1	3.0	2.1	4.8	3.4	2.3	5.2	3.8	2.5	5.8
Other direct anthropogenic sources	Fossil fuel and industry	0.9	0.8	1.1	0.9	0.9	1.0	0.9	0.8	1.0	1.0	0.8	1.1
	Waste and waste water	0.2	0.1	0.3	0.3	0.2	0.4	0.3	0.2	0.4	0.3	0.2	0.5
	Biomass burning	0.7	0.7	0.7	0.7	0.6	0.8	0.6	0.6	0.6	0.6	0.5	0.8
	sub-total	1.8	1.6	2.1	1.9	1.7	2.1	1.8	1.6	2.1	1.9	1.6	2.3
Indirect emissions from anthropogenic N additions	Inland waters, estuaries, coastal zones	0.4	0.2	0.5	0.4	0.2	0.5	0.4	0.2	0.6	0.5	0.2	0.7
	Atmospheric N deposition on land	0.6	0.3	1.2	0.7	0.4	1.4	0.7	0.4	1.3	0.8	0.4	1.4
	Atmospheric N deposition on ocean	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.2	0.1	0.1	0.2
	sub-total	1.1	0.6	1.9	1.2	0.7	2.1	1.2	0.6	2.1	1.3	0.7	2.2
Perturbed fluxes from climate/CO ₂ /land cover change	CO ₂ effect	-0.2	-0.3	0.0	-0.2	-0.4	0.0	-0.3	-0.5	0.1	-0.3	-0.6	0.1
	Climate effect	0.4	0.0	0.8	0.5	0.1	0.9	0.7	0.3	1.2	0.8	0.3	1.3
	Post-deforestation pulse effect	0.7	0.6	0.8	0.7	0.6	0.8	0.7	0.7	0.8	0.8	0.7	0.8
	Long-term effect of reduced mature forest area	-0.8	-0.8	-0.9	-0.9	-0.8	-1.0	-1.0	-0.9	-1.1	-1.1	-1.0	-1.1
	sub-total	0.1	-0.4	0.7	0.1	-0.5	0.7	0.2	-0.4	0.9	0.2	-0.6	1.1
Anthropogenic total		5.6	3.6	8.7	6.2	3.9	9.7	6.7	4.1	10.3	7.3	4.2	11.4
Natural fluxes													
Natural soils baseline		5.6	4.9	6.6	5.6	4.9	6.5	5.6	5.0	6.5	5.6	4.9	6.5
Ocean baseline		3.6	3.0	4.4	3.5	2.8	4.4	3.5	2.7	4.3	3.4	2.5	4.3
Natural (Inland waters, estuaries, coastal zones)		0.3	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4	0.3	0.3	0.4
Lightning and atmospheric production		0.4	0.2	1.2	0.4	0.2	1.2	0.4	0.2	1.2	0.4	0.2	1.2
Surface sink		-0.01	0.00	-0.3	-0.01	0.00	-0.3	-0.01	0.00	-0.3	-0.01	0.00	-0.3
Natural total		9.9	8.5	12.2	9.8	8.3	12.1	9.8	8.2	12.0	9.7	8.0	12.0
Bottom-up total source		15.5	12.1	20.9	15.9	12.2	21.7	16.4	12.3	22.4	17.0	12.2	23.5
Top-down Ocean								5.1	3.1	7.2	5.1	3.4	7.1
Top-down Land								10.8	9.3	12.5	11.8	10.6	13.8
Top-down total source								15.9	15.1	16.9	16.9	15.9	17.7
Top-down Statospheric sink								12.1	11.4	13.1	12.4	11.7	13.3
Observed atmospheric chemical sink*								13.3	12.2	14.4	13.5	12.4	14.6
Change in atmospheric abundance**								3.7	3.2	4.2	4.3	3.8	4.8
Atmospheric burden		1462	1442	1482	1493	1472	1514	1531	1510	1552	1555	1533	1577

497 Note: BU estimates include four categories of anthropogenic sources (red for agriculture, orange for
498 other direct anthropogenic sources, maroon for indirect emissions from anthropogenic N additions, and
499 brown for perturbed fluxes from climate/CO₂/land cover change) and one category for natural sources
500 and sinks (green). The sources and sinks of N₂O are given in Tg N yr⁻¹. The atmospheric burden is given
501 in Tg N. *calculated from satellite observations with a photolysis model (about 1% of this sink
502 occurs in the troposphere). **Calculated from the combined NOAA and AGAGE record of surface N₂O,
503 and adopting the uncertainty of the IPCC AR5 (Chapter 6)². Detailed information on calculating each
504 sub-category is shown in Supplementary Tables 1–13.

Fig. 1 Global N₂O budget for the recent decade (2007–2016). The red arrow represents direct emissions of N additions in the agricultural sector (Agriculture). The orange arrows represent emissions from other direct anthropogenic sources. The maroon arrows represent indirect emissions from anthropogenic N additions. The brown arrows represent perturbed fluxes from climate/CO₂/land cover change effects. The green arrows represent natural source. The anthropogenic and natural N₂O sources are derived from BU estimates. The blue arrows represent surface sink and observed atmospheric chemical sink of which about 1% occurs in the troposphere. The total budget (sources + sinks) does not exactly match the observed atmospheric accumulation, because each of the terms has been derived independently and we do not force top-down agreement by rescaling the terms. This imbalance readily falls within the overall uncertainty in closing the N₂O budget, as reflected in each of the terms. The N₂O sources and sinks are given in Tg N yr⁻¹.

Fig. 2 Regional N₂O sources in the recent decade (2007–2016) over 11 regions. The Earth's ice-free land is partitioned into ten regions: North America (NA), South America (SA), Europe (EU), Middle East (MIDE), Africa (AF), Russia (RUS), East Asia (EAS), South Asia (SAS), Southeast Asia (SEA), and Oceania (OCE). In each subplot from left to right: emissions from five sub-sectors using BU approaches: natural fluxes without ocean (green), direct emissions of N additions in the agricultural sector (Agriculture, red), other direct anthropogenic sources (orange), indirect emissions from anthropogenic N additions (maroon), and perturbed fluxes from climate/CO₂/land cover change (brown); the sum of these five categories by BU approaches (blue), and the estimates by TD approaches (gold). BU and TD estimates of ocean emissions are shown at the bottom left (from bottom to top: 30°–90°N, 30°S–30°N, and 90°–30°S). Error bars indicate the spread between the minimum and the maximum values. The center map shows the spatial distribution of 10-year average N₂O emissions from land and ocean based on the land and ocean models. Per capita N₂O emission (kg N capita⁻¹ yr⁻¹) during 2007–2016 is shown in Supplementary Fig. 2.

Fig. 3 Ensembles of regional anthropogenic N_2O emissions over the 1980–2016 period. The bar chart in the center shows the accumulated changes in regional and global N_2O emissions during the study period. Error bars indicate the 95% confidence interval for the average of accumulated changes. The Mann-Kendall test was performed to examine a monotonic increasing or decreasing trend in the estimated ensemble N_2O emissions for each region and the globe during 1980–2016. The accumulated changes were calculated from the linear regressed annual change rate (Tg N yr^{-2}) multiplied by 37 years. All regions except SEA show a significant increasing or decreasing trend in the estimated ensemble N_2O emissions during the study period (indicated by ** for each bar).

Fig. 4 Historical and projected global anthropogenic N₂O emissions and concentrations.

Global anthropogenic N₂O emissions (**a**, **b**) and concentrations (**c**, **d**) compared to the four representative concentration pathways (RCPs) in the IPCC AR5 (**a**, **c**, ref. ²) and the new marker scenarios based on the Shared Socioeconomic Pathways (SSPs) used in CMIP6 (**b**, **d**, ref. ⁴¹).

The historical data is represented as the mean of the BU and TD estimates of anthropogenic N₂O emissions, while the atmospheric concentration uses the three observation networks available, AGAGE, NOAA, and CSIRO. TD anthropogenic emissions were calculated by subtracting BU-derived natural fluxes. To aid the comparison, the four RCPs were shifted down so that the 2005 value is equal to the 2000–2009 average of the mean of TD and BU estimates. The SSPs are harmonized³ to match the historical emissions used in CMIP6⁴² and Extended Data Fig. 10 shows the unharmonized data.

Methods

Terminology. This study provides an estimation of the global N₂O budget considering all possible sources and all global change processes that can perturb the budget. A total of 18 sources and three sinks of N₂O are identified and grouped into six categories (Figure 1, Table 1):

- 1) Natural fluxes in absence of climate change and anthropogenic disturbances including Soil emissions, Surface sink, Ocean emissions, Lightning and atmospheric production, and Natural emission from inland waters, estuaries, coastal zones (inland and coastal waters),
- 2) Perturbed fluxes from climate/CO₂/land cover change including CO₂ effect, Climate effect, Post-deforestation pulse effect, and Long-term effect of reduced mature forest area,
- 3) Direct emissions of N additions in the agricultural sector (Agriculture) including emissions from direct application of synthetic N fertilizers and manure (henceforth Direct soil emissions), Manure left on pasture, Manure management, and Aquaculture,
- 4) Indirect emissions from anthropogenic N additions including atmospheric N deposition (NDEP) on land, atmospheric NDEP on ocean, and effects of anthropogenic loads of reactive N in inland waters, estuaries, coastal zones,
- 5) Other direct anthropogenic sources including Fossil fuel and industry, Waste and waste water, and Biomass burning, and
- 6) Two estimates of stratospheric sinks obtained from atmospheric chemistry transport models and observations, and one tropospheric sink (Table 1, Extended Data Fig. 2).

For the purpose of compiling national GHG inventories for country reporting to the climate convention, our anthropogenic N₂O emission categories are aligned with those used in UNFCCC reporting and IPCC 2006 methodologies (Supplementary Table 14). We also provide the detailed comparison of our methodology and quantification with the IPCC AR5 (see Supplementary Section 4; Supplementary Table 15).

Data synthesis. We consider global N₂O emission from land and ocean consisting of natural fluxes and anthropogenic emissions based on BU and TD approaches, however, the TD approach cannot separate natural and anthropogenic sources.

‘Natural soil baseline’ emissions were obtained from six terrestrial biosphere models (NMIP¹⁶, Supplementary Tables 16–17) and provided here reflect a situation without consideration of land use change (e.g., deforestation) and without consideration of indirect anthropogenic effects via global change (i.e., climate, elevated CO₂, and atmospheric N deposition). BU oceanic N₂O emissions were based on an inter-comparison of five global ocean biogeochemistry models (Supplementary Table 18). The natural emission from ‘Inland water, estuaries, coastal zones’ includes coastal upwelling⁵⁰ and inland and coastal waters that were obtained from Yao et al.³⁶, Maavara et al.³⁵, and Lauerwald et al.⁵¹. Since the data (rivers, reservoirs, and estuaries) provided by Maavara et al. and Lauerwald et al. are for the year 2000, we assume that these values are constant during 1980–2016. Yao et al.³⁶ provided annual riverine N₂O emissions using DLEM during the same period. Here, we averaged estimates from Yao et al. with that from Maavara et al.³⁵. In addition, we estimated N₂O emissions from global and regional reservoirs in the 2000s, and averaged their estimates with that from Maavara et al.³⁵ to represent emissions from reservoirs during 1980–2016. The estimate for global and regional estuaries and lakes is still based on the long-term averaged values provided by Maavara et al.³⁵ and Lauerwald et al.⁵¹, respectively. We considered the riverine emissions in the year 1900 as equivalent to the natural emission for the DLEM estimate assuming that the N load from land was negligible in that period⁵². We quantified the contribution of natural sources to total emission from reservoirs, lakes, and estuaries at 44% (36%–52%), with consideration of all N inputs (i.e., inorganic, organic, dissolved, particulate forms). We combined the estimate from

lightning with that from atmospheric production into an integrated category ‘Lightning and atmospheric production’. We make the simplification of considering the category ‘Lightning and atmospheric production’ as purely natural, however, atmospheric production is affected to some extent by anthropogenic activities through enhancing the concentrations of the reactive species NH_2 and NO_2 . This category is in any case very small and the anthropogenic enhancement effect is uncertain. Lightning produces NO_x , the median estimate of which is 5 Tg N yr^{-1} (ref. ⁵³). We assumed an EF of 1% (ref. ⁵⁴) and a global estimate of $0.05 (0.02\text{--}0.09) \text{ Tg N yr}^{-1}$ from lightning. Atmospheric production of N_2O results from the reaction of NH_2 with NO_2 (refs. ^{55,56}), N with NO_2 , and oxidation of N_2 by $\text{O}(^1\text{D})$ ⁵⁷, all of which constitute an estimated source of $0.3 (0.2\text{--}1.1) \text{ Tg N yr}^{-1}$. The estimate of ‘Surface sink’ was obtained from Schlesinger⁵⁸ and Syakila et al.⁵⁹.

The anthropogenic sources include four sub-sectors:

(a) Agriculture. It consists of four components: ‘Direct soil emissions’, ‘Manure left on pasture’, ‘Manure management’, and ‘Aquaculture’. Data for ‘Direct soil emissions’ were obtained as the ensemble mean of N_2O emissions from an average of three inventories (EDGAR v4.3.2, FAOSTAT, and GAINS), the SRNM/DLEM models, and the NMIP/DLEM models. The statistical model SRNM only covers cropland N_2O emissions, the same as the NMIP. Thus, we add the DLEM-based estimate of pasture N_2O emissions into the two estimates in cropland to represent direct agricultural soil emissions (i.e., SRNM/DLEM or NMIP/DLEM). The ‘Manure left on pasture’ and ‘Manure management’ emissions are the ensemble mean of EDGAR v4.3.2, FAOSTAT, and GAINS databases. Global N flows (i.e., fish feed intake, fish harvest, and waste) in freshwater and marine aquaculture were obtained from Beusen et al.³⁰ and Bouwman et al.^{60,61} based on a nutrient budget model for the period 1980–2016. We then calculated global aquaculture N_2O emissions through considering 1.8% loss of N waste in aquaculture, the same

EF used in Hu et al.⁶² and Macleod et al.³¹. The uncertainty range of the EF is from 0.5% (ref. ¹⁴) to 5% (ref. ⁶³), the same range used in the UNEP report⁹. The ‘Aquaculture’ emission for the period 2007–2016 was a synthesis data from Hu et al.⁶² in 2009, the FAO Report³¹ in 2013, and our calculations. The estimate of aquaculture N₂O emission prior to 2009 was from our calculations only.

The estimated direct emissions from agriculture have increased from 2.6 (1.8–4.1) Tg N yr⁻¹ in the 1980s to 3.8 (2.5–5.8) Tg N yr⁻¹ over the recent decade (2007–2016, Table 1). Specifically, direct soil emission from the application of fertilizers is the major source and increased at a rate of 0.27±0.01 Tg N yr⁻¹ per decade (P < 0.05; Table 1). Compared with the three global inventories (FAOSTAT, EDGAR v4.3.2, and GAINS), the estimates from process-based models (NMIP/DLEM^{15,16}) and a statistical model (SRNM)/DLEM^{15,17} exhibited a faster increase (Extended Data Fig. 4a). Over the past four decades, we also found a small but significant increase in emissions from livestock manure (i.e., manure left on pasture and manure management) at a rate of 0.1±0.01 Tg N yr⁻¹ per decade (P < 0.05; Extended Data Fig. 4b-c). Meanwhile, global aquaculture N₂O emissions increased 10-fold, however, this flux remains the smallest term in the global budget (Extended Data Fig. 4d).

(b) Other direct anthropogenic sources. It includes ‘Fossil fuel and industry’, ‘Waste and waste water’, and ‘Biomass burning’. Both ‘Fossil fuel and industry’ and ‘Waste and waste water’ are the ensemble means of EDGAR v4.3.2 and GAINS databases. The ‘Biomass burning’ emission is the ensemble mean of FAOSTAT, DLEM, and GFED4s databases.

Emissions from a combination of fossil fuel and industry, waste and waste water, and biomass burning increased from 1.8 (1.6–2.1) Tg N yr⁻¹ in the 1980s to 1.9 (1.6–2.3) Tg N yr⁻¹ over the period of 2007–2016 (Table 1). The waste and waste water emission showed a continuous

increase at a rate of $0.04 \pm 0.01 \text{ Tg N yr}^{-1}$ per decade ($P < 0.05$) (Extended Data Fig. 5c). Emissions from biomass burning, estimated based on three data sources (DLEM, GFED4s, and FAOSTAT), slightly decreased at a rate of $-0.03 \pm 0.04 \text{ Tg N yr}^{-1}$ per decade ($P = 0.3$) since the 1980s (Extended Data Fig. 5d). This item is largely affected by climate and land use change^{64,65}. Of the three data sources, the DLEM estimate exhibited significant inter-annual variability, especially during 1980–2000 when extreme fire events were detected in 1982, 1987, 1991, 1994, and 1998. The occurrences of these extreme fires were associated with El Niño-Southern Oscillation (ENSO) events, especially in Indonesia (e.g., ‘Great Fire of Borneo’ in 1982)⁶⁶. Since 1997, N_2O emissions from fires estimated by DLEM, GFED4s, and FAOSTAT were consistent in the inter-annual variability. All the three estimates showed a decreasing trend, agreeing well with satellite-observed decrease of global burned area^{64,65}.

(c) Indirect emissions from anthropogenic N additions. Data were obtained from various sources and considered N deposition on land and ocean (‘N deposition on land’ and ‘N deposition on ocean’), as well as the N leaching and runoff from upstream (‘Inland and coastal waters’). The emission from ‘N deposition on ocean’ was provided by Suntharalingam et al.⁶⁷, while emission from ‘N deposition on land’ was the ensemble mean of an average of three inventories: FAOSTAT/EDGAR v4.3.2, GAINS/EDGAR v4.3.2, and NMIP. FAOSTAT and GAINS documented the sector ‘Indirect agricultural N_2O emissions’ by separating estimates from N leaching or N deposition, while EDGAR v4.3.2 did not. Here, we treated ‘Indirect agricultural N_2O emissions’ from EDGAR v4.3.2 as ‘Inland and coastal waters’ emissions for data synthesis. Only EDGAR v4.3.2 provided an estimate of indirect emission from non-agricultural sectors, while both FAOSTAT and GAINS, following the IPCC guidelines, provided NH_3/NO_y volatilization from agricultural sectors. Here, we sum FAOSTAT or GAINS with

EDGAR v4.3.2 (i.e., FAOSTAT/EDGAR v4.3.2 or GAINS/EDGAR v4.3.2) to represent N deposition induced soil emissions from both agricultural and non-agricultural sectors. The N₂O emissions from ‘Inland and coastal waters’ consist of rivers, reservoirs, lakes, estuaries, and coastal zone, which is the ensemble mean of an average of three inventories (EDGAR v4.3.2, FAOSTAT, GAINS), and the mean of process-based models. The anthropogenic emission estimated by Yao et al.³⁶ considered annual N inputs and other environmental factors (i.e., climate, elevated CO₂, and land cover change). For long-term average in rivers, reservoirs, estuaries and lakes, we applied a mean of 56% (based on the ratio of anthropogenic to total N additions from land) to calculate anthropogenic emissions. Seagrass, mangrove, saltmarsh and intertidal N₂O emissions were undated from Murray et al⁶⁸. Coastal waters with low disturbance generally either have low N₂O emissions or act as a sink for N₂O^{69,70}. Here, coastal zone emissions were treated as anthropogenic emissions due to intensive human disturbances⁷¹.

N₂O emissions following transport of anthropogenic N additions via atmosphere and water bodies increased from 1.1 (0.6–1.9) Tg N yr⁻¹ in the 1980s to 1.3 (0.7–2.2) Tg N yr⁻¹ during 2007–2016 (Table 1). The N₂O emissions from inland and coastal waters increased at a rate of 0.03±0.00 Tg N yr⁻¹ per decade (P < 0.05). Such an increase was reported by all the three inventories (FAOSTAT, GAINS, and EDGAR v4.3.2) with FAOSTAT giving the largest estimate. In contrast, the DLEM-based estimate presented a divergent trend: first increasing from 1980–1998 and then slightly decreasing thereafter (Extended Data Fig. 6a). Emissions from atmospheric N deposition on oceans were relatively constant with a value of 0.1 (0.1–0.2) Tg N yr⁻¹, while a large increase in emissions was found from atmospheric N deposition on land, with 0.06±0.01 Tg N yr⁻¹ per decade (P < 0.05) reported in the three estimates (FAOSTAT/EDGAR v4.3.2, GAINS/EDGAR v4.3.2, and NMIP). The FAOSTAT agricultural source, together with

the EDGAR v4.3.2 industrial source, is consistent with NMIP estimates in the magnitude of N₂O emissions, with the latter estimating a slightly slower increase from 2010 to 2016 (Extended Data Fig. 6b).

(d) Perturbed fluxes from climate/CO₂/land cover change. Perturbed N₂O fluxes represent the sum of the effects of climate, elevated atmospheric CO₂, and land cover change. The estimate of climate and CO₂ effects on emissions was based on NMIP. The effect of land cover change on N₂O dynamics includes the reduction due to ‘Long-term effect of reduced mature forest area’ and the emissions due to ‘Post-deforestation pulse effect’. The two estimates were based on the book-keeping approach and the DLEM model simulation. The book-keeping method is developed by Houghton et al.⁷² for accounting for carbon flows due to land use. In this study, an observation dataset consisting of 18 tropical sites was collected to follow the book-keeping logic. The dataset covers N₂O emissions from a reference mature forest and their nearby converted pastures aged between one and 60 years. The average tropical forest N₂O emission rate of 1.974 kg N₂O-N ha⁻¹ yr⁻¹ was adopted as the baseline⁷³. Two logarithmic response curves of soil N₂O emissions (normalized to the baseline) after deforestation were developed: $y = -0.31 \ln(x) + 1.53$ ($R^2 = 0.30$) and $y = -0.454 \ln(x) + 2.21$ ($R^2 = 0.09$). The first logarithmic function uses data collected by a review analysis⁷⁴, based upon which the second one further considers observations from Verchot et al.²¹ and Keller and Reiners⁷⁵. In the first function, x (unit: year) indicates pasture age in years after deforestation and y (unitless; 0–1) indicates the ratio of pasture N₂O emission over the N₂O emission from the nearby reference mature forest. In the second function, x (unit: year) indicates secondary forest age and y (unitless; 0–1) indicates the ratio of secondary forest N₂O emission over that of a reference mature forest. This form of the response functions can effectively reproduce the short-lived increase in soil N₂O emissions after

initial forest clearing and the gradually declining emission rates of converted crops/pastures^{21,76}. Using these two curves and the baseline, we kept track of the N₂O reduction of tropical forests and the post-deforestation crop/pasture N₂O emissions at an annual time-scale. This book-keeping method was applied to the two deforestation area datasets (Supplementary Text 2.8), so we could investigate not only the difference caused by the two sets of land use data but also the difference between this empirical method and the process-based model. For land conversion from natural vegetation to croplands or pastures, DLEM uses a similar strategy to Houghton et al.⁷² and McGuire et al.⁷⁷ to simulate its influences on carbon and N cycles. Moreover, through using the sites of field observation from Davidson et al.²⁰ and Keller and Reiners⁷⁵, we estimated N₂O emission from secondary tropical forests based on the algorithm: $y = 0.0084x + 0.2401$ ($R^2 = 0.44$). x (unit: year) indicates secondary forest age and y (unitless; 0–1) indicates the ratio of secondary forest N₂O emission over that of a reference mature forest. The difference between primary forests and secondary forests were subtracted from natural soil emissions simulated by six terrestrial biosphere models in NMIP.

We calculated the ensemble of oceanic N₂O emission based on the BU approach (five ocean biogeochemical models; Supplementary Table 18) and the TD approach (five estimates from four inversion models; Supplementary Table 19), respectively. The atmospheric burden and its rate of change during 1980–2016 were derived from mean maritime surface mixing ratios of N₂O (refs. ^{78,79}) with a conversion factor of 4.79 Tg N/ppb (ref. ⁸⁰). Combining uncertainties in measuring the mean surface mixing ratios⁷⁸ and that of converting surface mixing ratios to a global mean abundance⁸⁰, we estimate a $\pm 1.4\%$ uncertainty in the burden. Annual change in atmospheric abundance is calculated from the combined NOAA and AGAGE record of surface N₂O and uncertainty is taken from the IPCC AR5 (ref. ²). There shows an agreement of the

stratospheric loss from atmospheric chemistry transport models (TD modeled chemical sink^{18,81}) and from satellite observations with a photolysis model (observed photochemical sink¹), which differ only by ~ 1 Tg N yr⁻¹. The satellite-based lifetime, 116 ± 9 years, gives an overall uncertainty in the annual loss of $\pm 8\%$. The tropospheric loss of N₂O from reaction with O(¹D) is included in observed atmospheric chemical sink (Table 1) and is small ($\sim 1\%$ of the stratospheric sink) with an estimated range of 0.1 to 0.2 Tg N yr⁻¹.

Comparison with the IPCC guidelines. The IPCC has provided guidance to quantify N₂O emissions, which is widely used in emission inventories for reporting to the UNFCCC. Over time the recommended approaches have changed, which is critical for estimating emissions from agricultural soils, the largest emission source. Previous global N₂O assessments^{52,82,83} based on the IPCC 1996 guidelines⁸⁴ attributed about 6.3 Tg N yr⁻¹ to the agricultural sector, including both direct and indirect emissions. This estimate is significantly larger than our results (Fig. 1; Table 1) derived from multiple methods, and is also larger than the most recent estimates from global inventories (EDGAR v4.3.2, FAOSTAT, and GAINS) that are based on the IPCC 2006 guidelines¹⁴. The main reason is that indirect emissions from leaching and groundwater were overestimated in previous studies⁸⁵. Correspondingly, projections of atmospheric N₂O concentrations based on these overestimated emissions⁸² led to biased estimates. For example, Mosier and Kroeze⁸² expected atmospheric N₂O concentrations to be 340–350 ppb in the year 2020, instead of 333 ppb⁵ as observed. Recently, the 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories has been published. It adopts the same approach for N application on soils, but considers impacts of different climate regimes. The new guidelines, based on a wealth of new scientific literature, proposed much smaller emissions from grazing animals by a factor of 5–7. Preliminary calculations we have made indicate that global

soil emissions based on these new guidelines may decrease by 20%–25%. Integrating estimates relying on the IPCC methodology with estimates by process-based models provides for a more balanced assessment in this paper. We also added information from assessments^{86,87} that derived agricultural emissions as the difference between atmospheric terms and other emissions like combustion, industry and nature, and they gave comparable magnitudes (4.3–5.8 Tg N yr⁻¹) to our bottom-up results.

Uncertainty. Current data analysis and synthesis of long-term N₂O fluxes are based on a wide variety of TD and BU methods. TD approaches, consisting of four inversion frameworks⁸⁸⁻⁹¹, provide a wide range of estimates largely due to systematic errors in the modelled atmospheric transport and stratospheric loss of N₂O. In addition, the emissions from TD analyses are dependent on the magnitude and distribution of the prior flux estimates to an extent that is strongly determined by the number of atmospheric N₂O measurements¹⁸. Inversions are generally not well constrained (and thus rely heavily on a priori estimates) in Africa, Southeast Asia, southern South America, and over the oceans, owing to the paucity of observations in these regions. The improvement of atmospheric transport models, more accurate priors, and more atmospheric N₂O measurements would reduce uncertainty in further TD estimates, particularly for ocean and regional emissions.

BU approaches are subject to uncertainties in various sources from land¹⁶ and oceans³². For process-based models (e.g. NMIP and ocean biogeochemical models), the uncertainty is associated with differences in model configuration as well as process parameterization^{16,32}. The uncertainty of estimates from NMIP could be reduced in multiple ways¹⁶. First, the six models in NMIP exhibited different spatial and temporal patterns of N₂O emissions even though they used the same forcings. Although these models have considered essential biogeochemical processes in

soils (e.g., biological N fixation, nitrification/denitrification, mineralization/immobilization, etc.)⁹², some missing processes such as freeze-thaw cycles and ecosystem disturbances should be included in terrestrial biosphere models to reduce uncertainties. Second, the quality of input datasets, specifically the amount and timing of N application, and spatial and temporal changes in distribution of natural vegetation and agricultural land, is critical for accurately simulating soil N₂O emissions. Third, national and global N₂O flux measurement networks¹⁷ could be used to validate model performance and constrain large-scale model simulations. Data assimilation techniques could be utilized to improve model accuracy.

Current remaining uncertainty in global ocean model estimates of N₂O emission includes the contribution of N₂O flux derived from the tropical oceanic low oxygen zones (e.g., the Eastern Equatorial Pacific, the northern Indian ocean) relative to the global ocean. These low oxygen zones are predominantly influenced by high yield N₂O formation processes (e.g., denitrification and enhanced nitrification). Regional observation-based assessments have also suggested that these regions may produce more N₂O than is simulated by the models³². The current generation of global ocean biogeochemistry models are not sufficiently accurate to represent the high N₂O production processes in low-oxygen zones, and their associated variability (see refs. ^{34,93,94} for more detail). Thus, precisely representing the local ocean circulation and associated biogeochemical fluxes of these regions could further reduce the uncertainty in estimates of global and regional oceanic N₂O emissions.

Regardless of the tier approach used, GHG inventories for agriculture suffer from high uncertainty in the underlying agriculture and rural data and statistics used as input, including statistics on fertilizer use, livestock manure availability, storage and applications, and nutrient, crop and soils management. For instance, animal waste management is an uncertain aspect, since

much of the manure is either not used, or employed as a fuel or building material, or may be discharged directly to surface water^{95,96}, with important repercussions for the calculated emissions. Furthermore, GHG inventories using default EFs show large uncertainties at local to global scales, especially for agricultural N₂O emissions, due to the poorly captured dependence of EFs on spatial diversity in climate, management, and soil physical and biochemical conditions^{2,22}. It is well known, for example from the IPCC guidelines, that higher-tier GHG inventories may provide more reasonable estimates by using the alternative EFs that are disaggregated by environmental factors and management-related factors⁹⁷. A large range of EFs have been used to estimate aquaculture N₂O emissions^{31,39,62,86} and long-term estimates of N flows in freshwater and marine aquaculture are scarce³⁰. Uncertainty also remains in several N₂O sources that have not yet been fully understood or quantified. To date, robust estimates of N₂O emissions from global peatland degradation are still lacking, although we have accounted for N₂O emissions due to the drainage of organic soils (histosols) obtained from FAOSTAT and GAINS databases^{28,43}. Recent evidence shows that permafrost thawing⁹⁸ and the freeze-thaw cycle⁹⁹ contribute to increasing N₂O emissions, which, however, have not been well established in the current estimates of the global N₂O budget.

Statistics. Through using the Mann-Kendall test in R-3.4.4, we checked the significance of trends in annual N₂O emissions from each sub-sector based on the BU approach.

References

- 50 Nevison, C. D., Lueker, T. J. & Weiss, R. F. Quantifying the nitrous oxide source from coastal upwelling. *Global Biogeochemical Cycles* **18**, GB1018 (2004).
- 51 Lauerwald, R. *et al.* Natural lakes are a minor global source of N₂O to the atmosphere. *Global Biogeochemical Cycles* **33**, 1564-1581 (2019).
- 52 Kroeze, C., Mosier, A. & Bouwman, L. Closing the global N₂O budget: a retrospective analysis 1500–1994. *Global Biogeochemical Cycles* **13**, 1-8 (1999).

839 53 Schumann, U. & Huntrieser, H. The global lightning-induced nitrogen oxides source.
840 *Atmospheric Chemistry and Physics* **7**, 3823-3907 (2007).

841 54 De Klein, C. *et al.* N₂O emissions from managed soils, and CO₂ emissions from lime and
842 urea application. *IPCC Guidelines for National Greenhouse Gas Inventories, Prepared*
843 *by the National Greenhouse Gas Inventories Programme* **4**, 1-54 (2006).

844 55 Dentener, F. J. & Crutzen, P. J. A three-dimensional model of the global ammonia cycle.
845 *Journal of Atmospheric Chemistry* **19**, 331-369 (1994).

846 56 Röckmann, T., Kaiser, J., Crowley, J. N., Brenninkmeijer, C. A. & Crutzen, P. J. The
847 origin of the anomalous or "mass-independent" oxygen isotope fractionation in
848 tropospheric N₂O. *Geophysical Research Letters* **28**, 503-506 (2001).

849 57 Kaiser, J. & Röckmann, T. Absence of isotope exchange in the reaction of N₂O+O(¹D)
850 and the global Δ¹⁷O budget of nitrous oxide. *Geophysical Research Letters* **32** (2005).

851 58 Schlesinger, W. H. An estimate of the global sink for nitrous oxide in soils. *Global*
852 *Change Biology* **19**, 2929-2931 (2013).

853 59 Syakila, A., Kroeze, C. & Slomp, C. P. Neglecting sinks for N₂O at the earth's surface:
854 does it matter? *Journal of Integrative Environmental Sciences* **7**, 79-87 (2010).

855 60 Bouwman, A. F. *et al.* Hindcasts and future projections of global inland and coastal
856 nitrogen and phosphorus loads due to finfish aquaculture. *Reviews in Fisheries Science*
857 **21**, 112-156 (2013).

858 61 Bouwman, A. F. *et al.* Global hindcasts and future projections of coastal nitrogen and
859 phosphorus loads due to shellfish and seaweed aquaculture. *Reviews in Fisheries Science*
860 **19**, 331-357 (2011).

861 62 Hu, Z., Lee, J. W., Chandran, K., Kim, S. & Khanal, S. K. Nitrous oxide (N₂O) emission
862 from aquaculture: a review. *Environmental science technology* **46**, 6470-6480 (2012).

863 63 Williams, J. & Crutzen, P. J. Nitrous oxide from aquaculture. *Nature Geoscience* **3**, 143
864 (2010).

865 64 Andela, N. *et al.* A human-driven decline in global burned area. *Science* **356**, 1356-1362
866 (2017).

867 65 Yang, J. *et al.* Spatial and temporal patterns of global burned area in response to
868 anthropogenic and environmental factors: Reconstructing global fire history for the 20th
869 and early 21st centuries. *Journal of Geophysical Research: Biogeosciences* **119**, 249-263
870 (2014).

871 66 Dennis, R. A review of fire projects in Indonesia (1982-1998). *Cifor* (1999).

872 67 Suntharalingam, P. *et al.* Quantifying the impact of anthropogenic nitrogen deposition on
873 oceanic nitrous oxide. *Geophysical Research Letters* **39**, L07605 (2012).

874 68 Murray, R. H., Erler, D. V. & Eyre, B. D. Nitrous oxide fluxes in estuarine environments:
875 response to global change. *Global Change Biology* **21**, 3219-3245 (2015).

876 69 Erler, D. V. *et al.* Applying cavity ring - down spectroscopy for the measurement of
877 dissolved nitrous oxide concentrations and bulk nitrogen isotopic composition in aquatic
878 systems: Correcting for interferences and field application. *Limnology and*
879 *Oceanography: Methods* **13**, 391-401 (2015).

880 70 Murray, R., Erler, D., Rosentreter, J. & Eyre, B. Seasonal and spatial N₂O concentrations
881 and emissions in three tropical estuaries. *Marine Chemistry* **221**, 103779 (2020).

882 71 Vernberg, F. J. & Vernberg, W. B. *The coastal zone: past, present, and future*. Univ of
883 South Carolina Press (2001).

884 72 Houghton, R. *et al.* Changes in the Carbon Content of Terrestrial Biota and Soils between
885 1860 and 1980: A Net Release of CO₂ to the Atmosphere. *Ecological monographs* **53**,
886 235-262 (1983).

887 73 Davidson, E. A. The contribution of manure and fertilizer nitrogen to atmospheric nitrous
888 oxide since 1860. *Nature Geoscience* **2**, 659-662 (2009).

889 74 van Lent, J., Hergoualc'h, K. & Verchot, L. V. Reviews and syntheses: Soil N₂O and NO
890 emissions from land use and land-use change in the tropics and subtropics: a meta-
891 analysis. *Biogeosciences* **12**, 7299-7313 (2015).

892 75 Keller, M. & Reiners, W. A. Soil-atmosphere exchange of nitrous oxide, nitric oxide, and
893 methane under secondary succession of pasture to forest in the Atlantic lowlands of Costa
894 Rica. *Global Biogeochemical Cycles* **8**, 399-409 (1994).

895 76 Melillo, J. M. *et al.* Nitrous oxide emissions from forests and pastures of various ages in
896 the Brazilian Amazon. *Journal of Geophysical Research: Atmospheres* **106**, 34179-34188
897 (2001).

898 77 McGuire, A. *et al.* Carbon balance of the terrestrial biosphere in the twentieth century:
899 Analyses of CO₂, climate and land use effects with four process-based ecosystem models.
900 *Global Biogeochemical Cycles* **15**, 183-206 (2001).

901 78 Dlugokencky, E., Steele, L., Lang, P. & Masarie, K. The growth rate and distribution of
902 atmospheric methane. *Journal of Geophysical Research: Atmospheres* **99**, 17021-17043
903 (1994).

904 79 Prather, M. *et al.* Annex II: Climate system scenario tables. *Cambridge, United Kingdom*
905 *and New York, NY, USA* (2013).

906 80 Prather, M. J., Holmes, C. D. & Hsu, J. Reactive greenhouse gas scenarios: Systematic
907 exploration of uncertainties and the role of atmospheric chemistry. *Geophysical Research*
908 *Letters* **39**, L09803 (2012).

909 81 Prather, M. J. & Hsu, J. Coupling of Nitrous Oxide and Methane by Global Atmospheric
910 Chemistry. *Science* **330**, 952-954 (2010).

911 82 Mosier, A. & Kroeze, C. Potential impact on the global atmospheric N₂O budget of the
912 increased nitrogen input required to meet future global food demands. *Chemosphere-*
913 *Global Change Science* **2**, 465-473 (2000).

914 83 Mosier, A. *et al.* Closing the global N₂O budget: nitrous oxide emissions through the
915 agricultural nitrogen cycle. *Nutrient cycling in Agroecosystems* **52**, 225-248 (1998).

916 84 IPCC. Revised 1996 IPCC guidelines for national greenhouse gas inventories. *Hayama,*
917 *Japan* (1997).

918 85 Nevison, C. in IPCC, Background Papers: IPCC Expert Meetings on Good Practice
919 Guidance and Uncertainty Management in National Greenhouse Gas Inventories. *IPCC*
920 *National Greenhouse Gas Inventories Programme, Technical Support Unit*. 381-397
921 (2000).

922 86 Crutzen, P. J., Mosier, A. R., Smith, K. A. & Winiwarter, W. N₂O release from agro-
923 biofuel production negates global warming reduction by replacing fossil fuels.
924 *Atmospheric Chemistry and Physics* **8**, 389-395 (2008).

925 87 Smith, K. A., Mosier, A. R., Crutzen, P. J. & Winiwarter, W. The role of N₂O derived
926 from crop-based biofuels, and from agriculture in general, in Earth's climate.
927 *Philosophical Transactions of the Royal Society of London B: Biological Sciences* **367**,
928 1169-1174 (2012).

- 88 Thompson, R. L. *et al.* Nitrous oxide emissions 1999 to 2009 from a global atmospheric inversion. *Atmospheric Chemistry and Physics* **14**, 1801-1817 (2014).
- 89 Wells, K. C. *et al.* Simulation of atmospheric N₂O with GEOS-Chem and its adjoint: evaluation of observational constraints. *Geoscience Model Development* **8**, 3179-3198 (2015).
- 90 Wilson, C., Chipperfield, M., Gloor, M. & Chevallier, F. Development of a variational flux inversion system (INVICAT v1. 0) using the TOMCAT chemical transport model. *Geoscientific Model Development* **7**, 2485-2500 (2014).
- 91 Patra, P. K. *et al.* Improved Chemical Tracer Simulation by MIROC4. 0-based Atmospheric Chemistry-Transport Model (MIROC4-ACTM). *Sola* **14**, 91-96 (2018).
- 92 Tian, H. Q. *et al.* The Global N₂O Model Intercomparison Project. *Bulletin of the American Meteorological Society* **99**, 1231-1252 (2018).
- 93 Suntharalingam, P. *et al.* Anthropogenic nitrogen inputs and impacts on oceanic N₂O fluxes in the northern Indian Ocean: The need for an integrated observation and modelling approach. *Deep Sea Research Part II: Topical Studies in Oceanography* **166**, 104-113 (2019).
- 94 Battaglia, G. & Joos, F. Marine N₂O Emissions From Nitrification and Denitrification Constrained by Modern Observations and Projected in Multimillennial Global Warming Simulations. *Global Biogeochemical Cycles* **32**, 92-121 (2018).
- 95 Galloway, J. *et al.* The impact of animal production systems on the nitrogen cycle. Vol. 1, *Island Press* (2010).
- 96 Steinfeld, H., Mooney, H. A., Schneider, F. & Neville, L. E. Livestock in a changing landscape, volume 1: drivers, consequences, and responses. Vol. 1, *Island Press* (2013).
- 97 IPCC. 2019 Refinement to the 2006 IPCC Guidelines for National Greenhouse Gas Inventories. *Hayama, Japan* (2019).
- 98 Elberling, B., Christiansen, H. H. & Hansen, B. U. High nitrous oxide production from thawing permafrost. *Nature Geoscience* **3**, 332-335 (2010).
- 99 Wagner-Riddle, C. *et al.* Globally important nitrous oxide emissions from croplands induced by freeze-thaw cycles. *Nature Geoscience* **10**, 279-283 (2017).
- 100 Suntharalingam, P. *et al.* Estimates of Oceanic Nitrous-oxide Emissions from Global Biogeochemistry Models. *American Geophysical Union, Fall Meeting 2018* (2018).
- 101 Janssens-Maenhout, G. *et al.* EDGAR v4.3.2 Global Atlas of the three major greenhouse gas emissions for the period 1970–2012. *Earth System Science Data* **11**, 959-1002 (2019).
- 102 Tubiello, F. *et al.* Estimating greenhouse gas emissions in agriculture: a manual to address data requirements for developing countries. *FAO, Rome* (2015).
- 103 Van Der Werf, G. R. *et al.* Global fire emissions estimates during 1997-2016. *Earth System Science Data* **9**, 697-720 (2017).
- 104 Dentener, F. Global maps of atmospheric nitrogen deposition, 1860, 1993, and 2050. *Data set. Available on-line (<http://daac.ornl.gov/>) from Oak Ridge National Laboratory Distributed Active Archive Center, Oak Ridge, TN, USA* (2006).
- 105 Riahi, K. *et al.* The Shared Socioeconomic Pathways and their energy, land use, and greenhouse gas emissions implications: An overview. *Global Environmental Change* **42**, 153-168 (2017).

Data availability

The relevant datasets of this study are archived in the box site of International Center for Climate and Global Change Research at Auburn University (<https://auburn.box.com/>). Source data for Figs. 1–4, Table 1, Extended Figs. 1–10 and Supplementary Information are provided with the paper. Additional description on data availability for atmospheric N₂O observations from NOAA, AGAGE and CSIRO networks is provided in the Supplementary Information. The data presented here are made available in the belief that their dissemination will lead to greater understanding and new scientific insights on the global and regional N₂O budgets and changes to it, and helping to reduce the uncertainties. As data are the result of initial processing to fit to the purpose of this publication, typically a wealth of underlying information is with the original data providers. Researchers interested to use results made available in the repository are encouraged, as good practice, to take advantage of underlying information by contacting the original data providers. If such a contact develops into a more intensive scientific discussion, further involvement including co-authorship should be considered.

Code availability

The relevant codes of this study are archived in the box site of International Center for Climate and Global Change Research at Auburn University (<https://auburn.box.com/>).

Acknowledgements

This paper is the result of a collaborative international effort under the umbrella of the Global Carbon Project (a project of Future Earth and a research partner of the World Climate Research Programme) and International Nitrogen Initiative. This research was made possible partly by Andrew Carnegie Fellowship Award no. G-F-19-56910; NSF grant nos. 1903722, 1243232 and 1922687; NASA grant nos. NNX14AO73G, NNX10AU06G, NNX11AD47G and NNX14AF93G; NOAA grant nos. NA16NOS4780207 and NA16NOS4780204; National Key R & D Program of China (grant nos. 2017YFA0604702 and 2018YFA0606001), National Natural Science Foundation of China (Grant no. 41961124006), CAS grants (KFJ-STSZDTP-0; SKLURE2017-1-6), and OUC-AU Joint Center Program. Additional funding support includes: E.T.B., P.R., G.P.P., R.L.T., P.S. acknowledge funding support from VERIFY project (EC H2020 grant no. 776810). P.S. also acknowledges funding from the EC H2020 grant No 641816 (CRESCENDO); A.I. acknowledges funding support from JSPS KAKENHI grant (no. 17H01867). G.B., F.J., and S.L. acknowledge support by Swiss National Science Foundation (#200020_172476) and by the EC H2020 grant no. 821003 (Project 4C) and no. 820989 (Project COMFORT). A.L. acknowledges support by DFG project SFB754/3. S.Z. acknowledges support by the EC H2020 grant no. 647204. K.C.W. and D.B.M. acknowledge support from NASA (IDS Grant #NNX17AK18G) and NOAA (Grant #NA13OAR4310086); P.A.R. acknowledges NASA Award NNX17AI74G, M.M. acknowledges support from the Scottish Government's Rural and Environment Science and Analytical Services Division (RESAS) Environmental Change Programme (2016-2021); B.E. acknowledges the support from ARC Linkage Grants LP150100519 and LP190100271; M.P. acknowledges US Department of Energy, DE-SC0012536; Lawrence Livermore National Laboratory, B628407 and NASA MAP program, NNX13AL12G; S.B. was supported by the EC H2020 with the CRESCENDO project (grant no. 641816) and by the COMFORT project (grant no. 820989). S.B. also acknowledges the support of the team in charge of the CNRM-CM climate model; F.Z. acknowledges the support from the

National Natural Science Foundation of China (41671464). Supercomputing time was provided by the Météo-France/DSI supercomputing center. P.K.P. is partly supported by Environment Research and Technology Development Fund (#2-1802) of the Ministry of the Environment, Japan. R.L. acknowledges support from the French state aid managed by the ANR under the "Investissements d'avenir" programme with the reference ANR-16-CONV-0003. NOAA ground-based observations of atmospheric N₂O are supported by NOAA's Climate Program Office under the Atmospheric Chemistry Carbon Cycle and Climate (AC4) theme. The AGAGE stations measuring N₂O are supported by NASA (USA) grants NNX16AC98G to MIT, and NNX16AC97G and NNX16AC96G to SIO, and by BEIS (UK) for Mace Head, NOAA (USA) for Barbados, and CSIRO and BoM (Australia) for Cape Grim. We also thank Dr. Steve Frolking and two anonymous reviewers for constructive comments and suggestions that have helped improve this paper. The statements made and views expressed are solely the responsibility of the authors.

Author contributions

Author contributions. H.T., R.L.T., J.G.C. and R.B.J. designed and coordinated the study. H.T., R.X., J.G.C., R.L.T., W.W., P.S., E.A.D., P.C., R.B.J., G.J.M., M.J.P., N.P., S.P., P.R., H.S., F.N.T., S.Z., F.Z., B.F. and G.P. conducted data analysis, synthesis and wrote the paper. R.L.T. led atmospheric inversions teaming with M.P.C., T.M., D.B.M., P.K.P., K.C.W., and C.W.; H.T. led land biosphere modeling teaming with P.C., H.S., S.Z., A.A., F.J., J.C., S.R.S.D., A.I., W.L., S.L., S.O., N.V., E.A.D., S.D. and W. Li; P.S. led ocean biogeochemical modeling teaming with G.B., L.B., S.B., E.T.B., F.J. and A.L.; P.R. led inland water and coastal modeling and synthesis teaming with B.D.E., G.G.L., R.L., T.M., P.A.R., H.T. and Y.Y.; A.F.B., J.W., M.M. provided data of N₂O flux in aquaculture. G.R.W. and J.Y. provided data of N₂O emissions from biomass burning. F.Z. provided cropland N₂O flux data from a statistical model and field observations. G.J.M., F.N.T. and W.W. provided N₂O inventory data. M.J.P. and D.J.R. provided data of stratospheric and tropospheric sinks. G.P.P. provided RCP and SSP scenarios data and analysis. B.H., E.D. and J.E. provided a global N₂O monitoring dataset of NOAA/ESRL GMD. R.P. and R.W. provided a global N₂O monitoring dataset of AGAGE stations. P.K. provided a global N₂O monitoring dataset of CSIRO. All coauthors reviewed and commented on the manuscript.

Competing interests: The authors declare no competing interests.

Additional information

Supplementary information is available for this paper at <https://>

Correspondence and requests for materials should be addressed to H.T.

tianhan@auburn.edu

Extended Data Fig. 1 Global mean growth rates and atmospheric concentration of N₂O.

Global mean growth rates (solid lines, during 1995–2017) and atmospheric N₂O concentration (dashed lines, during 1980–2017) are from the AGAGE⁶ (green), NOAA⁵ (orange), and CSIRO (blue) networks. Global mean growth rates were calculated with annual time steps and are shown as 12-month moving averages. Growth rates are not calculated prior to 1995 due to insufficient data and higher uncertainties on the measurements.

Extended Data Fig. 2 The methodology for data synthesis of global N₂O budget. BU and TD represent bottom-up and top-down methods, respectively. The color codes are the same as that used in Table 1 and Figs. 1–3. We utilize both approaches, including 22 BU and five TD estimates of N₂O fluxes from land and oceans. For sources estimated by BU, we include six process-based terrestrial biosphere modeling studies¹⁶; five process-based ocean biogeochemical models¹⁰⁰; one nutrient budget model^{30,60,61}; five inland water modeling studies^{35,36,50,51,68}; one statistical model SRNM based on spatial extrapolation of field measurements¹⁷; and four GHG inventories: EDGAR v4.3.2¹⁰¹, FAOSTAT¹⁰², GAINS⁴³, and GFED4s¹⁰³. In addition, previous literatures regarding estimates of ‘Surface sink’^{58,73}, ‘Lightning’^{53,54}, ‘Atmospheric production’^{56,57,104}, ‘Aquaculture’^{31,62}, and model-based ‘Tropospheric sink’⁸¹ and observed ‘Stratospheric sink’¹ are included in the current synthesis. ^aMacLeod et al.³¹ and Hu et al.⁶² provide global aquaculture N₂O emissions in 2013 and in 2009, respectively; and the nutrient budget model^{30,60,61} provides N flows in global freshwater and marine aquaculture over the period 1980–2016. ^bModel-based estimates of N₂O emissions from ‘Inland and coastal waters’ include rivers and reservoirs^{35,36}, lakes⁵¹, estuaries³⁵, coastal zones (i.e., seagrasses, mangroves, saltmarsh and intertidal saltmarsh)⁶⁸, and coastal upwelling⁵⁰.

Extended Data Fig. 3 Comparison of annual total N₂O emissions at global and regional scales estimated by BU and TD approaches. The blue lines represent the mean N₂O emission from BU methods and the shaded areas show minimum and maximum estimates; The gold lines represent the mean N₂O emission from TD methods and the shaded areas show minimum and maximum estimates.

Extended Data Fig. 4 Global agricultural N₂O emissions. **a**, Direct emission from agricultural soils associated with mineral fertilizer, manure and crop residue inputs, and cultivation of organic soils based on EDGAR v4.3.2, GAINS, FAOSTAT, NMIP/DLEM, and SRNM/DLEM estimates. NMIP/DLEM or SRNM/DLEM means the combination of N₂O emission by NMIP or SRNM from croplands with N₂O emission from intensively managed grassland (pasture) by DLEM. **b**, Direct emission from the global total area under permanent meadows and pasture, due to manure N deposition (left on pasture) based on EDGAR v4.3.2, FAOSTAT, and GAINS estimates. **c**, Emission from manure management based on FAOSTAT, GAINS, and EDGAR v4.3.2. **d**, Aquaculture N₂O emission based on a nutrient budget model³⁰, MacLeod et al.³¹, and Hu et al.⁶²; the solid line represents the ‘best estimate’ that is the product of EF (1.8%) and N waste from aquaculture provided by the nutrient budget model; the dashed lines represent the minimum and maximum values.

Extended Data Fig. 5 Global N₂O emission from other direct anthropogenic sources. a, Emission from fossil fuel combustion based on EDGAR v4.3.2 and GAINS estimates. **b,** Emission from industry based on EDGAR v4.3.2 and GAINS estimates. **c,** Emission from waste and waste water based on EDGAR v4.3.2 and GAINS estimates. **d,** Emission from biomass burning based on FAOSTAT, DLEM, and GFED4s estimates.

Extended Data Fig. 6 Global N₂O emissions from natural soils, inland and coastal waters and due to change in climate, atmospheric CO₂ and N deposition. **a**, Changes in global soil N₂O fluxes due to changing CO₂ and climate. **b**, Global natural soil N₂O emissions without consideration of land use change (e.g., deforestation) and without consideration of indirect anthropogenic effects via global change (i.e., climate, elevated CO₂, and atmospheric N deposition). The estimates are based on NMIP estimates during 1980–2016 including six process-based land biosphere models. Here, we also subtracted the difference between with and without consideration of secondary forests emissions that grow back after pasture or cropland abandonment from natural soil emissions based on NMIP estimates. The solid lines represent the ensemble and dashed lines show the minimum and maximum values. **c**, Global anthropogenic N₂O emission from inland waters, estuaries, coastal zones based on models (model-based), FAOSTAT, GAINS, and EDGAR v4.3.2 estimates. **d**, Emission due to atmospheric N deposition (NDEP) on land based on NMIP, FAOSTAT/EDGAR v4.3.2, and GAINS/EDGAR v4.3.2. FAOSTAT/EDGAR v4.3.2 or GAINS/EDGAR v4.3.2 means the combination of agricultural source from FAOSTAT or GAINS with non-agricultural source from EDGAR v4.3.2. A process-based model DLEM³⁶ and a mechanistic stochastic model^{35,51} were used to estimate N₂O emission from inland waters and estuaries, while site-level emission rates of N₂O were upscaled to estimate global N₂O fluxes from the global seagrass area⁶⁸.

Extended Data Fig. 7 Global N₂O dynamics due to land cover changes. The blue line represents the mean forest N₂O reduction caused by the long-term effect of reduced mature forest area (i.e., deforestation) and shaded areas show minimum and maximum estimates; the red line represents the mean N₂O emission from post-deforestation pulse effect (i.e., crop/pasture N₂O emissions from legacy N of previous forest soil, not accounting for new fertilizer N added to these crop/pasture lands) and shaded areas show minimum and maximum estimates; the gray line represents the mean net deforestation emission of N₂O and shaded areas show minimum and maximum estimates.

Extended Data Fig. 8 Global simulated N₂O emission anomaly due to climate effect and global annual land surface temperature anomaly during 1901–2016. Global N₂O emission anomalies are the ensemble of six process-based land biosphere models in NMIP. The temperature data were obtained from the CRU-NCEP v8 climate dataset (<https://vesg.ipsl.upmc.fr>). The above left figure **a**) shows the correlation between average global annual land surface temperature and simulated N₂O emissions (i.e., the result of SE6 experiment in NMIP¹⁶) considering annual changes in climate but keeping all other factors (i.e., N fertilizer, manure, NDEP, elevated CO₂, and land cover change) at the level of 1860. The above right figure **b**) shows the correlation between average global annual land surface temperature and simulated N₂O emissions (i.e., the result of SE1 experiment in NMIP¹⁶) considering annual changes in all factors during 1860–2016.

Extended Data Fig. 9 Direct soil emissions and agricultural product trades in Brazil. **a**, Red line shows the ensemble direct N_2O emissions from livestock manure based on EDGAR v4.3.2, GAINS, and FAOSTAT, the sum of ‘manure left on pasture’ and ‘manure management’; The gray columns show the amount of beef export by Brazil. **b**, Orange line shows the ensemble direct N_2O emissions from croplands due to N fertilization based on NMIP and SRNM; The gray columns show the amount of soybean and corn exports by Brazil. The data of beef and cereal product trades were adapted from the ABIEC (beef) and FAOSTAT (soybean and corn). Mmt yr^{-1} represents millions of metric tons per year.

Extended Data Fig. 10 An extension of Fig. 4 to provide a comparison of anthropogenic N₂O emissions (a) and atmospheric N₂O concentrations (b) in the unharmonized SSPs¹⁰⁵. The emission and concentration data are as in Fig. 4. The unharmonized emissions from the Integrated Assessment Models (IAMs)¹⁰⁵ show a large variation due to different input data and model assumptions. Comparison with Fig. 4b, d illustrates the modifications to the IAM scenario data for use in CMIP6. All baseline scenarios (SSP3-7.0 and SSP5-8.5; without climate policy applied) are shown in gray regardless of the radiative forcing level they reach in 2100.

Supplementary Information

A comprehensive quantification of global nitrous oxide sources and sinks

Hanqin Tian¹, Rongting Xu¹, Josep G. Canadell², Rona L. Thompson³, Wilfried Winiwarter^{4,5}, Parvatha Suntharalingam⁶, Eric A. Davidson⁷, Philippe Ciais⁸, Robert B. Jackson^{9,10,11}, Greet Janssens-Maenhout^{12,13}, Michael J. Prather¹⁴, Pierre Regnier¹⁵, Naiqing Pan^{1,16}, Shufen Pan¹, Glen P. Peters¹⁷, Hao Shi¹, Francesco N. Tubiello¹⁸, Sönke Zaehle¹⁹, Feng Zhou²⁰, Almut Arneth²¹, Gianna Battaglia²², Sarah Berthet²³, Laurent Bopp²⁴, Alexander F. Bouwman^{25,26,27}, Erik T. Buitenhuis^{6,28}, Jinfeng Chang^{8,29}, Martyn P. Chipperfield^{30,31}, Shree R. S. Dangal³², Edward Dlugokencky³³, James W. Elkins³³, Bradley D. Eyre³⁴, Bojie Fu^{16,35}, Bradley Hall³³, Akihiko Ito³⁶, Fortunat Joos²², Paul B. Krummel³⁷, Angela Landolfi^{38,39}, Goulven G. Laruelle¹⁵, Ronny Lauerwald^{8,15,40}, Wei Li^{8,41}, Sebastian Lienert²², Taylor Maavara⁴², Michael MacLeod⁴³, Dylan B. Millet⁴⁴, Stefan Olin⁴⁵, Prabir K. Patra^{46,47}, Ronald G. Prinn⁴⁸, Peter A. Raymond⁴², Daniel J. Ruiz¹⁴, Guido R. van der Werf⁴⁹, Nicolas Vuichard⁸, Junjie Wang²⁷, Ray F. Weiss⁵⁰, Kelley C. Wells⁴⁴, Chris Wilson^{30,31}, Jia Yang⁵¹ & Yuanzhi Yao¹

¹International Center for Climate and Global Change Research, School of Forestry and Wildlife Sciences, Auburn University, Auburn, AL, USA

²Global Carbon Project, CSIRO Oceans and Atmosphere, Canberra, Australian Capital Territory, Australia

³Norsk Institutt for Luftforskning, NILU, Kjeller, Norway

⁴International Institute for Applied Systems Analysis, Laxenburg, Austria

⁵Institute of Environmental Engineering, University of Zielona Góra, Zielona Góra, Poland.

⁶School of Environmental Sciences, University of East Anglia, Norwich, UK

⁷Appalachian Laboratory, University of Maryland Center for Environmental Science, Frostburg, MD, USA

⁸Laboratoire des Sciences du Climat et de l'Environnement, LSCE, CEA CNRS, UVSQ UPSACLAY, Gif sur Yvette, France

⁹Department of Earth System Science, Stanford University, Stanford, CA, USA

¹⁰Woods Institute for the Environment, Stanford University, Stanford, CA, USA

¹¹Precourt Institute for Energy, Stanford University, Stanford, CA, USA

¹²European Commission, Joint Research Centre (JRC), Ispra, Italy

¹³Ghent University, Faculty of Engineering and Architecture, Ghent, Belgium

¹⁴Department of Earth System Science, University of California Irvine, Irvine, CA, USA

¹⁵Department Geoscience, Environment & Society, Université Libre de Bruxelles, Brussels, Belgium

¹⁶State Key Laboratory of Urban and Regional Ecology, Research Center for Eco-Environmental Sciences, Chinese Academy of Sciences, Beijing, China

¹⁷CICERO Center for International Climate Research, Oslo, Norway

¹⁸Statistics Division, Food and Agriculture Organization of the United Nations, Via Terme di Caracalla, Rome, Italy

¹⁹Max Planck Institute for Biogeochemistry, Jena, Germany

- ²⁰Sino-France Institute of Earth Systems Science, Laboratory for Earth Surface Processes, College of Urban and Environmental Sciences, Peking University, Beijing, China
- ²¹Karlsruhe Institute of Technology, Institute of Meteorology and Climate Research/Atmospheric Environmental Research, Garmisch-Partenkirchen, Germany
- ²²Climate and Environmental Physics, Physics Institute and Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland
- ²³Centre National de Recherches Météorologiques (CNRM), Université de Toulouse, Météo-France, CNRS, Toulouse, France
- ²⁴LMD-IPSL, Ecole Normale Supérieure / PSL Université, CNRS; Ecole Polytechnique, Sorbonne Université, Paris, France
- ²⁵PBL Netherlands Environmental Assessment Agency, The Hague, The Netherlands
- ²⁶Department of Earth Sciences – Geochemistry, Faculty of Geosciences, Utrecht University, Utrecht, The Netherlands
- ²⁷Key Laboratory of Marine Chemistry Theory and Technology, Ministry of Education, Ocean University of China, Qingdao, China
- ²⁸Tyndall Centre for Climate Change Research, School of Environmental Sciences, University of East Anglia, Norwich, UK
- ²⁹College of Environmental and Resource Sciences, Zhejiang University, Hangzhou, China.
- ³⁰National Centre for Earth Observation, University of Leeds, Leeds, UK
- ³¹Institute for Climate and Atmospheric Science, School of Earth and Environment, University of Leeds, Leeds, UK
- ³²Woods Hole Research Center, Falmouth, MA, USA
- ³³NOAA Global Monitoring Laboratory, Boulder, CO, USA
- ³⁴Centre for Coastal Biogeochemistry, School of Environment Science and Engineering, Southern Cross University, Lismore, New South Wales, Australia
- ³⁵Faculty of Geographical Science, Beijing Normal University, Beijing, China
- ³⁶Center for Global Environmental Research, National Institute for Environmental Studies, Tsukuba, Japan
- ³⁷Climate Science Centre, CSIRO Oceans and Atmosphere, Aspendale, Victoria, Australia
- ³⁸GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany
- ³⁹Istituto di Scienze Marine, Consiglio Nazionale delle Ricerche (CNR), Rome, Italy
- ⁴⁰Université Paris-Saclay, INRAE, AgroParisTech, UMR ECOSYS, Thiverval-Grignon, France
- ⁴¹Ministry of Education Key Laboratory for Earth System modeling, Department of Earth System Science, Tsinghua University, Beijing, China
- ⁴²Yale School of Forestry and Environmental Studies, New Haven, CT, USA
- ⁴³Land Economy, Environment & Society, Scotland's Rural College (SRUC), Edinburgh, UK
- ⁴⁴Department of Soil, Water, and Climate, University of Minnesota, St Paul, MN, USA
- ⁴⁵Department of Physical Geography and Ecosystem Science, Lund University, Lund, Sweden
- ⁴⁶Research Institute for Global Change, JAMSTEC, Yokohama, Japan
- ⁴⁷Center for Environmental Remote Sensing, Chiba University, Chiba, Japan
- ⁴⁸Center for Global Change Science, Massachusetts Institute of Technology, Cambridge, MA, USA
- ⁴⁹Faculty of Science, Vrije Universiteit, Amsterdam, Netherlands.
- ⁵⁰Scripps Institution of Oceanography, University of California San Diego, La Jolla, USA
- ⁵¹Department of Forestry, Mississippi State University, Mississippi State, MS, USA

Supporting text

1. Data sources	1
2. Detailed description on multiple approaches	3
2.1 NMIP – Global N ₂ O Model Inter-comparison Project.....	3
2.2. The FAOSTAT inventory	5
2.3. The EDGAR v4.3.2 inventory.....	7
2.4. The GAINS inventory	8
2.5. The SRNM model	9
a. Flux upscaling model.....	9
b. Global cropland N ₂ O observation dataset	10
c. Gridded input datasets	11
2.6. Global N flow in aquaculture	11
2.7. Model-based ocean N ₂ O fluxes.....	12
2.8. Net N ₂ O emission from land cover change.....	13
a. Deforestation area and crop/pasture expansion	13
c. Secondary tropical forest emissions.	14
2.9. Inland water, estuaries, coastal zones.....	15
a. Riverine N ₂ O emission simulated by DLEM	15
b. The DLEM estimate on N ₂ O emission from global reservoirs	16
c. Mechanistic Stochastic Modeling of N ₂ O emissions from rivers, lakes, reservoirs and estuaries	16
d. Coastal zone emissions.....	18
2.10. Atmospheric inversion models.....	18
3. Atmospheric N ₂ O observations and growth rates for three different atmospheric networks (NOAA, AGAGE, and CISRO).....	21
4. Comparison with the IPCC AR5.....	23
5. Per capita N ₂ O emission at global and regional scales in the recent decade	24

Supporting tables

Supplementary Table 1 N ₂ O emissions from global agricultural soils based on multiple bottom-up approaches including the additions of mineral N fertilizer, manure and crop residues, and cultivation of organic soils.	32
Supplementary Table 2 N ₂ O emissions from global total area under permanent meadows and pasture, due to manure N deposition (left on pasture) based on EDGAR v4.3.2, FAOSTAT, and GAINS estimates.	32
Supplementary Table 3 N ₂ O emissions due to global manure management based on multiple bottom-up approaches.	33
Supplementary Table 4 Aquaculture N ₂ O emissions based on multiple sources.	33
Supplementary Table 5 Anthropogenic N ₂ O emissions from the global inland waters based on process-based models.	34
Supplementary Table 6 Anthropogenic N ₂ O emissions from the global inland waters based on multiple bottom-up approaches.	34
Supplementary Table 7 Natural N ₂ O emissions from the global inland waters based on process-based models.	35
Supplementary Table 8 Nitrous oxide emissions due to atmospheric N deposition on land based on multiple bottom-up approaches.	35
Supplementary Table 9 Global N ₂ O emissions from waste and waste water based on EDGAR v4.3.2 and GAINS estimates.	36
Supplementary Table 10 Global N ₂ O emissions from fossil fuel and industry based on multiple bottom-up approaches.	36
Supplementary Table 11 Global N ₂ O emissions from biomass burning based on multiple bottom-up approaches.	37
Supplementary Table 12 Global oceanic N ₂ O emissions based on multiple models.	37
Supplementary Table 13 Global N ₂ O emissions based on multiple top-down approaches.	38
Supplementary Table 14 Comparison of terminologies used in this study and previous reports.	39
Supplementary Table 15 Comparison of the global N ₂ O budget in this study with the IPCC AR5.	40
Supplementary Table 16 Simulation experiments in the NMIP (Tian et al. ^{1,17}).....	41
Supplementary Table 17 Information of NMIP models using in this study	41
Supplementary Table 18 Summary of models in ocean N ₂ O inter-comparison	42
Supplementary Table 19 Overview of the inversion frameworks that are included in the global N ₂ O budget.....	42

Supporting figures

Supplementary Fig. 1 Spatial distribution of ten study regions across the globe.	43
Supplementary Fig. 2 Per capita N ₂ O emission (kg N capita ⁻¹ yr ⁻¹) during 2007–2016.	44

1. Data sources

Bottom-up methods include process-based models (NMIP¹ including six process-based terrestrial biosphere models, DLEM-only² for pastureland, five ocean models³⁻⁸, one mechanistic stochastic model^{9,10}), four GHG emission databases [EDGAR v4.3.2¹¹, FAOSTAT¹², GAINS¹³, GFED4s¹⁴ (only for biomass burning), and one statistical model (SRNM) only for cropland soils¹⁵. The top-down approach includes four independent atmospheric inversion frameworks¹⁶. The NMIP result provides nitrous oxide (N₂O) emissions from natural and agricultural soils, defined as soils in agricultural land, during 1860–2016, with consideration of multiple environmental factors, such as climate, elevated atmospheric carbon dioxide (CO₂), land cover and land use change, atmospheric nitrogen (N) deposition, mineral N fertilizer, and manure N in cropland¹⁷. Mineral N fertilizer and manure N are mainly applied to cropland, while N deposition can reach soils under all land uses. Natural soil emissions were estimated by NMIP based on the ensemble mean of six models (Supplementary Table 16): (1) the Dynamic Land Ecosystem Model (DLEM)^{18,19}, (2) Land Processes and eXchanges model - Bern (LPX-Bern v1.4)^{20,21}, (3) O-CN²², (4) Organising Carbon and Hydrology In Dynamic Ecosystems (ORCHIDEE)²³, (5) Organising Carbon and Hydrology In Dynamic Ecosystems-Carbon Nitrogen Phosphorus (ORCHIDEE-CNP)²³, and (6) Vegetation Integrated Simulator for Trace gases (VISIT)^{24,25} (See more model information in Tian et al.^{1,17}). Agricultural soil emissions were from manure and fertilizer N applications on cropland during 1860–2016¹ and intensively managed grassland (pastures) during 1900–2014². For ‘Indirect emissions from anthropogenic N additions’, we considered emissions from atmospheric N deposition and ‘Inland and coastal waters’ N leaching/runoff including five sub-systems: rivers, lakes, estuaries, reservoirs, and coastal zones. Yao et al.²⁶ estimated N₂O emissions from rivers using the process-based model (DLEM) during 1900–2016 and provided estimates from reservoirs as well, while emissions determined from the stochastic mechanistic model of Maavara et al.¹⁰ and Lauerwald et al.⁹ for the river-reservoir-estuary continuum, and lakes, respectively, were in 2000. Coastal zone emissions were obtained from data compilation reported in Camillini et al.²⁷ and Murray et al.²⁸. The DLEM model also provided an estimate of N₂O emissions from biomass burning across various biomes (crop residue and savannas, peatland, tropical forest, temperate forest, and boreal forest) during 1860–2015. A nutrient budget model for shellfish and finfish²⁹⁻³¹ was used to calculate the nutrient flows in aquaculture

production systems. For computing the N₂O emission we consider the amount of N released to the environment, i.e. the difference between N intake and N in the harvested fish, which includes all the nutrient excretion. Estimates of oceanic N₂O fluxes are derived from an inter-comparison of five global ocean biogeochemistry models including Bern-3D³, NEMOv3.6-PISCESv2-gas⁴, NEMO-PlankTOM10⁵, UVic2.9⁶, and NEMO-PISCES 3.2⁷.

The EDGAR v4.3.2 applies the IPCC guidelines mostly at Tier-1, but integrates higher tier information based on available country reporting, mostly from Annex I countries. It provided data from 1970 to 2012. We updated the data to 2016 based on the global and regional trends between 2000 and 2012 for each individual category. In EDGAR v4.3.2, ‘Indirect emission from N deposition’ only represents non-agricultural activities. ‘Waste and waste water’ includes ‘Waste incineration’ and ‘Wastewater handling’. We merged ‘Transportation’, ‘Energy’, ‘Industry’, and ‘Residential and other sectors’ to represent the total emission from ‘Fossil fuel and industry’. Since the EDGAR v4.3.2 database did not provide the emission of ‘Biomass burning’ from land use outside of agriculture, here we did not include its estimate of ‘Agriculture waste burning’ into the data synthesis. The FAOSTAT emissions database of the Food and Agriculture Organization of the United Nations (FAO) covers emissions of N₂O from agriculture and land use by country and globally, from 1961 to 2017 for agriculture, and from 1990¹² for relevant land use categories, i.e. cultivation of histosols, biomass burning, etc., applying only Tier-1 coefficients³². In addition to the IPCC agriculture burning categories ‘Burning crop residues’ and ‘Burning savannah’, FAOSTAT also estimates N₂O emissions from deforestation fires, forest fires and peatland fires. Emissions from ‘Fossil fuel and industry’ are directly adapted from the EDGAR v4.3.2 emission inventory. The GAINS model¹³ provided N₂O emissions data every five-years (i.e., 1990, 1995, 2000, 2005, 2010, 2015). We assumed the change between each five-year estimate was linear. To avoid abrupt jumps between 1989 and 1990 during data synthesis, we linearly extrapolated data to 1980 through using estimates in 1990 and 1995 in each sub-sector. ‘Direct soil emissions’ are from synthetic N fertilizer, animal manure, cultivation of histosols, and crop residues. ‘Indirect emissions from anthropogenic N additions’ are from ‘N deposition on land’ and ‘Inland and coastal waters’ (i.e., lakes, rivers, and shelf seas). The source of ‘N deposition on land’ is mainly from agricultural activities, but it deposited on all global ice-free areas (i.e., agricultural land, forest land, other land uses). The ‘Energy’ emission includes conversion, industry, transport, and domestic. The ‘Industry’

emission includes nitric acid plants, adipic acid plants, and caprolactam plants. We merged ‘Energy’ and ‘Industry’ to represent ‘Fossil fuel and industry’ emissions. They also considered N₂O use, but we did not include this sector in the synthesis table. We merged ‘Composting’ and ‘Wastewater’ sectors into ‘Waste and waste water’ to make comparison with the EDGAR v4.3.2 database. In addition, the sector ‘Grazing’ was treated as ‘Manure left on pasture’ to make comparison. The GFED4s emission inventory¹⁴ provided N₂O emissions from ‘Biomass burning’ including agricultural waste and other biomass burning (i.e., Savanna, grassland, and shrubland fires, boreal forest fires, temperate forest fires, deforestation and degradation, and peatland fires) during 1997–2016.

The spatially-referenced non-linear model SRNM was fitted through considering environmental factors and N management practices to generate gridded annual EF maps at 5’ spatial resolution, and then to calculate global/regional N₂O emissions during 1901–2016 together with time-series N input datasets¹⁵. This database provides N₂O emissions from global and regional cropland with the application of synthetic N fertilizer and manure N for the period 1980–2016.

For the top-down constraints on the global and regional N₂O emissions for the period 1998–2016, we have used estimates from four independent atmospheric inversion frameworks (INVICAT, PyVAR, MIROC4-ACTM, and GEOSChem), all of which used the Bayesian inversion method. Here, two versions of PyVAR were run using different ocean priors (one high and one low) for determining the sensitivity to the ocean prior. These runs are denoted as PyVAR-1 and PyVAR-2, respectively. For the top-down global estimate, we used the original spatial resolution in each framework. For the top-down regional estimate, we interpolated the coarse resolution into $1^{\circ} \times 1^{\circ}$ to cover all land areas in the four frameworks (see details in section 2.10).

2. Detailed description on multiple approaches

2.1 NMIP – Global N₂O Model Inter-comparison Project

Ten process-based Terrestrial Biosphere Models (TBMs) participate in NMIP. In general, N₂O emissions from soil are regulated at two levels, which are the rates of nitrification and denitrification in the soil and soil physical factors regulating the ratio of N₂O to other nitrous

gases³³. For N input to land ecosystems, all ten models considered the atmospheric N deposition and biological fixation, nine models with crop N₂O module included N fertilizer use, but only six models considered manure as N input. For vegetation processes, all models included dynamic algorithms in simulating N allocation to different living tissues and vegetation N turnover, and simulated plant N uptake using the “Demand and Supply-driven” approach. For soil N processes, all ten models simulated N leaching according to water runoff rate; however, models are different in representing nitrification and denitrification processes and the impacts of soil chemical and physical factors. The differences in simulating nitrification and denitrification processes are one of the major uncertainties in estimating N₂O emissions. Algorithms associated with N₂O emissions in each participating model are briefly described in Appendix A of Tian et al.¹⁷.

All participating models are driven by consistent input datasets (i.e., climate, atmospheric CO₂ concentration, land cover change, atmospheric N deposition, mineral N fertilization, and manure N application) and implemented seven simulation experiments (SE0 – SE6; Supplementary Table 17) at the spatial resolution of 0.5° globally covering the period of 1861–2016 (ref. ¹). The SE1 includes all driving factors for models with manure addition, and the SE2 is the experiment including all the driving factors for models except manure N. In the SE0 simulation, driving forces were kept constant at the level in 1860 over the entire simulation period (1861–2016).

By comparing results from different model scenarios, it is possible to attribute the changed spatiotemporal variations of soil N₂O emissions to the variations of six natural and anthropogenic factors, namely, climate (CLIM, including precipitation, humidity, temperature and photosynthetic active radiation changes), atmospheric CO₂ concentration (CO₂), land cover change (LCC), atmospheric N deposition (NDEP), mineral N fertilizer use (NFER), and manure N use in cropland (MANN). In order to understand soil N₂O emissions dynamics caused by crop cultivation, we further separated the global and regional N₂O emissions into those derived from cropland soils and those from soils of other land ecosystems. All soils in other land ecosystems except cropland were treated as “natural soils” while model simulations were implemented in this study. Except for cropland, the current NMIP simulations do not include management

practices (such as grazing and forest logging) for other managed ecosystems such as pasture, planted forests and urban.

In this study, we aimed to attribute the impact of single factor on cropland N₂O emissions, thus participating models without providing SE2–SE6 and SE0 results in cropland were excluded. Here, we included estimates from six process-based models (Supplementary Table 16). Four models (DLEM, ORCHIDEE, ORCHIDEE-CNP, and VISIT) considered the effects of manure N application in cropland and ran all the seven simulation experiments (SE0–SE6), while the other two models (LPX-Bern and O-CN) did not include manure effects and ran six model experiments (all except SE1). We used four model results (i.e., DLEM, ORCHIDEE, ORCHIDEE-CNP, and VISIT) to calculate the manure N effect (SE1–SE2). Meanwhile, we used six model results (i.e., DLEM, LPX-Bern, O-CN, ORCHIDEE, ORCHIDEE-CNP, and VISIT) to calculate the effects of synthetic N fertilizer use (SE2–SE3) and atmospheric N deposition (SE3–SE4). The effect of N deposition in natural vegetation was calculated from the six models mentioned above.

2.2. The FAOSTAT inventory

The FAOSTAT emissions data are computed at Tier 1 following IPCC, 2006, Vol. 4. The overall equation is as follows:

Direct emissions are estimated at country level, using the formula:

$$Emission = A * EF \quad (1a)$$

where emission represents kg N yr⁻¹; *A* represents amount of N in the following items (annual synthetic N applications/manure applied to soils/manure left on pasture/manure treated in manure management systems/crop residue/biomass burned amount) in kg N yr⁻¹; *EF* = Tier 1, default IPCC emission factors, expressed in kg N/kg N.

Indirect emissions are estimated at country level, using the formula:

$$Emission = A_{v\&l} * EF \quad (1b)$$

where emission represents kg N yr^{-1} ; $A_{v\&l}$ represents the fraction of manure/synthetic N fertilizers that volatilize as NH_3 and NO_x and are lost through runoff and leaching in kg N yr^{-1} ; EF = Tier 1, default IPCC emission factors, expressed in kg N/kg N .

Synthetic N fertilizers: N_2O from synthetic fertilizers is produced by microbial processes of nitrification and denitrification taking place on the addition site (direct emissions), and after volatilization/redeposition and leaching processes (indirect emissions).

Manure management: The term manure includes both urine and dung (i.e., both liquid and solid material) produced by livestock. N_2O is produced directly by nitrification and denitrification processes in the manure, and indirectly by N volatilization and redeposition processes.

Manure applied to soils: N_2O is produced by microbial processes of nitrification and denitrification taking place on the application site (direct emissions), and after volatilization/redeposition and leaching processes (indirect emissions).

Manure left on pastures: N_2O is produced by microbial processes of nitrification and denitrification taking place on the deposition site (direct emissions), and after volatilization/redeposition and leaching processes (indirect emissions).

Crop Residue: N_2O emissions from crop residues consist of direct and indirect emissions from N in crop residues left on agricultural fields by farmers and from forages during pasture renewal (following the definitions in the IPCC guidelines³⁴). Specifically, N_2O is produced by microbial processes of nitrification and denitrification taking place on the deposition site (direct emissions), and leaching processes (indirect emissions).

Cultivation of organic soils: The FAOSTAT domain “Cultivation of organic soils” contains estimates of direct N_2O emissions associated with the drainage of organic soils – histosols – under cropland and grazed grassland.

Burning-savanna: N_2O emissions from the burning of vegetation biomass in the land cover types: Savanna, Woody Savanna, Open Shrublands, Closed Shrublands, and Grasslands.

Burning-crop residues: N_2O produced by the combustion of a percentage of crop residues burnt on-site. Burning-biomass: N_2O emissions from the burning of vegetation biomass in the land cover types: Humid tropical forest, other forest, and organic soils.

2.3. The EDGAR v4.3.2 inventory

The new online version, EDGAR v4.3.2 incorporates a full differentiation of emission processes with technology-specific emission factors and additional end-of-pipe abatement measures⁶ and as such updates and refines the emission estimates. The emissions are modelled based on latest scientific knowledge, available global statistics, and methods recommended by IPCC (2006)³⁴. Official data submitted by the Annex I countries to the United Nations Framework Convention on Climate Change (UNFCCC) and to the Kyoto Protocol are used to some extent, particularly regarding control measures implemented since 1990 that are not described by international statistics.

The N₂O emission factor for direct soil emissions of N₂O from the use of synthetic fertilizers and from manure used as fertilizers and from crop residues is taken from IPCC (2006)³⁴, that updated the default IPCC emission factor in the IPCC Good Practice Guidance (2000) with a 20% lower value. N₂O emissions from the use of animal waste as fertilizer are estimated taking into account both the loss of N that occurs from manure management systems before manure is applied to soils and the additional N introduced by bedding material. N₂O emissions from fertilizer use and CO₂ from urea fertilization are estimated based on IFA and FAO statistics.

N₂O emissions from manure management are based on distribution of manure management systems from Annex I countries reporting to the UNFCCC, Zhou et al.³⁵ for China and IPCC (2006)³⁴ for the rest of the countries.

Different N₂O emission factors are applied to tropical and non-tropical regions. N and dry matter content of agricultural residues are estimated from the cultivation area and yield for 24 crop types (two types of beans, barley, cassava, cereals, three types of peas, lentils, maize, millet, oats, two types of potatoes, pulses, roots and tubers, rice, rye, soybeans, sugar beet, sugar cane, sorghum, wheat and yams) from FAOSTAT (2014) and using emission factors of IPCC (2006)³⁴.

Indirect N₂O emissions from leaching and runoff of nitrate are estimated from N input to agricultural soils as described above. Leaching and runoff are assumed to occur in all agricultural areas except non-irrigated dryland regions, which are identified with maps of FAO Geonetwork (2011). The fraction of N lost through leaching and runoff is based on the study of Van Dreht et al.³⁶. The updated emission factor for indirect N₂O emissions from N leaching and run-off from

the IPCC (2006) guidelines is selected, while noting that it is 70% lower than the mean value of the 1996 IPCC Guidelines and the IPCC Good Practice Guidance (IPCC, 1997, 2000).

Indirect N₂O emissions from atmospheric deposition of N of NO_x and NH₃ emissions from non-agricultural sources, mainly fossil fuel combustion, are estimated using N in NO_x and NH₃ emissions from these sources as activity data, based on EDGAR v4.3.2 database for these gases. The same emission factor from IPCC (2006)³⁴ is used for indirect N₂O from atmospheric deposition of N from NH₃ and NO_x emissions, as for agricultural emissions.

2.4. The GAINS inventory

The methodology adopted for the estimation of current and future greenhouse gas emissions and the available potential for emission controls follows the standard methodology used by the Greenhouse Gas - Air Pollution Interactions and Synergies (GAINS) model³⁷. For a given year, emissions of each pollutant p are calculated as the product of the activity levels, the “uncontrolled” emission factor in the absence of any emission control measures, the efficiency of emission control measures and the application rate of such measures:

$$E_{i,p} = \sum_{j,a,t} E_{i,j,a,t,p} = \sum_{j,a,t} A_{i,j,a} ef_{i,j,a,p} (1 - eff_{t,p}) X_{i,j,a,t} \quad (2)$$

where subscripts i,j,a,t,p denote region, sector, activity, abatement technology, and pollutant, respectively; $E_{i,p}$ represents emissions of the specific pollutant p in region i ; A_j represents activity in a given sector j ; ef represents “uncontrolled” emission factor; eff represents reduction efficiency; X represents actual implementation rate of the considered abatement.

Results (emissions from all anthropogenic sources) are available in 5-year intervals from 1990 to 2050 (in some regions up to 2070) for each GAINS region, typically comprising one country to express areas of common legislation also with respect of air pollution. Very large countries have been further split along administrative areas, while in cases of limited data availability also groups of countries have been combined into GAINS regions.

For N₂O, the fate of emissions abatement is often connected with action taken to control other pollutants. For example, it may occur that after control (e.g., of NO_x emissions), N₂O emissions become higher than in the unabated case. To reflect this effect, negative reduction efficiencies would need to be used for N₂O. As it is difficult to communicate such negative numbers, GAINS

has resorted to present “controlled” emission factors instead, which describe the emission factor of a process after installation of abatement technology.

2.5. The SRNM model

a. Flux upscaling model

The SRNM model³⁸ was applied to simulate direct cropland-N₂O emissions. In SRNM, N₂O emissions were simulated from N application rates using a quadratic relationship, with spatially-variable model parameters that depend on climate, soil properties, and management practices. The original version of SRNM was calibrated using field observations from China only³⁹. In this study, we used the global N₂O observation dataset to train it to create maps of gridded annual emission factors of N₂O and the associated emissions at 5-minute resolution from 1901 to 2014¹⁵. The gridded EF and associated direct cropland-N₂O emissions are simulated based on the following equation:

$$E_{ijt} = \alpha_{ij} N_{ijt}^2 + \beta_{ij} N_{ijt} + \varepsilon_{ijt}, \quad \forall i \quad (3a)$$

where

$$\alpha_{ij} \sim N\left(\sum_k (x_k \lambda_{ijk}), \sigma_{ijk}^2\right), \quad \beta_{ij} \sim N\left(\sum_k (x_k \phi_{ijk}), \sigma_{ijk}'^2\right) \quad (3b)$$

$$\lambda_{ijk} \sim N(\mu_{ijk}, \omega_{ijk}^2), \quad \phi_{ijk} \sim N(\mu'_{ijk}, \omega_{ijk}'^2), \quad \varepsilon_{ijt} \sim N(0, \tau^2) \quad (3c)$$

and i denotes the sub-function of N₂O emission ($i=1, 2, \dots, I$) that applies for a sub-domain division Ω_i of six climate or soil factors, j represents the type of crop ($j=1-2$, 1 for upland crops and 2 for paddy rice), k is the index of climate or soil factors ($k=1-6$, i.e., soil pH, clay content, soil organic carbon, bulk density, the sum of cumulative precipitation and irrigation, mean daily air temperature). Ω_i denotes a set of the range of multiple x_k . E_{ijt} denotes direct N₂O emission flux (kg N ha⁻¹ yr⁻¹) estimated for crop type j in year t in the i th sub-domain, N_{ijt} is N application rate (kg N ha⁻¹ yr⁻¹), and α_{ij} and β_{ij} are defined as summation of the product of x_k and λ_{ijk} over k . The random terms λ and ϕ are assumed to be independent and normally distributed, representing

the sensitivity of α and β to x_k . ε is the model error. μ and μ' are the mean effect of x_k for α and β , respectively. σ , σ' , ω , ω' , and τ are standard deviations. Optimal sub-domain division, associated parameters mean values and standard deviations were determined by using the Bayesian Recursive Regression Tree version 2 (BRRT v2)³⁹⁻⁴¹, constrained by the extended global cropland-N₂O observation dataset. The detailed methodological approach of the BRRT v2 is described in Zhou et al.⁴¹

b. Global cropland N₂O observation dataset

We aggregated cropland N₂O flux observation data from 180 globally distributed observation sites from online databases, on-going observation networks, and peer-reviewed publications. Chamber-based observations were only included in this dataset. These data repositories are as follows: the NitroEurope, CarbonEurope, GHG-Europe (EU-FP7), GRACEnet, TRAGnet, NANORP, and 14 meta-analysis datasets⁴²⁻⁵⁵. Four types of data were excluded from our analysis: (i) observations without a zero-N control for background N₂O emission, (ii) observations from sites that used controlled-release fertilizers or nitrification inhibitors, (iii) observations not covering the entire crop growing season, (iv) observations made in laboratory or greenhouse. We then calculated cropland-N₂O emissions as the difference between observed N₂O emission (E) and background N₂O emission (E₀). Values of EF were estimated for each nonzero N application rate (N_a) as direct cropland-N₂O emission divided by N_a : $EF = (E - E_0)/N_a$. This yielded a global dataset of direct cropland-N₂O emissions, N-rate-dependent N₂O EFs and fertilization records from each site (i.e., 1,052 estimates for upland crops from 152 sites and 154 estimates for paddy rice from 28 sites), along with site-level information on climate, soils, crop type, and relevant experimental parameters. Total numbers of sites and total measurements in the dataset were more than doubled those for previous datasets of N₂O EF. The extended global N₂O observation network covered most of fertilized croplands, representing a wide range of environmental conditions globally. For each site in our dataset, the variables included four broad categories: N₂O emissions data, climate data (cumulative precipitation and mean daily air temperature), soil attributes (soil pH, clay content, SOC, BD), and management-related or experimental parameters (N application rate, crop type). More details on global cropland N₂O observation dataset can be found in Wang et al.¹⁵

c. Gridded input datasets

The updated SRNM model was driven by many input datasets, including climate, soil properties, N inputs (e.g., synthetic N fertilizer, livestock manure and crop residues applied to cropland), as well as the historical distribution of cropland. Cumulative precipitation and mean daily air temperature over the growing season were acquired from the CRU TS v3.23 climate dataset⁴³ (0.5-degree resolution), where growing season in each grid cell was identified following Sacks et al.⁵² The patterns of SOC, clay content, BD, and soil pH were acquired from the HWSD v1.2 (ref.⁵⁶, 1-km resolution). Both climate and soil properties were re-gridded at a resolution of 5' × 5' using a first-order conservative interpolation widely used in the CMIP5 model intercomparison⁵⁷. Annual cropland area at 5' spatial resolution from 1901 to 2014 was obtained from the History Database of the Global Environment (HYDE 3.2.1)⁵⁸. N inputs of synthetic fertilizers were generated based on sub-national statistics (i.e., county-, municipal, provincial or state-levels) of N-fertilizer consumption of 15,593 administrative units from 38 national statistical agencies and national statistics of the other 197 countries from FAOSTAT. N inputs of livestock manure and crop residues applied to cropland were provided by Zhang et al.⁵⁹ and FAOSTAT, respectively. To compute crop-specific N application rates, we allocated N inputs for upland crops and paddy rice based on the breakdown (or proportion) of total fertilizer use by crop from Rosas⁶⁰. Crop-specific N application rates (N_{ijt}) were finally resampled into grid maps at 5' spatial resolution following the dynamic cropland distributions of the HYDE 3.2.1. The assumption of a maximum combined synthetic + manure + crop residues N application rate was 1,000 kg N ha⁻¹, larger than the previous threshold⁶¹ that was only applied for the sum of synthetic fertilizers and manure.

2.6. Global N flow in aquaculture

We apply a nutrient budget model for shellfish and finfish⁶²⁻⁶⁴ to calculate the nutrient flows in aquaculture production systems. These flows comprise feed inputs, retention in the fish, and nutrient excretion. Individual species within crustaceans, seaweed, fish and molluscs are aggregated to the International Standard Statistical Classification of Aquatic Animals and Plants (ISSCAAP) groups⁶⁵, for which production characteristics are specified. Feed and nutrient conversion rates are used for each ISSCAAP group to calculate the feed and nutrient intake from

production data from FAO⁶⁵. Feed types include home-made aquafeeds and commercial compound feeds with different feed conversion ratios that also vary in time due to efficiency improvement; in addition, the model accounts for algae in ponds, that are often fertilized with commercial fertilizers or animal manure, consumed by omnivore fish species like carp. A special case is the filter-feeding bivalves that filter seston from the water column, and excrete pseudofeces, feces and dissolved nutrients. Based on production data and tissue/shell nutrient contents the model computes the nutrient retention in the fish. Using apparent digestibility coefficients, the model calculates outflows in the form of feces (i.e. particulate nutrients) and dissolved nutrients. Finally, nutrient deposition in pond systems and recycling is calculated. For computing the N₂O emission we consider the amount of N released to the environment, i.e. the difference between N intake and N in the harvested fish, which includes all the nutrient excretion. Since in pond cultures part of that N is managed, we made the amount of N recycling explicit, as well as ammonia emissions from ponds. This is to avoid double counting when computing N₂O emissions from crop production.

2.7. Model-based ocean N₂O fluxes

Oceanic N₂O is produced by microbial activity during organic matter cycling in the subsurface ocean; its production mechanisms display significant sensitivity to ambient oxygen level. In the oxic ocean, N₂O is produced as a byproduct during the oxidation of ammonia to nitrate, mediated by ammonia oxidizing bacteria and archaea. N₂O is also produced and consumed in sub-oxic and anoxic waters through the action of marine denitrifiers during the multi-step reduction of nitrate to gaseous N. The oceanic N₂O distribution therefore displays significant heterogeneity with background levels of 10-20 nmol/l in the well-oxygenated ocean basins, high concentrations (> 40 nmol/l) in hypoxic waters, and N₂O depletion in the core of ocean oxygen minimum zones (OMZs).

Oceanic N₂O emissions are estimated to account for up to a third of the pre-industrial N₂O fluxes to the atmosphere, however, the natural cycle of ocean N₂O has been perturbed in recent decades by inputs of anthropogenically derived nutrient (via atmospheric deposition and riverine fluxes), and by the impacts of climate change (via impacts on biological productivity and ocean deoxygenation).

Estimates of oceanic N₂O fluxes for the Global N₂O Budget synthesis are derived from an inter-comparison of five global ocean biogeochemistry models that include explicit representation of the oceanic N₂O cycle (Supplementary Table 18). Ocean biogeochemistry models include process representation of ocean circulation, nutrient cycling and trace-gas generation. In particular, the N₂O fluxes to the atmosphere are derived from N₂O cycle parameterizations embedded in the ocean biogeochemistry models and combined with a parameterization of gas-exchange across the air-sea interface. The models participating in this inter-comparison are taken from the recent studies of Battaglia and Joos³, Berthet et al.⁴, Buitenhuis et al.⁵, Landolfi et al.⁶, and Martinez-Rey et al.⁷.

The models differ in aspects of physical configuration (e.g., spatial resolution), meteorological forcing applied at the ocean surface, and in their parameterizations of ocean biogeochemistry; specific details on individual models are provided in the publications listed in Supplementary Table 18. Towards the N₂O budget synthesis, all modelling groups reported annual mean estimates of ocean-atmosphere N₂O fluxes for the period 1980–2016 (or for as many years as possible in that period). Fluxes were reported at the following spatial scales: (a) global; (b) Southern latitudes (90°–30°S); (c) Tropics (30°S–30°N); and (d) Northern latitudes (30°–90°N). In addition, four modelling groups reported annually averaged ocean N₂O fluxes at higher spatial resolution; i.e., gridded to a 1° × 1° resolution.

2.8. Net N₂O emission from land cover change

a. Deforestation area and crop/pasture expansion

Two sets of deforestation area were used to represent land cover changes during 1860–2016. The LUH2 v2h (land use harmonization, <http://luh.umd.edu>) land use forcing data were used to derive the deforestation area and its partition between crops and pastures from 1860–2016. LUH2 categorizes forest lands into forested primary land and potentially forested secondary land, while croplands are divided into C3 annual crops, C3 perennial crops, C4 annual crops, C4 perennial crops, and C3 N-fixing crops. In the empirical computation, all sub-classes within each land use type were treated the same. Thus only the annual transition area from forests to croplands or managed pasture was needed.

In the process-based estimates, the model requires input of the plant functional types (PFTs) of the forests (e.g., tropical broadleaf evergreen forest and tropical broadleaf deciduous forest), and the species of croplands (e.g., wheat and rice). Thus, a potential vegetation map and the accompanied composition ratio map of each natural PFT acquired from the Synergetic Land Cover Product (SYNMAP) were jointly used with LUH2 v2h to generate the historical spatial distribution of PFTs.

b. Methods

Here we ran the DLEM model with varying climate and CO₂ but hold other factors constant to estimate forest baseline emissions and unfertilized crop/pasture emissions from 1860-2016. The climate data were acquired from CRU-NCEP v7 (<https://vesg.ipsl.upmc.fr>), which is a fusion of the CRU and NCEP/NCAR reanalysis products at a spatial resolution of $0.5^\circ \times 0.5^\circ$ and a daily time-step. The atmospheric CO₂ data were obtained from NOAA GLOBLVIEW-CO2 dataset (<https://www.esrl.noaa.gov>), which are derived from atmospheric and ice core measurements. In the tropical area, both estimates from the DLEM model and the bookkeeping method were adopted, whereas in extra-tropical area, we only adopted the DLEM outputs.

c. Secondary tropical forest emissions

There are not many published studies on N₂O emissions from secondary tropical forests that grow back after crop or pasture abandonment. A recent meta-analysis by Sullivan et al.⁶⁶ lumps together all forms of N "gas loss" including NO and N₂O, so it does not address N₂O specifically. It also reviews the data for secondary forests across the tropics and shows that eight N cycling parameters, including N gas loss and some other parameters that overlap with those measured by Davidson et al.⁶⁷ and Keller and Reiners⁶⁸, recover only gradually during secondary tropical forest succession. Their meta-analysis of the N gas loss parameter showed a significant positive slope, indicating gradually increasing gas loss rates with age after initiation of secondary forest regrowth⁶⁶. Keller and Reiners⁶⁸ showed a gradual recovery of soil nitrate and soil emissions of N₂O and nitric oxide (NO) during 20 years of secondary forest succession. As shown, N₂O emissions did not return to the level of the primary forest after about 20 years of secondary forest succession. Davidson et al.⁶⁷ found that it takes 40–70 years of secondary forest succession for N₂O emissions to approach levels of the primary forest. This is also consistent with other trends of related N cycling parameters, such as the nitrate:ammonium ratio, soil

nitrate, litter mass:N, litterfall N:P, and foliar ^{15}N . In this study, through using the sites of field observation from Davidson et al.⁶⁷ and Keller and Reiners⁶⁸, we estimated N_2O emission from secondary tropical forests based on the algorithm: $y=0.0084x + 0.2401$ ($R^2 = 0.44$). x (unit: year) indicates secondary forest age and y (unitless; 0–1) indicates the ratio of secondary forest N_2O emission over that of a reference mature forest. The difference between primary forests and secondary forests were subtracted from natural soil emissions simulated by six land-surface models in NMIP.

2.9. Inland water, estuaries, coastal zones

a. Riverine N_2O emission simulated by DLEM

Here we developed a riverine N_2O module within a scale adaptive water transport model and coupled with the DLEM model¹⁵. The land surface module of DLEM-simulated N species (NO_3^- , NH_4^+ , DON and PON) leaching from soils when N inputs were into the water transport model. The river routine module within the DLEM is a fully distributed water transport model, which explicitly calculated the flow routine cell-to-cell based on hydraulics methods. The water quality module built into the water transport module can simulate the carbon lateral transportation, biogeochemical reactions (e.g., decomposition of organic matter, nitrification, denitrification), CO_2 degassing and physical deposition of particle organic matter and has been successfully applied in the Gulf of Mexico and the U.S. east coast⁶⁹⁻⁷⁵. Specifically, by introducing sub-grid routine processes technology into the model, the scale adaptive water transport module can effectively address the physical and biogeochemical processes of the small streams within a grid cell, which has been overly simplified in earth system models. We validated global N fluxes based on GEMS-GLORI world river discharge database. The newly developed riverine N_2O module receives dissolved N_2O from land and groundwater, atmosphere wet deposition, and calculate the dynamics of dissolved N_2O concentration and fluxes in both small streams and large rivers. Here, we validated the annual mean riverine N_2O concentration, ground water N_2O concentration, and riverine N_2O emissions globally based on literature survey. DLEM simulated results all agree well with the observations.

b. The DLEM estimate on N₂O emission from global reservoirs

We assumed the reservoirs were linked to rivers, and thus these aquatic systems shared the similar N₂O emission rates in the large-scale studies. We therefore estimate the reservoir surface-area from the Global Reservoirs and Dams (GRanD) database. In riverine N₂O fluxes estimations, we have two N₂O fluxes rates: one is the emission from the large river channel, and the other one is the emission from small rivers within the grid cell. We obtained the upstream area of each dam from the GRanD database and overlaid with the area raster of the 0.5° cell. If the upstream area of a dam is less than the area of its belonging 0.5° grid cell, we considered the dam was located at the small streams within the grid cell and the fluxes of that dam equal to the small river N₂O fluxes of that grid. On the contrary, if the upstream area was larger than the area of the grid cell, the dam is located at the large river channel, thus the fluxes of that dam equal to the riverine N₂O fluxes of the main channel in that grid cell. Align with uncertainty analysis in the riverine N₂O estimations, we overlaid the surface area of dams with riverine N₂O emission rate estimates from the nine-uncertainty experiments to get the reservoir N₂O emissions. We calculated the average as the final reported value.

c. Mechanistic Stochastic Modeling of N₂O emissions from rivers, lakes, reservoirs and estuaries

In our calculations, we used a process-oriented model recently developed to estimate N₂O emissions from inland waters, including rivers, reservoirs and estuaries¹⁰. To estimate N₂O emissions from lakes⁹, we applied the same approach to a global lake dataset⁷⁶. Based on a spatially explicit representation of water bodies and point and non-point sources of N and phosphorus (P), this model quantifies the global scale spatial patterns in inland water N₂O emissions in a consistent manner at 0.5° resolution. The methodology is based on the application of a stochastic Monte Carlo-based model to estimate average annual rates of primary production, ammonification, nitrification, denitrification, N fixation and burial of N in sediments as well as N₂O production and emission generated by nitrification and denitrification. Because of the scarcity of observations, the Monte Carlo approach is a necessary step to generate predictive equations for the N budget and N₂O emission of each inland water body based on inputs of total N (TN) and total P (TP) from the watershed and water residence times in a given river segment, lake, reservoir or estuary^{9,10}. In situ N cycling processes for each specific water body worldwide cannot be predicted due to the lack of parameter constraints or data at this fine granularity.

Instead, the model is fed with hypothetical but realistic combinations of physical and biogeochemical parameters through the use of probability density functions (PDFs) approximating the global statistical distribution of those parameters as derived from literature values and databases. A Monte Carlo analysis of the model is then performed, in which parameters are stochastically selected from the pre-assigned PDFs. After several thousand iterations spanning the entire parameter space of physical and biogeochemical characteristics, a database of hypothetical worldwide N dynamics, including N₂O production and emissions, is generated for river, lake, reservoir, and estuarine systems. Then, global relationships relating N processes and N₂O emissions to TN and TP loads and water residence time are fitted from the database and applied for the global upscaling.

To calculate the cascading loads of TN and TP delivered to each water body along the river–reservoir–estuary continuum, we spatially routed all reservoirs from the GRand database⁷⁷, with river networks from Hydrosheds 15s⁷⁸ and, at latitudes above 50°N, Hydro1K (USGS, 2000), which were in turn connected to estuaries as represented in the “Worldwide Typology of Nearshore Coastal Systems” of Dürr et al.⁷⁹. In addition, the global data base HydroLAKES⁷⁶ was used to topologically connect 1.4 million lakes with a minimum surface area of 0.1 km² within the river network. Note that besides natural lakes, HydroLAKES includes updated information on 6,796 reservoirs from the GRand data base, which was used in the study of Maavara et al.¹⁰. In order to estimate the TN and TP loads to each water body, we then relied on a spatially explicit representation of TN and TP mobilization from the watershed into the river network (see Maavara et al. for details^{80,81}).

For the estimation of N₂O emission, we applied two distinct model configurations, respectively named DS1 and DS2 in Maavara et al.¹⁰. DS1 estimates N₂O emissions from denitrification and nitrification based on an EF of 0.9%, which is in the mean of published values⁸², and the assumption that N₂O production equals N₂O emissions¹⁰. For DS2, the reduction of N₂O to N₂ during denitrification if N₂O is not evading sufficiently rapidly from the water body is taken into account. The fluxes in the model represent lumped sediment–water column rates and were resolved at the annual timescale. The use of water residence time as independent variable in both the mechanistic model and the upscaling process introduces an important kinetic refinement to existing global N₂O emission estimates. Rather than applying an average EF (directly scaling N₂O emissions to N inputs) to all water bodies, the use of water

residence time explicitly adjusts for the extent of N₂O production and emission that is kinetically possible within the timeframe available in a given water body. Simulated N₂O emission rates were evaluated against measurement-based upscaling methods applied to reservoirs⁸³ and rivers⁸⁴ as well as against observation-driven regional estimates of lake N₂O emissions based on literature data⁹.

d. Coastal zone emissions

The average of net N₂O fluxes from three seagrass species²⁷ (seagrasses, mangroves, saltmarsh and intertidal) was scaled to the global seagrass area²⁸. The mangrove data from Murray et al.²⁸ was updated with water-air and sediment-air N₂O fluxes from Maher et al.⁸⁵ and Murray et al.⁸⁶. The average sediment-air N₂O flux and the average water-air N₂O flux were each applied for 12 hours a day (see Rosentreter et al.⁸⁷), and scaled to the global mangrove area²⁸. Murray et al.²⁸ saltmarsh data was updated with sediment-air N₂O fluxes from Yang et al.⁸⁸, Chmura et al.⁸⁹, Welti et al.⁹⁰ and Roughan et al.⁹¹ and scaled to the global saltmarsh area²⁸. Murray et al.²⁸ intertidal data was updated with sediment-air N₂O fluxes from Moseman-Valtierra et al.⁹² and Sun et al.⁹³ and scaled to the global intertidal area⁹⁴.

2.10. Atmospheric inversion models

Emissions were estimated using four independent atmospheric inversion frameworks (see Supplementary Table 19). The frameworks all used the Bayesian inversion method, which finds the optimal emissions, that is, those, which when coupled to a model of atmospheric transport, provide the best agreement to observed N₂O mixing ratios while being guided by the prior estimates and their uncertainty. In other words, the optimal emissions are those that minimize the cost function:

$$J(\mathbf{x}) = \frac{1}{2}(\mathbf{x} - \mathbf{x}_b)^T \mathbf{B}^{-1}(\mathbf{x} - \mathbf{x}_b) + \frac{1}{2}(\mathbf{y} - H(\mathbf{x}))^T \mathbf{R}^{-1}(\mathbf{y} - H(\mathbf{x})) \quad (5a)$$

where $\mathbf{x}$ and $\mathbf{x}_b$ are, respectively, vectors of the optimal and prior emissions, $\mathbf{B}$ is the prior error covariance matrix, $\mathbf{y}$ is a vector of observed N₂O mixing ratios, $\mathbf{R}$ is the observation error covariance matrix, and $H(\mathbf{x})$ is the model of atmospheric transport (for details on the inversion

method see⁹⁵). The optimal emissions, $\mathbf{x}$, were found by solving the first order derivative of equation (5a):

$$J'(\mathbf{x}) = \mathbf{B}^{-1}(\mathbf{x} - \mathbf{x}_b) + (H'(\mathbf{x}))^T \mathbf{R}^{-1}(\mathbf{y} - H(\mathbf{x})) = 0 \quad (5b)$$

where $(H'(\mathbf{x}))^T$ is the adjoint model of transport. In frameworks INVICAT, PyVAR and GEOSChem, equation (5b) was solved using the variational approach⁹⁶⁻⁹⁸, which uses a descent algorithm and computations involving the forward and adjoint models. In framework MIROC4-ACTM, equation (5b) was solved directly by computing a transport operator, $\mathbf{H}$ from integrations of the forward model, such that $\mathbf{H}\mathbf{x}$ is equivalent to $H(\mathbf{x})$, and taking the transpose of $\mathbf{H}$ ⁹⁹.

Each of the inversion frameworks used a different model of atmospheric transport with different horizontal and vertical resolutions (see Supplementary Table 19). The transport models TOMCAT and LMDz5, used in INVICAT and PyVAR respectively, were driven by ECMWF ERA-Interim wind fields, MIROC4-ACTM, was driven by JRA-55 wind fields, and GEOSChem was driven by MERRA-2 wind fields. While INVICAT, PyVAR, and GEOSChem optimized the emissions at the spatial resolution of the transport model, MIROC4-ACTM optimized the error in the emissions aggregated into 84 land and ocean regions. All frameworks optimized the emissions with monthly temporal resolution. The transport models included an online calculation of the loss of N₂O in the stratosphere due to photolysis and oxidation by O(¹D) resulting in mean atmospheric lifetimes of between 118 and 129 years, broadly consistent with recent independent estimates of the lifetime of 116±9 years (ref. ¹⁰⁰).

All inversions used N₂O measurements of discrete air samples from the National Oceanic and Atmospheric Administration Carbon Cycle Cooperative Global Air Sampling Network (NOAA). In addition, discrete measurements from the Commonwealth Scientific and Industrial Research Organisation network (CSIRO) as well as in-situ measurements from the Advanced Global Atmospheric Gases Experiment network (AGAGE), the NOAA CATS network, and from individual sites operated by University of Edinburgh (UE), National Institute for Environmental Studies (NIES) and the Finnish Meteorological Institute (FMI) were included in INVICAT, PyVAR and GEOSChem. Measurements from networks other than NOAA were corrected to the NOAA calibration scale, NOAA-2006A, using the results of the WMO Round Robin inter-

comparison experiment (<https://www.esrl.noaa.gov/gmd/ccgg/wmorrr/>), where available. For AGAGE and CSIRO, which did not participate in the WMO Round Robins, the data at sites where NOAA discrete samples are also collected were used to calculate a linear regression with NOAA data, which was applied to adjust the data to the NOAA-2006A scale. For the remaining CSIRO sites where there were no NOAA discrete samples, the mean regression coefficient and offset from all other CSIRO sites were used. The inversions used the discrete sample measurements without averaging, and hourly or daily means of the in-situ measurements, depending on the particular inversion framework.

Each framework applied its own method for calculating the observation space uncertainty, the square of which gives the diagonal elements of the observation error covariance matrix **R**. The observation space uncertainty accounts for measurement and model representation errors and is equal to the quadratic sum of these terms. Typical values for the observation space uncertainty were between 0.3 and 0.5 ppb for all inversion frameworks.

Prior emissions were based on estimates from terrestrial biosphere and ocean biogeochemistry models as well as from inventories. INVICAT, PyVAR and GEOSChem used the same prior estimates for emissions from natural and agricultural soils from the model OCN v1.1²² and for biomass burning emissions from GFEDv4.1s. For non-soil anthropogenic emissions (namely those from energy, industry and waste sectors), INVICAT and GEOSChem used EDGAR v4.2FT2010 and PyVAR used EDGAR v4.3.2. MIROC4-ACTM used the VISIT model^{24,25} for emissions from natural soils and EDGAR 4.2 for all anthropogenic emissions, including agricultural burning, but did not explicitly include a prior estimate for wildfire emissions.

Three different prior estimates for ocean emissions were used: 1) from the ocean biogeochemistry model, NEMO-PlankTOM5¹⁰¹, with a global total of 6.6 Tg N yr⁻¹, 2) from the updated version of this model, NEMO-PlankTOM10⁵ with a global total of 3.7 Tg N yr⁻¹, and 3) from the MIT ocean general circulation model, as described by Manizza et al.¹⁰² with a global total of 3.8 Tg N yr⁻¹.

Prior uncertainties were estimated in all the inversion frameworks for each grid cell (INVICAT, PyVAR and GEOSChem) or for each region (MIROC4-ACTM) and square of the uncertainties formed the diagonal elements of the prior error covariance matrix **B**. INVICAT, PyVAR and GEOSChem estimated the uncertainty as proportional to the prior value in each grid

cell, but MIROC4-ACTM set the uncertainty uniformly for the land regions at 1 Tg N yr⁻¹ and for the ocean regions at 0.5 Tg N yr⁻¹.

3. Atmospheric N₂O observations and growth rates for three different atmospheric networks (NOAA, AGAGE, and CISRO)

The monthly atmospheric N₂O abundances and their growth rates are derived from three different atmospheric observational networks (AGAGE, CISRO and NOAA) (Extended Data Fig. 1).

For atmospheric N₂O observations from the NOAA network¹⁰³, we used global mean mixing ratios from the GMD combined dataset during 1980–2017 based on measurements from five different measurement programs [HATS old flask instrument, HATS current flask instrument (OTTO), CCGG group Cooperative Global Air Sampling Network (<https://www.esrl.noaa.gov/gmd/ccgg/flask.php>), HATS *in situ* (RITS program), and HATS *in situ* (CATS program)]. CCGG provides uncertainties with each measurement (see site files: ftp://aftp.cmdl.noaa.gov/data/greenhouse_gases/n2o/flask/surface/). Global means are derived from flask and *in situ* measurements obtained by gas chromatography with electron capture detection, from 4–12 sites (fewer sites in the earlier years), weighted by representative area. Monthly mean observations from different NOAA measurement programs are statistically combined to create a long-term NOAA/ESRL GMD dataset. Uncertainties (1 sigma) associated with monthly estimates of global mean N₂O, are ~1 ppb from 1977–1987, 0.6 ppb from 1988–1994, 0.3–0.4 ppb from 1995–2000, and 0.1 ppb from 2001–2017. NOAA data are generally more consistent after 1995, with standard deviations on the monthly mean mixing ratios at individual sites of ~0.5 ppb from 1995–1998, and 0.1–0.4 ppb after 1998. A detailed description of these measurement programs and the method to combine them are available via <https://www.esrl.noaa.gov/gmd/hats/combined/N2O.html>.

The Advanced Global Atmospheric Gases Experiment (AGAGE) global network (and its predecessors ALE and GAGE)¹⁰⁴ has made continuous high frequency gas chromatographic measurements of N₂O at five globally distributed sites since 1978. AGAGE includes two types of instruments [i.e., a gas chromatograph with multiple detectors (GC-MD) and a gas

chromatograph with preconcentration and mass spectrometric analysis (Medusa GC-MS)]. The measurement precision for N₂O improved from about 0.35% in ALE to 0.13% in GAGE¹⁰⁵ and 0.05% in AGAGE¹⁰⁴. We used the global mean of N₂O measurements from the GC-MD during 1980–2017. Further information on AGAGE stations, instruments, calibration, uncertainties and access to data is available at the AGAGE website: <http://agage.mit.edu>.

The CSIRO flask network¹⁰⁶ consists of nine sampling sites distributed globally and has been in operation since 1992. Flask samples are collected approximately every two weeks and shipped back to CSIRO GASLAB for analysis. Samples were analyzed by gas chromatography with electron capture detection (GC-ECD). One Shimadzu gas chromatograph, labelled “Shimadzu-1” (S1) was used over the entire length of the record and the measurement precision for N₂O from this instrument is about 0.1%. N₂O data from the CSIRO global flask network are reported on the NOAA-2006A N₂O scale and are archived at the World Data Centre for Greenhouse Gases (WDCGG: <https://gaw.kishou.go.jp/>). Nine sites from the CSIRO network were used to calculate the annual global N₂O mole fractions. Smooth curve fits to the N₂O data from each of these sites were calculated using the technique outlined in Thoning et al.¹⁰⁷, using a short-term cut-off of 80 days. The smooth curve fit data were then placed on an evenly spaced latitude (5 degree) versus time (weekly) grid using the Kriging interpolation technique. Finally, the gridded data were used to calculate the global annual average mole fractions weighted by latitude.

We plotted the atmospheric globally averaged N₂O abundances and the associated growth rates for the three global atmospheric networks NOAA, AGAGE, and CSIRO during 1980–2017 (see Extended Data Fig. 1). We see remarkably consistent global mean N₂O estimated from the three different networks, increasing from 301.0±0.1 ppb in 1980 to 329.9±0.4 ppb in 2017. Growth rates of N₂O are also remarkably consistent among the three measurement networks. After a period in the late 1990s in which the growth rate averaged about 0.8 ppb yr⁻¹, the global growth rate fell to ~0.6 ppb yr⁻¹ and then gradually increased to nearly 1 ppb yr⁻¹ by 2013–2017. Interannual variability in the N₂O growth rate was higher prior to 1995 (not shown) than after 1995, which may be an artifact of less precise measurements due to changes in instrumental precision and measurement frequency over the study period. Additional discussion on uncertainties associated with measurement errors and emission errors in inversions can be found in Chen and Prinn¹⁰⁸ and Thompson et al¹⁶.

4. Comparison with the IPCC AR5

Our methodology significantly differs from past approaches summarized in the IPCC AR5. Most of the estimates used in the AR5 (e.g., natural sources) directly inherited or adopted with minor revisions data from studies conducted mainly in the 1990s. Some estimates used in the IPCC AR5 (e.g., atmospheric deposition on land) were from a review by Syakila and Kroeze¹⁰⁹, which depended on empirical methods and simple assumptions.

Compared to the findings reported in the IPCC AR5, our budget includes several new sources (e.g., aquaculture, deforestation/post-deforestation, the effects of environmental factors, natural sources of inland and coastal waters) and one additional (tropospheric) sink for N₂O (Table 6). We report natural sources of N₂O emissions from inland and coastal waters with a value of 0.3 Tg N yr⁻¹. The total source of N₂O in our study is 0.9 Tg N yr⁻¹ smaller than that in the IPCC AR5, while our estimate of anthropogenic N₂O emissions is 0.4 Tg N yr⁻¹ larger in the recent decade (Supplementary Table 15). Our larger estimate of anthropogenic emissions is associated with environmental effects (0.2, with a range of -0.6 to 1.1 Tg N yr⁻¹, based on NMIP simulations), and a 0.4 Tg N yr⁻¹ larger estimate of atmospheric N deposition emissions (based on modeling results and inventories, Table 1). In contrast, our estimate of direct emissions from agriculture [(3.8 (2.5–5.8) Tg N yr⁻¹, plus aquaculture, a minor contribution] is 0.3 Tg N yr⁻¹ smaller than reported in the IPCC AR5.

Natural sources in our study are 1.3 Tg N yr⁻¹ smaller than those reported by the IPCC AR5 for 2007–2016 and the range is significantly reduced. The mean NMIP estimate of global natural soil emission [5.6 (4.9–6.6) Tg N yr⁻¹] is 1.0 Tg N yr⁻¹ smaller compared to those in the IPCC AR5 estimate [6.6 (3.3–9.0) Tg N yr⁻¹]. The reduction in uncertainty in NMIP estimates may result from calibration of terrestrial biosphere models in NMIP against in situ observations across the globe¹, while the AR5 estimate, essentially inherited from the AR4 synthesis, was based on results from a single simple model by Bouwman et al.¹¹⁰.

In this study, global oceanic N₂O emission is derived from an ensemble of global ocean biogeochemistry models. Our estimate [3.4 (2.5–4.3) Tg N yr⁻¹] is 0.4 Tg N yr⁻¹ smaller and the uncertainty range is significantly smaller than reported in the IPCC AR5 (1.8–9.4 Tg N yr⁻¹). The larger AR5 range was determined using an analysis of Atlantic Ocean surface measurements (Rhee et al.¹¹¹; the Atlantic is not a region of significant N₂O emission) as the lower bound, and the upper bound was the maximal value of N₂O production from a global empirically based analysis¹¹². The parameterizations governing marine productivity and N₂O yield in our five ocean models have been constrained by a variety of datasets characterizing marine biogeochemical process rates, and the model simulations of ocean N₂O have been evaluated against global biogeochemical databases (e.g., see Battaglia and Joos³ and Buitenhuis et al.⁵ for more detail). The smaller range of ocean N₂O emission reported in this study includes advances in modeling such factors as quantification of global marine export production, improved constraints on N₂O yield parameters (particularly in the well-oxygenated ocean), and more comprehensive evaluation of modeled biogeochemical distributions.

The estimated N₂O production through atmospheric chemistry is 0.2 Tg N yr⁻¹ smaller than reported in the IPCC AR5. The observed stratospheric sink of N₂O in this study is 0.9 Tg N yr⁻¹ smaller than in the IPCC AR5, wherein stratospheric N₂O destruction was tuned to be consistent with the difference between the total source and the observed atmospheric N₂O growth rate. In our study, stratospheric sinks were obtained from atmospheric chemistry transport models and the recent post-AR5 study by Prather et al.^{16,100,113} who calculated N₂O stratospheric loss (& lifetime) based on satellite observations combined with simple photolysis models using observed atmospheric temperature, O₂, and O₃. Our uncertainties in the atmospheric loss of N₂O (±1.1 Tg N yr⁻¹) are slightly larger than those of the AR5 (±0.9 Tg N yr⁻¹). In our study, annual change in atmospheric abundance is calculated from the combined NOAA and AGAGE record of surface N₂O and uncertainty (±0.5 Tg N yr⁻¹) is taken from the IPCC AR5 (ref. ¹¹⁴).

5. Per capita N₂O emission at global and regional scales in the recent decade

Per capita N₂O emission is calculated using global and regional emissions divided by the numbers of global and regional population¹¹⁵ (see Supplementary Fig. 2). Global per capita emissions from top-down and bottom-up approaches were on average ~2 kg N capita⁻¹ yr⁻¹ in the

recent decade. Bottom-up estimates show that per capita natural fluxes including natural soils and inland and coastal waters were the largest source, followed by agriculture and other direct anthropogenic sources. South America and Oceania have ~2 times and ~6 times higher per capita emissions than the global average, respectively. Africa and Russia also have higher per capita N₂O emissions than the global value contributed primarily by natural fluxes and to a minor extent by other direct anthropogenic sources (Africa: Biomass burning; Russia: Fossil fuel and industry and Biomass burning). In addition, North America and Europe show higher than global per capita emissions from agriculture and other direct anthropogenic sources (primary from Fossil fuel and industry). Middle East, East Asia, South Asia, and Southeast Asia show lower than global per capita emissions from all sources.

References:

- 1 Tian, H. Q. *et al.* Global soil nitrous oxide emissions since the preindustrial era estimated by an ensemble of terrestrial biosphere models: Magnitude, attribution, and uncertainty. *Global Change Biology* **25**, 640-659 (2019).
- 2 Dangal, S. R. *et al.* Global nitrous oxide emissions from pasturelands and rangelands: Magnitude, spatio-temporal patterns and attribution. *Global Biogeochemical Cycles* **33**, 200-222 (2019).
- 3 Battaglia, G. & Joos, F. Marine N₂O Emissions From Nitrification and Denitrification Constrained by Modern Observations and Projected in Multimillennial Global Warming Simulations. *Global Biogeochemical Cycles* **32**, 92-121 (2018).
- 4 Berthet, S. *et al.* Evaluation of an Online Grid-Coarsening Algorithm in a Global Eddy-Admitting Ocean Biogeochemical Model. *Journal of Advances in Modeling Earth Systems* **11**, 1759-1783 (2019).
- 5 Buitenhuis, E. T., Suntharalingam, P. & Le Quéré, C. Constraints on global oceanic emissions of N₂O from observations and models. *Biogeosciences* **15**, 2161-2175 (2018).
- 6 Landolfi, A., Somes, C. J., Koeve, W., Zamora, L. M. & Oschlies, A. Oceanic nitrogen cycling and N₂O flux perturbations in the Anthropocene. *Global Biogeochemical Cycles* **31**, 1236-1255 (2017).
- 7 Martinez-Rey, J., Bopp, L., Gehlen, M., Tagliabue, A. & Gruber, N. Projections of oceanic N₂O emissions in the 21st century using the IPSL Earth system model. *Biogeosciences* **12**, 4133-4148 (2015).
- 8 Suntharalingam, P. *et al.* Estimates of Oceanic Nitrous-oxide Emissions from Global Biogeochemistry Models. *American Geophysical Union, Fall Meeting 2018* (2018).
- 9 Lauerwald, R. *et al.* Natural lakes are a minor global source of N₂O to the atmosphere. *Global Biogeochemical Cycles* **33**, 1564-1581 (2019).
- 10 Maavara, T. *et al.* Nitrous oxide emissions from inland waters: Are IPCC estimates too high? *Global Change Biology* **25**, 473-488 (2019).
- 11 Janssens-Maenhout, G. *et al.* EDGAR v4.3.2 Global Atlas of the three major greenhouse gas emissions for the period 1970–2012. *Earth System Science Data* **11**, 959-1002 (2019).
- 12 Tubiello, F. *et al.* Estimating greenhouse gas emissions in agriculture: a manual to address data requirements for developing countries. FAO, Rome (2015).
- 13 Winiwarter, W., Höglund-Isaksson, L., Klimont, Z., Schöpp, W. & Amann, M. Technical opportunities to reduce global anthropogenic emissions of nitrous oxide. *Environmental research letters* **13**, 014011 (2018).
- 14 Van Der Werf, G. R. *et al.* Global fire emissions estimates during 1997-2016. *Earth System Science Data* **9**, 697-720 (2017).
- 15 Wang, Q. *et al.* Data-driven estimates of global nitrous oxide emissions from croplands. *National Science Review* **00**, 1-12 (2019).
- 16 Thompson, R. L. *et al.* Acceleration of global N₂O emissions seen from two decades of atmospheric inversion. *Natural Climate Change* **9**, 993-998 (2019).
- 17 Tian, H. Q. *et al.* The Global N₂O Model Intercomparison Project. *Bulletin of the American Meteorological Society* **99**, 1231-1252 (2018).
- 18 Tian, H. *et al.* Global methane and nitrous oxide emissions from terrestrial ecosystems due to multiple environmental changes. *Ecosystem Health and Sustainability* **1**, 1-20 (2015).
- 19 Xu, R. *et al.* Preindustrial nitrous oxide emissions from the land biosphere estimated by using a global biogeochemistry model. *Climate of the Past* **13**, 977 (2017).
- 20 Stocker, B. D. *et al.* Multiple greenhouse-gas feedbacks from the land biosphere under future climate change scenarios. *Nature Climate Change* **3**, 666, doi:10.1038/nclimate1864 (2013).
- 21 XU-RI & PRENTICE, I. C. Terrestrial nitrogen cycle simulation with a dynamic global vegetation model. *Global Change Biology* **14**, 1745-1764 (2008).

- 22 Zaehle, S., Ciais, P., Friend, A. D. & Prieur, V. Carbon benefits of anthropogenic reactive nitrogen offset by nitrous oxide emissions. *Nature Geoscience* **4**, 601-605 (2011).
- 23 Goll, D. S. *et al.* A representation of the phosphorus cycle for ORCHIDEE (revision 4520). *Geosci. Model Dev.* **10**, 3745-3770 (2017).
- 24 Inatomi, M., Ito, A., Ishijima, K. & Murayama, S. Greenhouse Gas Budget of a Cool-Temperate Deciduous Broad-Leaved Forest in Japan Estimated Using a Process-Based Model. *Ecosystems* **13**, 472-483 (2010).
- 25 Ito, A., Nishina, K., Ishijima, K., Hashimoto, S. & Inatomi, M. Emissions of nitrous oxide (N₂O) from soil surfaces and their historical changes in East Asia: a model-based assessment. *Progress in Earth and Planetary Science* **5**, 55, doi:10.1186/s40645-018-0215-4 (2018).
- 26 Yao, Y. *et al.* Increased global nitrous oxide emissions from streams and rivers in the Anthropocene. *Natural Climate Change* **10**, 138-142 (2020).
- 27 Camillini, N., Rossentreter, J., Erler, D., Glud, R. N. & Eyre, B. Blue carbon burial in seagrass meadows is partially offset by CH₄ and N₂O fluxes. *Science Advance* (Submitted).
- 28 Murray, R. H., Erler, D. V. & Eyre, B. D. Nitrous oxide fluxes in estuarine environments: response to global change. *Global Change Biology* **21**, 3219-3245 (2015).
- 29 Beusen, A. H., Bouwman, A. F., Van Beek, L. P., Mogollón, J. M. & Middelburg, J. J. Global riverine N and P transport to ocean increased during the 20th century despite increased retention along the aquatic continuum. *Biogeosciences* **13**, 2441-2451 (2016).
- 30 Bouwman, A. F. *et al.* Hindcasts and future projections of global inland and coastal nitrogen and phosphorus loads due to finfish aquaculture. *Reviews in Fisheries Science* **21**, 112-156 (2013).
- 31 Bouwman, A. F. *et al.* Global hindcasts and future projections of coastal nitrogen and phosphorus loads due to shellfish and seaweed aquaculture. *Reviews in Fisheries Science* **19**, 331-357 (2011).
- 32 Tubiello, F. N. *et al.* The FAOSTAT database of greenhouse gas emissions from agriculture. **8**, 015009 (2013).
- 33 Davidson, E. A., Erickson, H. E., Verchot, L. V., Keller, M. & Veldkamp, E. Testing a Conceptual Model of Soil Emissions of Nitrous and Nitric Oxides: Using two functions based on soil nitrogen availability and soil water content, the hole-in-the-pipe model characterizes a large fraction of the observed variation of nitric oxide and nitrous oxide emissions from soils. *BioScience* **50**, 667-680 (2000).
- 34 IPCC. 2006 IPCC Guidelines for National Greenhouse Gas Inventories., (Japan on behalf of the IPCC, Hayama, Japan, 2006).
- 35 Zhou, J., Jiang, M. & Chen, G. Estimation of methane and nitrous oxide emission from livestock and poultry in China during 1949–2003. *Energy Policy* **35**, 3759-3767 (2007).
- 36 Van Drecht, G., Bouwman, A. F., Knoop, J. M., Beusen, A. H. W. & Meinardi, C. R. Global modeling of the fate of nitrogen from point and nonpoint sources in soils, groundwater, and surface water. *Global Biogeochemical Cycles* **17**, 1115, doi:10.1029/2003gb002060 (2003).
- 37 Amann, M. *et al.* Cost-effective control of air quality and greenhouse gases in Europe: Modeling and policy applications. *Environmental Modelling & Software* **26**, 1489-1501 (2011).
- 38 Zhang, X. *et al.* Managing nitrogen for sustainable development. *Nature* **528**, 51, doi:doi:10.1038/nature15743 (2015).
- 39 Gao, S. *et al.* Quantifying nitrogen leaching response to fertilizer additions in China's cropland. *Environmental pollution* **211**, 241-251 (2016).
- 40 Hou, X. *et al.* Detection and attribution of nitrogen runoff trend in China's croplands. *Environmental pollution* **234**, 270-278 (2018).
- 41 Zhou, F. *et al.* New model for capturing the variations of fertilizer-induced emission factors of N₂O. *Global Biogeochemical Cycles* **29**, 885-897 (2015).
- 42 Decock, C. Mitigating nitrous oxide emissions from corn cropping systems in the midwestern US: potential and data gaps. *Environmental science & technology* **48**, 4247-4256 (2014).

- 43 Harris, I., Jones, P. D., Osborn, T. J. & Lister, D. H. Updated high-resolution grids of monthly climatic observations—the CRU TS3. 10 Dataset. *International journal of climatology* **34**, 623-642 (2014).
- 44 Helgason, B. *et al.* Toward improved coefficients for predicting direct N₂O emissions from soil in Canadian agroecosystems. *Nutrient Cycling in Agroecosystems* **72**, 87-99 (2005).
- 45 Hénault, C. *et al.* Predicting in situ soil N₂O emission using NOE algorithm and soil database. *Global Change Biology* **11**, 115-127 (2005).
- 46 Hickman, J. E., Scholes, R. J., Rosenstock, T. S., Garcia-Pando, C. P. & Nyamangara, J. Assessing non-CO₂ climate-forcing emissions and mitigation in sub-Saharan Africa. *Current Opinion in Environmental Sustainability* **9**, 65-72 (2014).
- 47 Kim, D.-G., Giltrap, D. & Hernandez-Ramirez, G. Background nitrous oxide emissions in agricultural and natural lands: a meta-analysis. *Plant and soil* **373**, 17-30 (2013).
- 48 Kim, D.-G., Hernandez-Ramirez, G. & Giltrap, D. Linear and nonlinear dependency of direct nitrous oxide emissions on fertilizer nitrogen input: A meta-analysis. *Agriculture, Ecosystems & Environment* **168**, 53-65 (2013).
- 49 Lehuger, S. *et al.* Predicting and mitigating the net greenhouse gas emissions of crop rotations in Western Europe. *Agricultural and forest meteorology* **151**, 1654-1671 (2011).
- 50 Leppelt, T. *et al.* Nitrous oxide emission budgets and land-use-driven hotspots for organic soils in Europe. *Biogeosciences* (2014).
- 51 Rochette, P. & Janzen, H. H. Towards a revised coefficient for estimating N₂O emissions from legumes. *Nutrient Cycling in Agroecosystems* **73**, 171-179 (2005).
- 52 Sacks, W. J., Deryng, D., Foley, J. A. & Ramankutty, N. Crop planting dates: an analysis of global patterns. *Global Ecol Biogeogr* **19**, 607-620 (2010).
- 53 Shcherbak, I., Millar, N. & Robertson, G. P. Global metaanalysis of the nonlinear response of soil nitrous oxide (N₂O) emissions to fertilizer nitrogen. *Proceedings of the National Academy of Sciences* **111**, 9199-9204 (2014).
- 54 Stehfest, E. & Bouwman, L. N₂O and NO emission from agricultural fields and soils under natural vegetation: summarizing available measurement data and modeling of global annual emissions. *Nutrient Cycling in Agroecosystems* **74**, 207-228 (2006).
- 55 Walter, K. *et al.* Direct nitrous oxide emissions from oilseed rape cropping—a meta-analysis. *GCB Bioenergy* **7**, 1260-1271 (2015).
- 56 Berdanier, A. B. & Conant, R. T. Regionally differentiated estimates of cropland N₂O emissions reduce uncertainty in global calculations. *Global change biology* **18**, 928-935 (2012).
- 57 Yang, H. *et al.* Regional patterns of future runoff changes from Earth system models constrained by observation. *Geophysical Research Letters* **44**, 5540-5549 (2017).
- 58 Goldewijk, K. K., Beusen, A., Doelman, J. & Stehfest, E. Anthropogenic land use estimates for the Holocene—HYDE 3.2. *Earth System Science Data* **9**, 927-953 (2017).
- 59 Zhang, B. *et al.* Global manure nitrogen production and application in cropland during 1860–2014: a 5 arcmin gridded global dataset for Earth system modeling. *Earth System Science Data* **9**, 667-678 (2017).
- 60 Rosas, F. Fertilizer use by crop at the country level (1990–2010). *CARD Working Papers* **555** (2012).
- 61 Carlson, K. M. *et al.* Greenhouse gas emissions intensity of global croplands. *Nature Climate Change* **7**, 63 (2017).
- 62 Bouwman, L. *et al.* Mariculture: significant and expanding cause of coastal nutrient enrichment. *Environmental Research Letters* **8**, 044026, doi:044010.041088/041748-049326/044028/044024/044026 (2013).
- 63 Bouwman, A. F. *et al.* Hindcasts and future projections of global inland and coastal nitrogen and phosphorus loads due to finfish aquaculture. *Reviews in Fisheries Science* **21**, 112-156 (2013).
- 64 Bouwman, A. F. *et al.* Global hindcasts and future projections of coastal nitrogen and phosphorus loads due to shellfish and seaweed aquaculture. *Reviews in Fisheries Science* **19**, 331-357 (2011).

- 65 FAO. FishStatJ - Software for Fishery and Aquaculture Statistical Time Series.
<http://www.fao.org/fishery/statistics/software/fishstatj/en>. (Fisheries and Aquaculture Information and Statistics Service, Food and Agriculture Organization of the United Nations, retrieved 22 May 2019, Rome, 2019).
- 66 Sullivan, B. W. *et al.* Biogeochemical recuperation of lowland tropical forest during succession. *Ecology* **100**, e02641, doi:10.1002/ecy.2641 (2019).
- 67 Davidson, E. A. *et al.* Recuperation of nitrogen cycling in Amazonian forests following agricultural abandonment. *Nature* **447**, 995 (2007).
- 68 Keller, M. & Reiners, W. A. Soil-atmosphere exchange of nitrous oxide, nitric oxide, and methane under secondary succession of pasture to forest in the Atlantic lowlands of Costa Rica. *Global Biogeochemical Cycles* **8**, 399-409 (1994).
- 69 Liu, M. *et al.* Long-term trends in evapotranspiration and runoff over the drainage basins of the Gulf of Mexico during 1901–2008. *Water Resources Research* **49**, 1988-2012 (2013).
- 70 Ren, W. *et al.* Century-long increasing trend and variability of dissolved organic carbon export from the Mississippi River basin driven by natural and anthropogenic forcing. *Global Biogeochemical Cycles* **30**, 1288-1299 (2016).
- 71 Ren, W. *et al.* Large increase in dissolved inorganic carbon flux from the Mississippi River to Gulf of Mexico due to climatic and anthropogenic changes over the 21st century. *Journal of Geophysical Research: Biogeosciences* **120**, 724-736 (2015).
- 72 Tao, B. *et al.* Increasing Mississippi river discharge throughout the 21st century influenced by changes in climate, land use, and atmospheric CO₂. *Geophysical Research Letters* **41**, 4978-4986 (2014).
- 73 Tian, H. *et al.* Climate extremes dominating seasonal and interannual variations in carbon export from the Mississippi River Basin. *Global Biogeochemical Cycles* **29**, 1333-1347 (2015).
- 74 Tian, H. *et al.* Anthropogenic and climatic influences on carbon fluxes from eastern North America to the Atlantic Ocean: A process-based modeling study. *Journal of Geophysical Research: Biogeosciences* **120**, 757-772 (2015).
- 75 Yang, Q. *et al.* Increased nitrogen export from eastern North America to the Atlantic Ocean due to climatic and anthropogenic changes during 1901–2008. *Journal of Geophysical Research: Biogeosciences* **120**, 1046-1068 (2015).
- 76 Messenger, M. L., Lehner, B., Grill, G., Nedeva, I. & Schmitt, O. Estimating the volume and age of water stored in global lakes using a geo-statistical approach. *Nature Communications* **7**, 13603 (2016).
- 77 Lehner, B. *et al.* High-resolution mapping of the world's reservoirs and dams for sustainable river-flow management. *Frontiers in Ecology and the Environment* **9**, 494-502 (2011).
- 78 Lehner, B., Verdin, K. & Jarvis, A. New global hydrography derived from spaceborne elevation data. *Eos, Transactions, American Geophysical Union* **89**, 93-94 (2008).
- 79 Dürr, H. H. *et al.* Worldwide Typology of Nearshore Coastal Systems: Defining the Estuarine Filter of River Inputs to the Oceans. *Estuaries and Coasts* **34**, 441-458 (2011).
- 80 Bouwman, A., Beusen, A. H. & Billen, G. J. G. B. C. Human alteration of the global nitrogen and phosphorus soil balances for the period 1970–2050. *Global Biogeochemical Cycles* **23**, GB0A04, doi:10.1029/2009GB003576 (2009).
- 81 Van Drecht, G., Bouwman, A., Harrison, J. & Knoop, J. Global nitrogen and phosphate in urban wastewater for the period 1970 to 2050. *Global Biogeochemical Cycles* **23**, GB0A03 (2009).
- 82 Beaulieu, J. J. *et al.* Nitrous oxide emission from denitrification in stream and river networks. *Proceedings of the National Academy of Sciences* **108**, 214-219 (2011).
- 83 Deemer, B. R. *et al.* Greenhouse gas emissions from reservoir water surfaces: a new global synthesis. **66**, 949-964 (2016).
- 84 Hu, M., Chen, D. & Dahlgren, R. A. Modeling nitrous oxide (N₂O) emission from rivers: A global assessment. *Global Change Biology* **22**, 3566-3582 (2016).

- 85 Maher, D. T., Sippo, J. Z., Tait, D. R., Holloway, C. & Santos, I. R. Pristine mangrove creek
waters are a sink of nitrous oxide. *Scientific reports* **6**, 25701 (2016).
- 86 Murray, R., Erler, D., Rosentreter, J., Maher, D. & Eyre, B. A seasonal source and sink of nitrous
oxide in mangroves: Insights from concentration, isotope, and isotopomer measurements.
Geochimica et Cosmochimica Acta **238**, 169-192 (2018).
- 87 Rosentreter, J. A., Maher, D. T., Erler, D. V., Murray, R. H. & Eyre, B. D. Methane emissions
partially offset “blue carbon” burial in mangroves. *Science advances* **4**, eaao4985 (2018).
- 88 Yang, W. H. & Silver, W. L. Gross nitrous oxide production drives net nitrous oxide fluxes
across a salt marsh landscape. *Global Change Biology* **22**, 2228-2237 (2016).
- 89 Chmura, G. L., Kellman, L., Van Ardenne, L. & Guntenspergen, G. R. Greenhouse gas fluxes
from salt marshes exposed to chronic nutrient enrichment. *PloS one* **11**, e0149937 (2016).
- 90 Welti, N., Hayes, M. & Lockington, D. Seasonal nitrous oxide and methane emissions across a
subtropical estuarine salinity gradient. *Biogeochemistry* **132**, 55-69 (2017).
- 91 Roughan, B. L., Kellman, L., Smith, E. & Chmura, G. L. Nitrous oxide emissions could reduce
the blue carbon value of marshes on eutrophic estuaries. *Environmental Research Letters* **13**,
044034 (2018).
- 92 Moseman-Valtierra, S. *et al.* Substantial nitrous oxide emissions from intertidal sediments and
groundwater in anthropogenically-impacted West Falmouth Harbor, Massachusetts.
Chemosphere **119**, 1281-1288 (2015).
- 93 Sun, W., Sun, Z., Mou, X., Sun, W. & Hu, X. Nitrous oxide emissions from intertidal zone of the
Yellow River estuary in autumn and winter during 2011–2012. *Estuaries and coasts* **40**, 145-159
(2017).
- 94 Murray, N. J. *et al.* The global distribution and trajectory of tidal flats. *Nature* **565**, 222 (2019).
- 95 Tarantola, A. *Inverse problem theory and methods for model parameter estimation*. Vol. 89
(Society for Industrial and Applied Mathematics 2005).
- 96 Thompson, R. L. *et al.* Nitrous oxide emissions 1999 to 2009 from a global atmospheric
inversion. *Atmospheric Chemistry and Physics* **14**, 1801-1817 (2014).
- 97 Wells, K. C. *et al.* Simulation of atmospheric N₂O with GEOS-Chem and its adjoint: evaluation
of observational constraints. *Geoscience Model Development* **8**, 3179-3198 (2015).
- 98 Wilson, C., Chipperfield, M., Gloor, M. & Chevallier, F. Development of a variational flux
inversion system (INVICAT v1. 0) using the TOMCAT chemical transport model. *Geoscientific
Model Development* **7**, 2485-2500 (2014).
- 99 Patra, P. K. *et al.* Improved Chemical Tracer Simulation by MIROC4. 0-based Atmospheric
Chemistry-Transport Model (MIROC4-ACTM). *Sola* **14**, 91-96 (2018).
- 100 Prather, M. J. *et al.* Measuring and modeling the lifetime of nitrous oxide including its variability.
Journal of Geophysical Research: Atmospheres **120**, 5693-5705 (2015).
- 101 Suntharalingam, P. *et al.* Quantifying the impact of anthropogenic nitrogen deposition on oceanic
nitrous oxide. *Geophysical Research Letters* **39**, L07605, doi:10.1029/2011GL050778 (2012).
- 102 Manizza, M., Keeling, R. F. & Nevison, C. D. On the processes controlling the seasonal cycles of
the air–sea fluxes of O₂ and N₂O: A modelling study. *Tellus B: Chemical and Physical
Meteorology* **64**, 18429, doi:10.3402/tellusb.v64i0.18429 (2012).
- 103 Hall, B., Dutton, G. & Elkins, J. The NOAA nitrous oxide standard scale for atmospheric
observations. *Journal of Geophysical Research: Atmospheres* **112**, D09305,
doi:10.1029/2006JD007954 (2007).
- 104 Prinn, R. G. *et al.* History of chemically and radiatively important atmospheric gases from the
Advanced Global Atmospheric Gases Experiment (AGAGE). *Earth System Science Data* **10**,
985-1018 (2018).
- 105 Prinn, R. *et al.* Atmospheric emissions and trends of nitrous oxide deduced from 10 years of
ALE–GAGE data. *Journal of Geophysical Research: Atmospheres* **95**, 18369-18385 (1990).
- 106 Francey, R. J. *et al.* The CSIRO (Australia) measurement of greenhouse gases in the global
atmosphere. 97-111 (World Meteorological Organization, Tokyo, Japan, 2003).

- 107 Thoning, K. W., Tans, P. P. & Komhyr, W. D. Atmospheric carbon dioxide at Mauna Loa Observatory: 2. Analysis of the NOAA GMCC data, 1974–1985. *Journal of Geophysical Research: Atmospheres* **94**, 8549-8565 (1989).
- 108 Chen, Y. H. & Prinn, R. G. Estimation of atmospheric methane emissions between 1996 and 2001 using a three-dimensional global chemical transport model. *Journal of Geophysical Research: Atmospheres* **111**, 307, doi:10.1029/2005JD006058 (2006).
- 109 Syakila, A. & Kroeze, C. The global nitrous oxide budget revisited. *Greenhouse Gas Measurement and Management* **1**, 17-26 (2011).
- 110 Bouwman, A., Fung, I., Matthews, E. & John, J. Global analysis of the potential for N₂O production in natural soils. *Global Biogeochemical Cycles* **7**, 557-597 (1993).
- 111 Rhee, T. S., Kettle, A. J. & Andreae, M. O. Methane and nitrous oxide emissions from the ocean: A reassessment using basin-wide observations in the Atlantic. *Journal of Geophysical Research: Atmospheres* **114**, doi:10.1029/2008jd011662 (2009).
- 112 Bianchi, D., Dunne, J. P., Sarmiento, J. L. & Galbraith, E. D. Data-based estimates of suboxia, denitrification, and N₂O production in the ocean and their sensitivities to dissolved O₂. *Global Biogeochemical Cycles* **26**, doi:10.1029/2011gb004209 (2012).
- 113 Prather, M. J. & Hsu, J. Coupling of Nitrous Oxide and Methane by Global Atmospheric Chemistry. *Science* **330**, 952-954 (2010).
- 114 Ciais, P. *et al.* in *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* 465-570 (Cambridge University Press, 2014).
- 115 FAOSTAT. *Food and Agriculture Organization of the United Nations Statistics-Population*, (2020).
- 116 Hu, Z., Lee, J. W., Chandran, K., Kim, S. & Khanal, S. K. Nitrous oxide (N₂O) emission from aquaculture: a review. *Environmental science technology* **46**, 6470-6480 (2012).
- 117 MacLeod, M., Hasan, M. R., Robb, D. H. F. & Mamun-Ur-Rashid, M. Quantifying and mitigating greenhouse gas emissions from global aquaculture. FAO, Rome, (2019).

Supplementary Table 1 N₂O emissions from global agricultural soils based on multiple bottom-up approaches including the additions of mineral N fertilizer, manure and crop residues, and cultivation of organic soils. Unit: Tg N yr⁻¹

Data sources		1980s	1990s	2000s	2007-2016
Process-based models	NMIP/DLEM Mean	1.7	2.1	2.4	2.8
	NMIP/DLEM Min	0.9	1.1	1.3	1.4
	NMIP/DLEM Max	2.6	3.1	3.4	3.8
Statistical model plus DLEM	SRNM/DLEM	1.3	1.6	1.9	2.1
	EDGAR v4.3.2	1.3	1.5	1.7	1.9
Inventories	GAINS	1.5	1.6	1.7	1.9
	FAOSTAT	1.2	1.5	1.7	1.9
Mean		1.5	1.8	2.0	2.3
Min		0.9	1.1	1.3	1.4
Max		2.6	3.1	3.4	3.8

Supplementary Table 2 N₂O emissions from global total area under permanent meadows and pasture, due to manure N deposition (left on pasture) based on EDGAR v4.3.2, FAOSTAT, and GAINS estimates. Unit: Tg N yr⁻¹

Data sources	1980s	1990s	2000s	2007-2016
EDGAR v4.3.2	1.0	1.1	1.2	1.3
GAINS	0.7	0.7	0.8	0.9
FAOSTAT	1.0	1.1	1.2	1.3
Mean	0.9	1.0	1.1	1.2
Min	0.7	0.7	0.8	0.9
Max	1.0	1.1	1.2	1.3

Supplementary Table 3 N₂O emissions due to global manure management based on multiple bottom-up approaches. Unit: Tg N yr⁻¹

Data sources	1980s	1990s	2000s	2007-2016
EDGAR v4.3.2	0.2	0.2	0.2	0.2
GAINS	0.4	0.4	0.5	0.5
FAOSTAT	0.2	0.2	0.2	0.2
Mean	0.3	0.3	0.3	0.3
Min	0.2	0.2	0.2	0.2
Max	0.4	0.4	0.5	0.5

Supplementary Table 4 Aquaculture N₂O emissions based on multiple sources. Unit: Tg N yr⁻¹

Data sources	Emission factor (%)	1980s	1990s	2000s	2007-2016*
Hu et al. ¹¹⁶	1.8	N/A	N/A	N/A	0.1
MacLeod et al. ¹¹⁷	1.8	N/A	N/A	N/A	0.1
Bouwman et al.	1.8	0.01	0.03	0.1	0.1
Bouwman et al. _Min	0.5	0.00	0.01	0.02	0.02
Bouwman et al. _Max	5.0	0.03	0.1	0.2	0.2

* Estimates in Hu et al.¹¹⁶ and Macleod et al.¹¹⁷ were in 2009 and 2013, respectively. N/A represents data are not available.

Supplementary Table 5 Anthropogenic N₂O emissions from the global inland waters based on process-based models. Unit: Tg N yr⁻¹

Data sources/sectors	1980s	1990s	2000s	2007-2016
River_DLEM	0.2	0.2	0.2	0.2
River_Maavara	0.03	0.03	0.03	0.03
River_Mean	0.1	0.1	0.1	0.1
Reservoirs_DLEM	0.05	0.05	0.05	0.05
Reservoirs_Maavara	0.03	0.03	0.03	0.03
Reservoirs_Mean	0.04	0.04	0.04	0.04
Estuaries_Maavara	0.1	0.1	0.1	0.1
Lake_Lauerwald	0.02	0.02	0.02	0.02
Blue carbon_Murray	0.1	0.1	0.1	0.1
Total_Mean	0.3	0.3	0.3	0.3
Total_Min	0.2	0.2	0.2	0.2
Total_Max	0.4	0.4	0.4	0.4

Supplementary Table 6 Anthropogenic N₂O emissions from the global inland waters based on multiple bottom-up approaches. Unit: Tg N yr⁻¹

Data sources	1980s	1990s	2000s	2007-2016
FAOSTAT	0.4	0.4	0.5	0.6
GAINS	0.4	0.4	0.5	0.6
EDGAR v4.3.2	0.5	0.5	0.6	0.7
Model-based	0.3	0.3	0.3	0.3
Mean	0.4	0.4	0.4	0.5
Min	0.2	0.2	0.2	0.2
Max	0.5	0.5	0.6	0.7

Supplementary Table 7 Natural N₂O emissions from the global inland waters based on process-based models. Unit: Tg N yr⁻¹

Data sources/sectors	1980s	1990s	2000s	2007-2016
River_DLEM	0.1	0.1	0.1	0.1
River_Maavara	0.02	0.02	0.02	0.02
River_Mean	0.04	0.04	0.04	0.04
Reservoirs_DLEM	0.04	0.04	0.04	0.04
Reservoirs_Maavara	0.03	0.03	0.03	0.03
Reservoirs_Mean	0.03	0.03	0.03	0.03
Estuaries_Maavara	0.05	0.05	0.05	0.05
Lake_Lauerwald	0.02	0.02	0.02	0.02
Blue carbon_Murray	0.2	0.2	0.2	0.2
Total_Mean	0.3	0.3	0.3	0.3
Total_Min	0.3	0.3	0.3	0.3
Total_Max	0.4	0.4	0.4	0.4

Supplementary Table 8 Nitrous oxide emissions due to atmospheric N deposition on land based on multiple bottom-up approaches. Unit: Tg N yr⁻¹

Data sources	1980s	1990s	2000s	2007-2016
EDGAR v4.3.2	0.3	0.3	0.3	0.4
FAOSTAT	0.3	0.3	0.3	0.4
GAINS	0.3	0.3	0.3	0.4
FAOSTAT/EDGAR v4.3.2	0.6	0.6	0.7	0.8
GAINS/EDGAR v4.3.2	0.6	0.6	0.7	0.7
NMIP_Mean	0.6	0.7	0.8	0.8
NMIP_Min	0.3	0.4	0.4	0.4
NMIP_Max	1.2	1.4	1.3	1.4
Mean	0.6	0.7	0.7	0.8
Min	0.3	0.4	0.4	0.4
Max	1.2	1.4	1.3	1.4

Supplementary Table 9 Global N₂O emissions from waste and waste water based on EDGAR v4.3.2 and GAINS estimates. Unit: Tg N yr⁻¹

Data sources	1980s	1990s	2000s	2007-2016
EDGAR v4.3.2	0.1	0.2	0.2	0.2
GAINS	0.3	0.4	0.4	0.5
Mean	0.2	0.3	0.3	0.3
Min	0.1	0.2	0.2	0.2
Max	0.3	0.4	0.4	0.5

Supplementary Table 10 Global N₂O emissions from fossil fuel and industry based on multiple bottom-up approaches. Unit: Tg N yr⁻¹

Data sources	Sectors	1980s	1990s	2000s	2007-2016
EDGAR v4.3.2	Energy	0.1	0.1	0.1	0.2
	Transportation	0.2	0.2	0.2	0.2
	Others_residential	0.1	0.2	0.2	0.2
	Industry	0.7	0.5	0.5	0.5
GAINS	Energy	0.3	0.4	0.5	0.5
	Industry	0.5	0.5	0.4	0.3
	Mean	0.9	0.9	0.9	1.0
	Min	0.8	0.9	0.8	0.8
	Max	1.1	1.0	1.0	1.1

Supplementary Table 11 Global N₂O emissions from biomass burning based on multiple bottom-up approaches.
Unit: Tg N yr⁻¹

Fire categories	Data sources	1980s	1990s	2000s	2007-2016
Crop residues and savannas	GFED4s		0.4	0.4	0.4
	FAOSTAT		0.4	0.4	0.3
	DLEM	0.3	0.3	0.3	0.4
	Mean	0.3	0.4	0.4	0.4
Tropical forests and Deforestation*	GFED4s		0.1	0.1	0.1
	FAOSTAT		0.1	0.1	0.1
	DLEM	0.2	0.2	0.2	0.2
	Mean	0.2	0.1	0.1	0.1
Peatland	GFED4s		0.04	0.01	0.01
	FAOSTAT		0.1	0.1	0.1
	DLEM	0.04	0.05	0.02	0.02
	Mean	0.04	0.06	0.04	0.04
Boreal and temperate forests	GFED4s		0.1	0.1	0.1
	FAOSTAT		0.1	0.1	0.1
	DLEM	0.1	0.1	0.1	0.1
	Mean	0.1	0.1	0.1	0.1
Total_Mean		0.7	0.7	0.6	0.6
Total_Min		0.7	0.5	0.4	0.5
Total_Max		0.7	0.9	0.8	0.8

* DLEM estimates represent burning of tropical forests that are caused by natural and deforestation fires.

Supplementary Table 12 Global oceanic N₂O emissions based on multiple models. Unit: Tg N yr⁻¹

Model	1980s	1990s	2000s	2007-2016
Bern-3D	4.4	4.4	4.3	4.3
NEMOv3.6-PISCESv2-gas	3.3	3.2	3.3	3.4
NEMO-PlankTOM10	3.0	2.8	2.7	2.5
UVic2.9	3.3	3.2	3.2	3.1
NEMO-PISCES 3.2	4.0	3.9	3.9	3.8
Mean	3.6	3.5	3.5	3.4
Min	3.0	2.8	2.7	2.5
Max	4.4	4.4	4.3	4.3

Supplementary Table 13 Global N₂O emissions based on multiple top-down approaches. Unit: Tg N yr⁻¹

Name	Category	2000s	2007-2016
INVICAT	Land	9.7	10.6
	Ocean	7.2	7.1
	Total	16.9	17.7
PyVAR-1	Land	9.4	10.6
	Ocean	6.4	6.4
	Total	15.8	17.0
PyVAR-2	Land	11.8	12.7
	Ocean	4.0	4.3
	Total	15.8	17.0
MIROC4- ACTM	Land	12.5	13.8
	Ocean	3.1	3.4
	Total	15.7	17.1
GEOSChem	Land	10.6	11.3
	Ocean	4.5	4.6
	Total	15.1	15.9
Mean	Land	10.8	11.8
	Ocean	5.1	5.1
	Total	15.9	16.9
Min	Land	9.4	10.6
	Ocean	3.1	3.4
	Total	15.1	15.9
Max	Land	12.5	13.8
	Ocean	7.2	7.1
	Total	16.9	17.7

Supplementary Table 14 Comparison of terminologies used in this study and previous reports.

GCP Terminology (in this study)		IPCC AR5 (IPCC, 2013)	National GHG inventories (used by UNFCCC according to IPCC, 2006 and IPCC, 2019)	UNFCCC / IPCC 2006 Source sector
<i>Anthropogenic sources</i>				
Direct emissions of N additions in the agricultural sector (Agriculture)	Direct soil emissions (mineral N and manure fertilization, cultivation of organic soils, and crop residue returns)	Agriculture	Direct N ₂ O emissions from managed soils (except due to grazing animals)	3Da without 3Da3
	Manure left on pasture		Urine and dung deposited by grazing animals	3Da3
	Manure management		Manure management	3B
	Aquaculture	---	---	---
Other direct anthropogenic sources	Fossil fuel and industry	Fossil fuel combustion and industrial processes	Energy and industrial processes	1, 2
	Waste and waste water	Human excreta	Waste	5
	Biomass burning (from crop residue, grassland, shrubland and savannas; peat fires, tropical forests, boreal forests, and temperate forests)	Biomass and biofuel burning	Prescribed burning of savannas, field burning of agricultural residues	3E, 3F
Indirect emissions from anthropogenic N additions	Inland and coastal waters (rivers, lakes, reservoirs, estuaries, and coastal zones)	Rivers, estuaries, coastal zones	Indirect emissions due to leaching and runoff	3Db2
	Atmospheric N deposition on land	Atmospheric deposition on land	Indirect emissions due to atmospheric deposition (of agricultural as well as other anthropogenic compounds emitted)	part of 3Db1
	Atmospheric N deposition on ocean	Atmospheric deposition on ocean		part of 3Db1
Perturbed fluxes from climate/CO ₂ /land cover change	CO ₂ effect	---	---	---
	Climate effect	---	---	---
	Post-deforestation pulse effect	---	---	---
	Long-term effect of reduced mature forest area	---	---	---
<i>Natural sources and sinks</i>				
Natural soils baseline		Soils under natural vegetation	---	---
Ocean baseline		Oceans	---	---
Natural (rivers, lakes, reservoirs, estuaries, and coastal upwelling)		---	---	---
Lightning and atmospheric production	Lightning		---	---
	Atmospheric chemistry		---	---
Soil/wetland surface sink		Surface sink	---	---

Supplementary Table 15 Comparison of the global N₂O budget in this study with the IPCC AR5.

	This study (2007–2016)	IPCC AR5 (2006/2011)
Bottom-up budget		
Anthropogenic Sources		
Fossil fuel combustion and industry	1.0 (0.8–1.1)	0.7 (0.2–1.8)
Agriculture (incl. Aquaculture)	3.8 (2.5–5.8)	4.1 (1.7–4.8)
Biomass and biofuel burning	0.6 (0.5–0.8)	0.7 (0.2–1.0)
Wastewater	0.3 (0.2–0.5)	0.2 (0.1–0.3)
Rivers, estuaries, and coastal zones	0.5 (0.2–0.7)	0.6 (0.1–2.9)
Atmospheric N deposition on ocean	0.1 (0.1–0.2)	0.2 (0.1–0.4)
Atmospheric N deposition on land	0.8 (0.4–1.4)	0.4 (0.3–0.9)
Other indirect effects from CO ₂ , climate and land-use change	0.2 (-0.6–1.1)	
Total Anthropogenic	7.3 (4.2–11.4)	6.9 (2.7–12.1)
Natural Sources and Sinks		
Rivers, estuaries, and coastal zones	0.3 (0.3–0.4)	
Oceans	3.4 (2.5–4.3)	3.8 (1.8–9.4)
Soils under natural vegetation	5.6 (4.9–6.5)	6.6 (3.3–9.0)
Atmospheric chemistry	0.4 (0.2–1.2)	0.6 (0.3–1.2)
Surface sink	-0.01 (0– -0.3)	-0.01 (0– -1)
Total natural	9.7 (8.0–12.0)	11.0 (5.4–18.6)
Total bottom-up source	17.0 (12.2–23.5)	17.9 (8.1–30.7)
Observed growth rate	4.3 (3.8–4.8)	3.6 (3.5–3.8)
Tropospheric sink	0.1 (0.1–0.2)	
Stratospheric sink*	13.4 (12.3–14.4)	14.3 (4.3–28.7)
Atmospheric inversion		
Atmospheric loss	12.4 (11.7–13.3)	11.9 (11.0–12.8)
Total source	16.9 (15.9–17.7)	15.8 (14.8–16.8)

Note: * Calculated from satellite observations combined with simple photolysis models in our study.

Supplementary Table 16 Simulation experiments in the NMIP (Tian et al.^{1,17})

	CLIM	CO ₂	LCC	NDEP	NFER	MANN
SE0	1901-1920*	1860	1860	1860	1860	1860
SE1	1901-2016	1860-2016	1860-2016	1860-2016	1860-2016	1860-2016
SE2	1901-2016	1860-2016	1860-2016	1860-2016	1860-2016	1860
SE3	1901-2016	1860-2016	1860-2016	1860-2016	1860	1860
SE4	1901-2016	1860-2016	1860-2016	1860	1860	1860
SE5	1901-2016	1860-2016	1860	1860	1860	1860
SE6	1901-2016	1860	1860	1860	1860	1860

Note: CLIM: climate condition; CO₂: atmospheric CO₂ concentration; LCC: land cover change; NDEP: atmospheric N deposition; NFER: mineral N fertilizer use; and MANN: manure N use in cropland. SE0: baseline and control run with repeated climate forcing from 1901-1920; SE1: CLIM+CO₂+LCLU+NDEP+NFER+MANN; SE2: CLIM+CO₂+LCLU+NDEP+NFER; SE3: CLIM+ CO₂+LCLU+NDEP; SE4: CLIM+ CO₂+LCLU; SE5: CLIM+ CO₂; SE6: CLIM. “1901-1920*” denotes that variable is constant at the level of 20-year average; “1860” denotes that variable is constant at the level of 1860; and “1860-2016” denotes that variable changes with time over the study period.

Supplementary Table 17 Information of NMIP models using in this study

Model	Contact	Affiliation	Publication
DLEM	Hanqin Tian	Auburn University	Tian et al. ¹⁸ , Xu et al. ¹⁹
LPX-Bern	Sebastian Lienert/ Fortunat Joos	University of Bern, Switzerland	Stocker et al. ²⁰ , Xu-Ri & Prentice ²¹
O-CN	Sönke Zaehle	Max Planck Institute for Biogeochemistry	Zaehle et al. ²²
ORCHIDEE	Nicolas Vuichard	IPSL – LSCE, France	Goll et al. ²³
ORCHIDEE- CNP	Jinfeng Chang/ Daniel Goll	IPSL – LSCE, France	
VISIT	Akihiko Ito	National Institute for Environmental Studies, Japan	Inatomi et al. ²⁴ , Ito et al. ²⁵

Supplementary Table 18 Summary of models in ocean N₂O inter-comparison

Group	Model	Native resolution (Lon × Lat × Depth)	Publication
U. Bern	Bern-3D	9° × 4.5° × 32 levels	Battaglia and Joos ³
CNRM	NEMOv3.6- PISCESv2-gas	1° × 1° × 75 levels	Berthet et al. ⁴
UEA	NEMO-PlankTOM10	2° × (0.5°–2°) × 30 levels	Buitenhuis et al. ⁵
GEOMAR	UVic2.9	3.6° × 1.8° × 19 levels	Landolfi et al. ⁶
IPSL	NEMO-PISCES 3.2	2° × (0.5°–2°) × 30 levels	Martinez-Rey et al. ⁷

Supplementary Table 19 Overview of the inversion frameworks that are included in the global N₂O budget.

Name	ACTM	Method	Resolution of state vector	ACTM horizontal resolution	ACTM vertical levels	Ocean prior
INVICAT	TOMCAT	4D-Var	5.625°×5.625°	5.625°×5.625°	60	1 (high)
PyVAR-1	LMDz5	4D-Var	3.75°×1.875°	3.75°×1.875°	39	1 (high)
PyVAR-2	LMDz5					2 (low)
MIROC4-ACTM	MIROC4-ACTM	Bayesian analytical	84 regions	2.8°×2.8°	67	3 (low)
GEOSChem	GEOSChem	4D-Var	5°×4°	5°×4°	47	2 (low)

Supplementary Fig. 1 Spatial distribution of ten study regions across the globe.

Supplementary Fig. 2 Per capita N₂O emission (kg N capita⁻¹ yr⁻¹) during 2007–2016. Annual population was obtained from FAOSTAT¹¹⁵ (<http://www.fao.org/faostat/en/#data/OA>).