A vertical strip on the left side of the slide showing a microscopic view of marine diatoms. The diatoms are elongated, rod-like structures with intricate patterns of silica, appearing in various orientations and colors (yellow, green, blue) against a dark background. A scale bar in the bottom left corner indicates 50 µm.

Declining silica availability – a challenge in the North Atlantic region?

-The illustrated abstract-

Kriste Makareviciute-Fichtner^{1,2}, Birte Matthiessen¹,
Heike K. Lotze³, and Ulrich Sommer¹

¹GEOMAR Helmholtz Center for Ocean Research Kiel, Marine Ecology,
Germany (kmakareviciute@geomar.de)

²HOSST Research School, GEOMAR Helmholtz-Centre for Ocean Research Kiel,
Wischofstraße 1-3, 24148 Kiel, Germany

³Department of Biology, Dalhousie University, Halifax, Canada

Introduction

Understanding how changes in limiting nutrient availability affect life in the oceans requires interdisciplinary efforts. Here we illustrate this with an example of silicon, one of the most common elements on land which bioavailable form, silicic acid ($\text{Si}(\text{OH})_4$), is a limiting nutrient for silicifying primary producers, such as diatoms.

Introduction

- Silicic acid concentrations have declined in several parts of North Atlantic region (↓)
- Many coastal areas of the North Atlantic region also face decreased relative availability of silicon due to increased riverine supply of nitrogen and phosphorus and stable or declining loads of silicon.

- Both declining silicic acid concentrations and declining silicon to nitrogen (Si:N) ratios limit the growth of diatoms, which are major primary producers contributing up to a quarter of global primary production.

Methods

To assess the effects of declining silicon availability on phytoplankton communities we conducted a mesocosm experiment manipulating Si:N ratios and copepod grazing pressure on phytoplankton communities from the Baltic Sea.

Results

Declining Si:N ratio affected not only diatom abundance and relative biomass but also their species composition and overall plankton diversity.

This non-metric multidimensional scaling (NMDS) ordination shows that phytoplankton communities cluster based on grazing pressure and as a gradient of Si:N ratio treatments

Lowered Si:N ratio $\Rightarrow$ lower phytoplankton biomass, different species composition, lower diversity (but this can be upgraded by grazers!)

Our results illustrate the importance of silicon in structuring community composition at the base of temperate marine food webs. Changes in silicic acid concentrations and Si:N ratios, therefore, may have far-reaching consequences on oceanic primary production and planktonic food webs.

Discussion

The decline in silicon concentrations in polar and subpolar North Atlantic waters is attributed to natural multi-decadal variability but is likely amplified by reduced ocean mixing due to increased water temperatures, illustrating the need of international efforts to curb global climate change.

The decline in Si:N ratios in coastal oceans also highlights the need for further reduction of nutrient pollution and improved river basin management.

This may require interdisciplinary and international approaches to manage anthropogenic perturbations of the silicon cycle.

Thank you for having a look!

If you have questions, feel free to ask during our
session chat or contact me at
kmakareviciute@geomar.de