

BERICHTE

aus dem Fachbereich Geowissenschaften
der Universität Bremen

No. 270

Ruhland, G., Fischer, G. and cruise participants

REPORT AND PRELIMINARY RESULTS OF POSEIDON CRUISE 365 (LEG 1+2)

**Leg 1: Las Palmas – Las Palmas
13.04.2008 – 16.04.2008**

**Leg 2: Las Palmas – Las Palmas
18.04.2008 – 29.04.2008**



Berichte, Fachbereich Geowissenschaften, Universität Bremen, No. 270,
40 Seiten, Bremen 2009

ISSN 0931-0800

The "Berichte aus dem Fachbereich Geowissenschaften" are produced at irregular intervals by the Department of Geosciences, Bremen University.

They serve for the publication of experimental works, Ph.D.-theses and scientific contributions made by members of the department.

Reports can be ordered from:

Monika Bachur

Forschungszentrum Ozeanränder, RCOM

Universität Bremen

Postfach 330 440

D 28334 BREMEN

Phone: (49) 421 218-65516

Fax: (49) 421 218-65515

e-mail: MBachur@uni-bremen.de

<http://elib3.suub.uni-bremen.de/publications/diss/html>

Citation:

Ruhland, G. , G. Fischer and cruise participants

Report and preliminary results of R/V POSEIDON Cruise 365 (Leg 1+2). Leg 1: Las Palmas - Las Palmas,
13.4. - 16.4.2008. Leg 2: Las Palmas - Las Palmas, 18.4. - 29.4.2008.

Berichte, Fachbereich Geowissenschaften, Universität Bremen, No. 270, 40 pages. Bremen, 2009.

ISSN 0931-0800

Contents: Leg 1 and 2

1.	Narrative	2
2.	Preliminary results.....	5
2.1.	Atmospheric Sciences	5
2.1.1.	Aerosol sampling	5
2.2.	Biogeochemistry and Oceanography (Leg 1).....	6
2.2.1.	Surveys at ESTOC and off Cape Blanc	6
2.2.2.	Testing the new optical nitrate sensor (NICO project)	19
2.3.	Marine Microbiology	24
2.3.1	Particle composition and size-related settling velocity.....	24
2.4.	Oceanography and Marine Geology	26
2.4.1	CTD-O ₂ -chlorophyll-fluorescence probes	26
2.4.2.	Sampling for chlorophyll-a measurements	27
2.4.3.	Particle fluxes measured with sediment traps	28
2.4.4.	Particle distribution measured with optical systems (ParCa).....	30
2.4.5.	Particle investigations with the ROV <i>Cherokee</i>	32
2.5.	Marine Techology	35
2.5.1.	Recovery of the DOLAN buoy	35
2.5.2.	Recovery of the PACT mooring	36
2.6.	References	37
2.7.	Acknowledgements	38
3.	Station List (Leg 1 and 2)	39

1. Narrative

RV Poseidon left the port of Las Palmas in the morning of April 13th, 2008 towards the DOLAN buoy position in the north of the island of Tenerife. In the afternoon the PACT mooring close to the buoy was successfully recovered. Afterwards a CTD cast for post-calibration had been done. The next morning (April 14th) the recovery of the whole buoy mooring was started. This work could be finished at noon. Due to the fact that the project time ended, the buoy was not redeployed. The ship steamed towards the ESTOC monthly monitoring position. On the way, an XBT was launched. In the late afternoon, the two CTD casts for the monthly monitoring were done. Overnight the ship steamed to the ESTOC sediment trap mooring. On April 15th this mooring was released and successfully recovered. All three particle traps provided full sets of sample bottles. At noon, a CTD cast to 800 m water depth was launched. Afterwards a NOAA drifter was deployed. At 14:00h ESTOC sediment trap mooring was redeployed after maintenance. Overnight, RV Poseidon steamed back to Las Palmas and arrived there in the morning of April 16th.

RV Poseidon left the port of Las Palmas (Gran Canaria) on April, 18, 2008 at 8:30 pm for Leg 2 sailing in southwesterly direction to the study area off Cape Blanc (Mauritania). We planned to perform biogeochemical, marine-chemical, microbial and micropaleontological studies off Cape Blanc as well as the exchange of two sediment trap moorings which were deployed during RV MERIAN cruise the year before. Additionally, a particle camera was planned to be launched to measure the distribution and size of marine snow aggregates. The ROV *Cherokee* was scheduled to be deployed for the observation of larger particles in the water column.

On Sunday, 20th of April, we reached the first study site about 200 nm off Cape Blanc at about 21°17.N and 20°49.W. Here, we recovered the sediment trap mooring CB-18 within three hours; all instruments had worked. The study site is located at the edge of the coastal upwelling filament off Cape Blanc. After launching the rosette with the shipboard SBE-3-CTD (additionally equipped with oxygen and chlorophyll sensors) to 800 m water depth, the particle camera ParCa with a SBE-19 CTD (with oxygen and chlorophyll fluorescence sensors) was launched to 2000 m water depth. On early monday morning, April 21, a deep rosette-CTD station to 2000 m was performed. The new sediment trap mooring CB-19 was then redeployed on the former position of CB-18 within two hours.

We then sailed to the second station having increasingly bad weather with strong winds and a high swell from the north. Therefore, we only launched an XBT to 1800 m water depth. We sailed further eastwards crossing station 3 (mooring site CBi-5) and finally decided to launch a shallow rosette-CTD down to 700 m at station 4 (about 20°44.N, 18°W). During the night, we moved back to the mooring site CBi-5/6 (station 3). After deploying a ParCa-CTD to 2500 m water depth, we

decided to release the mooring CBi-5 on Wednesday morning, April 23, during relatively rough weather conditions. The upper MSD trap with 40 sample containers had worked perfectly, the lower trap had failed. Lateron, we sailed to the south to station 5, launching an XBT to 1800 m, a shallow rosette-CTD to 800 m as well as a ParCa-CTD to 2500 m. During the night we moved northwards to the mooring site CBi-5/6. In the early morning of Thursday, April 24, we deployed a shallow and a deep rosette-CTD. As the weather improved, we could deploy the new mooring CBi-6 in the afternoon, now equipped with an additional MSD platform with an FSI-CTD and a video camera system. This system should record the particle distribution and transport over an annual cycle. Additionally, a high resolution MSD sediment trap with 40 cups was deployed in the upper part water column, the lower water column will be sampled by a classical 20-cup sediment trap. In the afternoon, we sailed to station 4 to launch the ParCa-CTD to 750 m water depth. Afterwards, we sailed northwards to station 7. As the wind and the swell further decreased, we were able to deploy the ROV *Cherokee* for about 5 hours. We studied four water levels at about 50, 100, 250 and 400 m water depths for the estimation of *in situ* particle settling rates using the settling chamber which is mounted to the ROV. The new settling chamber worked perfectly. The station work was completed with a shallow rosette-CTD and the ParCa-CTD down to 2000 m. Late Friday night, April 25th, we sailed northwards to Las Palmas, Gran Canaria. We reached Las Palmas earlier as planned in the early Monday morning of April 28th, 2008.




Fig. 1. Cruise track of POS 365 (Leg 1 and 2). The location of the DOLAN buoy and the sediment trap moorings CI/ESTOC, CBI and CB are also shown.

2. Preliminary Results

2.1. Atmospheric Sciences

2.1.1. Aerosol sampling (*Inka Meyer*)

Terrigenous sediments deposited in marine sediments are a mixture of a pelagic component brought in by the wind and a hemipelagic component supplied by rivers and from the shelf. The analysis of eolian dust allows the estimation of aridity in eolian source regions and the intensity of the transporting winds through grain-size measurements. Eolian dust can be used to reconstruct changes in continental climate. For validating the terrigenous sediment fraction in marine sediment cores, present-day dust samples were collected during the cruise.

The collecting was done with two dust-collectors which were placed on the observation deck. An engine inside both dust-collectors sucks the surrounding air. This air passes through a filter in which the dust is collected. The dust-collectors are connected with a vane to prevent that particles from the funnel reach the filters. If the dust collectors have no connection with the sensor, the engines stops automatically. Two kinds of filters are used for several investigations. The glassfibre-filter is for analysing the organics in the dust, while the cellulose-filters will be used for grain-size measurements, chemical and mineralogical analysis. Seven samples were taken on the second leg (Table 1).

Table 1. Samples collected for dust investigations.

Collection time 2008	Sample identification
19-20/04	GlassFib01, CellFib01
21-22/04	GlassFib02, CellFib02
22-23/04	GlassFib03, CellFib03
23-24/04	GlassFib04, CellFib04
25-26/04	GlassFib05, CellFib05
26-27/04	GlassFib06, CellFib06
27-28/04	GlassFib07, CellFib07

2.2. Biogeochemistry and Oceanography (Leg 1)

2.2.1. Surveys at ESTOC and off Cape Blanc (*M. Villagarcía, D.V. Moreno*)

Objectives and scientific questions

Along P365a the ANIMATE/ ESTOC mooring was recovered for the last time. It had been put in place in November 2007 during P360 and the sensors had been in the water for about 5 months.

At the same time, it was necessary to do the biogeochemical monthly samplings at the ESTOC station (European Station for Time series in the Ocean Canary islands), as it has been continuously done since 1994. A calibration cast before the final recovery of the mooring with CTD/Rosette were also made to accomplish the requirements of the sensors being recovered.




Fig. 2. Position of the CTD stations (blue circles) made along Poseidon P365 (Leg 1).

At the beginning of the cruise, the DOLAN/ANIMATE mooring had to be recovered, hence a rosette/CTD cast to 1000m was made in order to have a calibration of the chemical sensors before their recovery (station #212_2, 13th April 2008). The same cast was used to collect water for the sediment trap mooring that was going to be replaced.

Then, after the DOLAN mooring with the chemical sensors from ANIMATE was recovered, the ESTOC station monthly sampling took place by means of two profiles done on 14th and 15th of April 2008 (station 215_01, sampled to the bottom and 217_01 sampled to 800m) (see station list).

The nutrient sensor NAS-3X[®] number 2625 was recovered with data of 5 months, the preliminary analysis shows in general good data with some gaps where the data was unacceptable. One of the

standards did not provide adequate response, only the standard with a concentration 2 of nitrate was used to calculate the nitrate and nitrite data.

Table 2. List of stations sampled along the cruise P365 (Leg 1), Las Palmas- DOLAN- Las Palmas (O=oxygen, CO₂ = Carbon dioxide, N=nutrients, S=salinity, C=chlorophyll “a”, HPLC = pigments, Inc= Incidences) .

Date Time (hh:mm)	St. #, CTD, Cast	Depth sta,m	Lat. N	Long. W	Depth Ro,db	Ni sk b ot	Depth samp, db	PARAMETERS							
								O	CO ₂	N	S	C	HPLC	Inc	
13.04. 2008 18:54	212, 02	3634	29°11.53'	15°56.05'	1000			Station to collect water fro traps and for pre-recovery calibration of mooring sensors							
						1	1000	Sediment traps water							
						2	1000	Sediment traps water							
						3	200	✓		✓	✓	✓	✓	✓	
						4	150	✓		✓		✓	✓	✓	
						5	100	✓		✓		✓	✓	✓	
						6	90	✓		✓		✓	✓	✓	
						7	80	✓		✓		✓	✓	✓	
						8	70	✓		✓		✓	✓	✓	
						9	55	✓		✓		✓	✓	✓	
						10	40	✓		✓		✓	✓	✓	
						11	25	✓		✓	✓	✓	✓	✓	
						12	10	✓		✓		✓	✓	✓	
14.04. 2008 16:15	215, 01	3643	29°10.00'	15°20.00'	3500			ESTOC April 2008, Deep profile							
						1	3500	✓	✓	✓	✓	✓			
						2	3000	✓	✓	✓	✓	✓			
						3	2800	✓	✓	✓	✓	✓			
						4	2500	✓	✓	✓	✓	✓			
						5	2000	✓	✓	✓	✓	✓			
						6	1800	✓	✓	✓	✓	✓			
						7	1500	✓	✓	✓	✓	✓			
						8	1300	✓	✓	✓	✓	✓			
						9	1200	✓	✓	✓	✓	✓			
						10	1100	✓	✓	✓	✓	✓			
						11	1000	✓	✓	✓	✓	✓			
						12	800	✓	✓	✓	✓	✓			
15.04. 2008 11:40	217, 01	3595	29°10.02'	15°20.05'	800			ESTOC April 2008, Shallow profile							
						1	800	✓	✓	✓	✓	✓			
						2	600	✓	✓	✓	✓	✓			
						3	400	✓	✓	✓	✓	✓			
						4	300	✓	✓	✓	✓	✓			
						5	200	✓	✓	✓	✓	✓	✓	✓	
						6	150	✓	✓	✓	✓	✓	✓	✓	
						7	125	✓	✓	✓	✓	✓	✓	✓	
						8	100	✓	✓	✓	✓	✓	✓	✓	
						9	75	✓	✓	✓	✓	✓	✓	✓	
						10	50	✓	✓	✓	✓	✓	✓	✓	
						11	25	✓	✓	✓	✓	✓	✓	✓	
						12	10	✓	✓	✓	✓	✓	✓	✓	

Physical (CTD, salinity) and biochemical (oxygen, CO₂, nutrients, chlorophyll and pigments) parameters were measured in order to characterize the water masses present in the water column (Table 2). Some of the parameters (oxygen, chlorophyll and pigments filtration) were analysed on board after sampling, and others were taken frozen to the ICCM (nutrients and filters from chlorophyll).

Methods

Water Sampling

Samples were collected immediately after the Niskin bottles were on board from each depth. The sampling sequence was as follows:

- 1.) Oxygen: was taken in glass bottles of about 125 ml of volume which were previously cleaned and washed with HCl acid and was fixed at once; then it was kept for at least six hours according to WOCE regulations and finally it was analysed at the laboratory on board the ship.
- 2.) Carbon system measurements: only taken at ESTOC, and in this case pH and alkalinity: samples were taken in glass bottles and were fixed immediately on board.
- 3.) Nutrients: were taken in polypropylene bottles which were previously cleaned and washed with HCl acid and were completely dry. Samples were immediately frozen at -20°C, analysing them as soon as possible after arrival at the laboratory. Freezing the samples is a common practice; it does not or only in a non-significant way affect the nitrate and nitrite and the phosphate values (by a slight decrease) and is not noticeable in the silicate values (Kremling and Wenck, 1986; McDonald and McLunghlin, 1982).
- 4.) Salinity: samples were taken in dark glass bottles which were previously cleaned and washed with HCl acid. Then, they were kept in boxes to protect them from light till analysis on land.
- 5.) Chlorophyll: samples of one liter of water were taken. The chlorophyll samples were filtered immediately and the filters were frozen subsequently at -20°C. Their analyses takes place at the ICCM home laboratory.
- 6.) Pigments: samples of one liter of water were taken and filtered immediately after sampling took place and the filters were deep frozen in liquid nitrogen to be maintained at -80°C. Their analyses takes place at the ICCM home laboratory.

All samples were taken using the procedures established in the WOCE Operations Manual, WHP Office Report WHPO 91-1/WOCE Report No.68/91.

Analysis

Dissolved Oxygen :The samples were analysed using the method described in the WOCE Operations Manual, WHP Office Report No. 68/91; the final titration point was detected using a Metrohm 665 Dosimat Oxygen Auto-Titrator Analyser.

Carbonate system measurements

The pH_t in total scale ($\text{mol} (\text{kg-SW})^{-1}$) was measured following the spectrophotometric technique of Clayton and Byrne (1993) using the m-cresol purple indicator (DOE, 1994). 0.0047 pH units were added to the pH experimental values in order to take into consideration the recommendations by Lee et al. (2000). A system similar to that described by Bellerby et al. (1995) was developed in our lab. The pH_t measurements were carried out using a Hewlett Packard Diode Array spectrophotometer in a 25°C-thermostated 1-cm flow-cell using a Peltier system. A stopped-flow protocol was used to analyse seawater previously thermostated to 25°C for a blank determination at 730, 578 and 434 nm. The flow was restarted, and the indicator injection valve switched on to inject 10 µl dye through a mixing coil (2 m). Three photometric measurements were carried out for each injection in order to remove all dye effect on the seawater pH_t measurement. Repeatedly, seawater measurements of the different Certified Reference Materials (CRM provided by Dr. Dickson, Scripps Institution of Oceanography) samples gave a standard deviation of ± 0.0015 ($n = 54$).

The total alkalinity of seawater (A_T) was determined by titration with HCl to the carbonic acid end point using two similar potentiometric systems, as described in more detail by Mintrop et al. (2000). In order to yield an ionic strength similar to open ocean seawater, the HCl solution (25 l, 0.25 M) was made from concentrated analytical grade HCl (Merck®, Darmstadt, Germany) in 0.45 M NaCl. The acid was standardised by titrating weighed amounts of Na_2CO_3 dissolved in 0.7 M NaCl solutions. The total alkalinity of seawater was evaluated from the proton balance at the alkalinity equivalence point, $\text{pH}_{\text{equiv}} = 4.5$, according to the exact definition of total alkalinity (Dickson, 1981). The performance of the titration systems was monitored by titrating different samples of certified reference material (CRM, batch 42) with known inorganic carbon and A_T values. The agreement between our data and CRM values was within $\pm 1.5 \mu\text{mol kg}^{-1}$. Total inorganic carbon (C_T) is computed from experimental values of pH_t and total alkalinity, using the carbonic acid dissociation constants of Mehrbach after Dickson and Millero (1987). This set of constants presented the best agreement between $C_T(\text{pH}, A_T)$ calculations and certified C_T values for CRM, batch 42, with a C_T residual of $\pm 3 \mu\text{mol kg}^{-1}$, $n=54$ (Millero, 1995, Lee et al., 1997).

Nutrients

The nutrients determination was performed with a segmented continuous-flow autoanalyser, a Skalar® San Plus System (ICCM).

Nitrate and Nitrite: The automated procedure for the determination of nitrate and nitrite is based on the cadmium reduction method; the sample is passed through a column containing granulated copper-cadmium to reduce the nitrate to nitrite (Wood et al., 1967), using ammonium chloride as pH controller and complexer of the cadmium cations formed (Strickland and Parsons, 1972). The optimal column preparation conditions are described by several authors (Nydahl, 1976; Garside, 1993).

Phosphate: Orthophosphate concentration is understood as the concentration of reactive phosphate (Riley and Skirrow, 1975) and according to Koroleff (1983a) is a synonym of “dissolved inorganic phosphate”. The automated procedure for the determination of phosphate is based on the following reaction: ammonium molybdate and potassium antimony tartrate react in an acidic medium with diluted solution of phosphate to form an antimony-phospho-molybdate complex. This complex is reduced to an intensely blue-coloured complex, ascorbic acid. The complex is measured at 880nm. The basic methodology for this anion determination is given by Murphy and Riley (1962); the used methodology is the one adapted by Strickland and Parsons (1972).

Silicate: The determination of the soluble silicon compounds in natural waters is based on the formation of the yellow coloured silicomolybdic acid; the sample is acidified and mixed with an ammonium molybdate solution forming molybdsilicic acid. This acid is reduced with ascorbic acid to a blue dye, which is measured at 810nm. Oxalic acid is added to avoid phosphate interference. The used method is described in Koroleff (1983b).

Phytoplankton pigments: Pigments were measured using fluorimetric analysis, following the methodology described by Welschmeyer (1994). The determination was achieved using a fluorometer TURNER 10-AU-000.

Salinity: Samples were measured with a salinometer, model Autosal 8400a, whose measurement range was between 0.005-42 (psu), with an accuracy of ± 0.003 , according to the manufacturer. It was calibrated following the manufacturer's information and standardizing it with IAPSO Standard Seawater. Salinity values were calculated as practical salinity according to Unesco (1978, 1984).

Pigments: Samples are defrozen in an ultrasound bath and extraction is done with methanol before being analysed by HPLC (High Performance Liquid Chromatography). The extraction process used

was described by Wright et al.(1991); this procedure was proposed as the JGOFS (Joint Global Ocean Flux Study) standard methodology for phytoplankton pigment determination (1994). Some modifications for improving the method were done by Wright in 2006.

Preliminary Results

This cruise took place during spring, which corresponds in surface waters and for this area, to the time of the year when the mixed layer of about 125-150 m that was formed during the late winter disappears, and it starts the seasonal thermocline formation which is shown by the slope profile found (Figure 3). If we look at the results of the preceding year, the 2007 sampling in the spring took place along P320 in March, and hence temperature values were lower and the mixed layer was more homogeneous (Figure 4).




Fig. 3. T/S diagram for the stations sampled (left) and potential temperature versus pressure showing the breakage of the mixed layer (right).

The only station sampled for deeper waters was ESTOC and there the subsurface waters encountered below the mixed layer represent the temperature and salinity values that characterise the NACW (North Atlantic Central Water) straight line. The range of values were 12- 17.5°C and 35.6- 36.6 for temperature and salinity, respectively. (Figure 3). At intermediate waters level, no presence of Mediterranean Water (MW) was found at ESTOC during the cruise (about 7-11°C of temperature), since all salinity values encountered were below 35.5 units, which is the limit of the Mediterranean Water (see example in Llinás et al., 2003). However, at these depths, the Antarctic Intermediate Water (AAIW) was found and from Figure 3 it seems there was a mixing of both types of water masses, giving a salinity maximum of about 35.5 in that water depth.




Fig. 4. Mixed layer of stations including ESTOC (in red) from cruise P320 sampling at ESTOC in March 2007 (left). Nitrate and nitrite values from the stations made along P365 (Leg 1) (right).

Nutrients for the upper 200m are shown in Figures 4 and 5. The nitracline is detected at 100m just below the chlorophyll “a” maximum (Figure 6), above these depths there is depletion of nitrates and nitrites and phosphates due to the photosynthetic activity of the surface waters.




Fig. 5. Phosphate values taken at both stations made along P365 (Leg 1) for the upper 200m (left) and silicate values from the same stations (right).

The silicate values are slightly higher than the other two nutrients, a feature characteristic for this area. Below 100m there is a strong increase of nutrients values, detecting around 6.5 $\mu\text{mol/l}$ of nitrate and nitrite, 0.35 $\mu\text{mol/l}$ of phosphates and 2 $\mu\text{mol/l}$ of silicates at 200m. These nutrients values have been compared with the 10-year nutrients time series that ICCM has been taking from ESTOC, and the data measured are within the expected range for the spring in this area.




Fig. 6. Chlorophyll “a” sampled at both stations made along P365 (Leg 1) for the upper 200m (left) and oxygen values for the same stations (right).

The chlorophyll „a“ is plotted in Figure 6 for ESTOC and the calibration station. Values ranged between 0 and 0.45 µg/l at latitude 29°N, showing a maxima at 70m of depth for ESTOC and about 90 m for station 212_2. The variability found for the maxima at each station is related to thermocline formation effects already mentioned. The mixed layer was loosing its homogeneity along the cruise days, and if there were inputs of nutrients from below, the amount of phytoplankton increases producing greater fluorescence values at different depths. An increase in phytoplankton standing stock is also shown in the increase of the oxygen maximum at about 50 m in one of the stations (Figure 6).

The cruise P365 Leg 2 had the following aims for ICCM:

- Seven expandable bathythermograph (XBT, T5, 1830 m) were launched to characterize the water column between the south of the islands and the most westerly station of Cape Blanc (Table 3).
- To carry out hydrographic investigations in the Cape Blanc area, 6 stations were made at this latitude, some to 2000 m and other down to 800 m. Samples for oxygen, nutrients, salinity and chlorophyll “a” were taken. Oxygen values were determined on board (Table 4).
- In Cape Blanc stations were the CTD was only done until 800 m, a XBT was launched (in stations: GeoB 12908(St2a), GeoB 12911(St4) and GeoB 12914 (St5) (Table 3).

Table 3. List of XBT's launched.

File name	Date (dd/mm/yy)	Latitude N	Longitude W
P365B_XBT_1	18/04/08	27° 00.000	16° 03.110
P365B_XBT_2	19/04/08	26° 02.574	16° 51.389
P365B_XBT_3	19/04/08	25° 05.148	17° 39.271
P365B_XBT_4	19/04/08	24° 07.722	18° 26.776
P365B_XBT_5	19/04/08	23° 10.296	19° 13.923
P365B_XBT_6	20/04/08	22° 12.870	20° 00.730
P365B_XBT_7	20/04/08	21° 16.797	20° 47.480
P365B_XBT_St2a	21/04/08	20° 59.485	19° 40.853
P365B_XBT_St4	23/04/08	20° 04.848	18° 58.960
P365B_XBT_St5	25/04/08	21° 18.462	18° 49.649




Fig. 7. Location of stations done by the ICCM and their positions off Cape Blanc, Mauritania.

Table 4. List of stations sampled by ICCM along 365 (Leg 2). DP= Deep Profile, SP= Shallow Profile, Chloroph= chlorophyll “a”, Inc= Incidences/ Comments. NW= not enough water available for some parameter.

Station Name: GeoB12907-2 (St1) SP:E227		Date: 20/04/08		Time: 20:03			
Niskin	Depth	Oxygen	Nutrients	Longitude: 20° 48.136 W	Salinity	Chloroph.	Inc.
1	800	✓	✓				
2	600	✓	✓		✓		
3	400	✓	✓				
4	300	✓	✓				
5	200	✓	✓			✓	
6	150	✓	✓			✓	
7	125	✓	✓			✓	
8	100	✓	✓			✓	
9	75	✓	✓			✓	
10	50	✓	✓			✓	
11	25	✓	✓			✓	
12	10	✓	✓	✓			✓

Table 4. continued

Station Name: GeoB12907-4 (St1)		Date: 21/04/08		Time: 06:11					
DP: E228	Latitude: 21° 16.908 N	Niskin	Depth	Oxygen	Nutrients	Longitude: 20° 48.136 W	Salinity	Chloroph.	Inc.
		1	2000						
		2	2000	√	√		√		
		3	1800	√	√				
		4	1600	√	√				
		5	1400	√	√				
		6	1200	√	√				
		7	1000	√	√				
		8	800	√	√		√		
		9	500						
		10	500	√	√				
		11	35						
		12	35	√	√			√	
Station Name: GeoB12909-1(St 2)		Date: 22/04/08		Time: 17:46					
SP: E231	Latitude: 21° 16.908 N	Niskin	Depth	Oxygen	Nutrients	Longitude: 20° 48.136 W	Salinity	Chloroph.	Inc.
		1	700						
		2	700	√	√		√		
		3	600	√	√				
		4	400						
		5	400	√	√				
		6	300	√	√				
		7	200	√	√			√	
		8	100	√	√			√	
		9	50	√	√			√	
		10	35						
		11	35	√	√		√	√	
		12	10	√	√			√	
Station Name: GeoB12911-2 (St4)		Date: 23/04/08		Time: 19:00					
SP: E234	Latitude: 20° 04.694 N	Niskin	Depth	Oxygen	Nutrients	Longitude: 18° 58.725 W	Salinity	Chloroph.	Inc.
		1	800	√	√				
		2	600	√	√		√		
		3	400	√	√				
		4	300	√	√				
		5	200	√	√			√	
		6	150	√	√			√	
		7	125	√	√			√	
		8	100	√	√			√	
		9	75	√	√			√	
		10	50	√	√			√	
		11	25	√	√		√	√	
		12	10	√	√			√	
Station Name: GeoB12912-1 (St3)		Date: 24/04/08		Time: 8:00					
SP: 235	Latitude: 20° 44.999 N	Niskin	Depth	Oxygen	Nutrients	Longitude: 18° 41.964 W	Salinity	Chloroph.	Inc.
		1	800	√	√				
		2	600	√	√		√		
		3	400	√	√				
		4	300	√	√				
		5	200	√	√			√	
		6	150	√	√			√	
		7	125	√	√			√	
		8	100	√	√			√	
		9	75	√	√			√	
		10	50	√	√			√	
		11	25	√	√		√	√	
		12	10	√	√			√	

Table 4. continued

Station Name: GeoB12912-2 (St3)		Date: 24/04/08		Time: 9:30					
DP: 235	Latitude: 20° 44.973 N	Niskin	Depth	Oxygen	Nutrients	Longitude: 18° 41.991 W	Salinity	Chloroph.	Inc.
		1	2500						
		2	2500	√	√		√		
		3	2200	√	√				
		4	2000	√	√				
		5	1800						
		6	1800	√	√				
		7	1500	√	√				
		8	1300	√	√				
		9	1000	√	√				
		10	800	√	√		√		
		11	75						
		12	75						

Station Name: GeoB12914-3 (St5)		Date: 25/04/08		Time: 18:46					
SP: 238	Latitude: 21° 19.83 N	Niskin	Depth	Oxygen	Nutrients	Longitude: 18° 49.58 W	Salinity	Chloroph.	Inc.
		1	800	√	√				
		2	600	√	√		√		
		3	400	√	√				
		4	300	√	√				
		5	200	√	√			√	
		6	150	√	√			√	
		7	100	√	√			√	
		8	75	√	√			√	
		9	50	√	√			√	
		10	25	√	√		√	√	
		11	10	√	√			√	
		12	Neptuno Gift						

Water Sampling

Samples were collected immediately after Niskin bottles were on board from each depth. The sampling sequence was as follows:

- Oxygen: was taken in glass bottles of about 125 ml of volume which were previously cleaned and washed with HCl acid and with the seawater which is going to be taken. Oxygen was fixed at once; then it was kept for at least six hours before the determination according to WOCE regulations and finally it was analysed at the laboratory on board the ship.
- Nutrients: were taken in polypropylene bottles which were previously cleaned and washed with HCl acid and were completely dry. In the moment of taking samples, bottles were cleaned with the seawater which is going to be taken. Samples were immediately frozen at least at -20°C, analysing them as soon as possible after arrival at the laboratory on land at the ICCM.
- Salinity: samples were taken in dark glass bottles which were previously cleaned and washed with HCl acid and before taking samples with that seawater. Then, they were kept in boxes to protect them from light till analysis on land at the ICCM.

- **Chlorophylla:** samples of one litre of water were taken. The chlorophyll samples were filtered immediately (500 ml, the rest of water is for cleaning the system) and the filters were frozen subsequently at least at -20°C. Their analysis will be done at the ICCM.

Analysis

- **Dissolved Oxygen:** samples were analysed using the method described in the WOCE Operations Manual, WHP Office Report No. 68/91; the final titration point was detected using a Metrohm 665 Dosimat Oxygen Auto-Titrator Analyser.
- **Nutrients:** nutrient determination will be performed with a segmented continuous-flow autoanalyser, a Skalar® San Plus System.
- **Phytoplankton Pigments:** pigments will be measured using fluorometric analysis, following the methodology described by Welschmeyer (1994). The determination will be achieved using a fluorometer TURNER 10-AU-000.
- **Salinity:** samples will be measured with a salinometer, model Autosal 8400a, whose measurement range was between 0.005-42 (psu), with an accuracy of ± 0.003 , according to the manufacturer.

Preliminary Results

On board only the XBT's results and oxygen measurements were obtained and the results are only preliminary (Fig. 8). Initial visual correction for errors was made; further checking will be done once the results of the parameters analysed on land are obtained.

Oxygen Results




Fig. 8. Oxygen profiles done on POS 365 (Leg 2): profiles down to 2500 m (left) and down to 800 m water depth (right).

XBT's Results

Values for the seven XBT launched during the transect Las Palmas-Cape Blanc are showed in Figure 9. Deep values show the general temperature values found for this subtropical area (4°-8°C). Antarctic Intermediate Water is sometimes detected around 600-800 m. Cooler subsurface waters are found to the south and east, probably due to the influence of the upwelled water encountered in the Cape Blanc area.




Fig. 9. Temperature versus depth for the seven XBT's of the transect (left) and for the three XBT stations off Cape Blanc (right).

2.2.2. Testing the new optical nitrate sensor (NICO project) (E. Kopiske)

This cruise covers the ROV test of the PROPS sensor with the ROV *Cherokee*. The “NICO” Project (Nitrate Continuous Observation Sensor) shall continue the development of the PROPS sensor in order to have an optical nitrate sensor for long term use in moorings in the marine science. The part of the MARUM is the practical use of the sensor on seagoing expeditions in several applications. The sensor was used in the laboratory with a flow-through-cuvette for the analysis of the samples from the rosette. The sensor was also mounted on the MARUM ROV *Cherokee* in order to get online photometric spectra during a ROV dive. The online-spectra can not only be used for nitrate

measurement but also for a quantitative measurement of other substances like yellow substance (Gelbstoff).

Offline measurements in the lab

92 water samples were collected during the cruise at 8 CTD stations (Table 5). Two bottles were taken from every sample, one was measured on board and the other was frozen at -40°C for a later reference analysis at the IFM-GEOMAR in Kiel. The bottles were rinsed with distilled water before the cruise and three times with the sample water before filling them with the sample. The second bottle was analyzed in the laboratory with the PROPS sensor by the use of a flow-through-cuvette. The cuvette was rinsed with approx. 50 ml of the sample before the analysis. Three measurements were taken with renewed water in the cuvette. The temperature was measured with a “Testo 826-T4” infrared temperature probe in order to have an uncontaminated sample.

In situ measurements with the ROV Cherokee

The PROPS sensor has never been used in an online and profiling configuration until now. The big advantage is that the spectra can be viewed online and that a substance analysis for nitrate or other substances can be viewed in parallel. The sensor was mounted on the ROV on the port forward corner of the ROV. The optical path length was 20 mm for this test. The connection was possible with a adapter bottle which converts the sensor interface to Schilling Seanet Hub standard. The connection to the ROV was made with a standard Schilling Seanet cable. The ROV provided the 24V power supply and an RS232 interface for the PROPS sensor. A Laptop with an USB to RS232 adapter was used for the GUI. The operator was sitting beside the ROV pilot in the ROV control lab. The standard msda_xe software was used to operate the sensor and to view the online data. The configuration of the sensor was to take measurements automatically every minute. The recorded spectra have been displayed as an overlay of 99 spectra. This allows the operator to analyze the change in the signal while descending. The recorded spectra show the increase of nitrate with depth and also a signal which was interpreted as yellow substance (Gelbstoff). This was the first time we had an online sensor for chemical parameters on the ROV. Only one ROV station was done due to high swell.

Table 5. List of samples taken for the NICO project.

Internal label	Ship_station #	GEOB #	Water depth
#1-215_01-3500	215-1	12903-1	3500
#2-215_01-3000	215-1	12903-1	3000
#3-215_01-2800	215-1	12903-1	2800
#4-215_01-2500	215-1	12903-1	2500
#5-215_01-2000	215-1	12903-1	2000

Table 5 continued

#6-215_01-1800	215-1	12903-1	1800
#7-215_01-1500	215-1	12903-1	1500
#8-215_01-1300	215-1	12903-1	1300
#9-215_01-1200	215-1	12903-1	1200
#10-215_01-1100	215-1	12903-1	1100
#11-215_01-1000	215-1	12903-1	1000
#12-215_01-800	215-1	12903-1	800
#1-217_01-800	217-1	12905-1	800
#2-217_01-600	217-1	12905-1	600
#3-217_01-400	217-1	12905-1	400
#4-217_01-300	217-1	12905-1	300
#5-217_01-200	217-1	12905-1	200
#6-217_01-150	217-1	12905-1	150
#7-217_01-125	217-1	12905-1	125
#8-217_01-100	217-1	12905-1	100
#9-217_01-75	217-1	12905-1	75
#10-217_01-50	217-1	12905-1	50
#11-217_01-25	217-1	12905-1	25
#12-217_01-10	217-1	12905-1	10
#1-3	227-1	12907-2	800
#2-3	227-1	12907-2	600
#3-3	227-1	12907-2	400
#4-3	227-1	12907-2	300
#5-3	227-1	12907-2	200
#6-3	227-1	12907-2	150
#7-3	227-1	12907-2	125
#8-3	227-1	12907-2	100
#9-3	227-1	12907-2	75
#10-3	227-1	12907-2	50
#11-3	227-1	12907-2	25
#12-3	227-1	12907-2	10
#1-4	232-1	12909-1	700
#2-4	232-1	12909-1	700
#3-4	232-1	12909-1	600
#4-4	232-1	12909-1	400
#5-4	232-1	12909-1	400
#6-4	232-1	12909-1	300
#7-4	232-1	12909-1	200
#8-4	232-1	12909-1	100
#9-4	232-1	12909-1	50
#10-4	232-1	12909-1	35
#11-4	232-1	12909-1	35
#12-4	232-1	12909-1	10
#1-5	234-2	12911-2	800
#2-5	234-2	12911-2	600
#3-5	234-2	12911-2	400
#4-5	234-2	12911-2	300
#5-5	234-2	12911-2	200
#6-5	234-2	12911-2	150
#7-5	234-2	12911-2	125

Table5 continued

#8-5	234-2	12911-2	100
#9-5	234-2	12911-2	75
#10-5	234-2	12911-2	50
#11-5	234-2	12911-2	25
#12-5	234-2	12911-2	10
#1-6	235-1	12912-1	800
#2-6	235-1	12912-1	600
#3-6	235-1	12912-1	400
#4-6	235-1	12912-1	300
#5-6	235-1	12912-1	200
#6-6	235-1	12912-1	150
#7-6	235-1	12912-1	125
#8-6	235-1	12912-1	100
#9-6	235-1	12912-1	75
#10-6	235-1	12912-1	50
#11-6	235-1	12912-1	25
#12-6	235-1	12912-1	10
#1-7	235-2	12912-2	2500
#2-7	235-2	12912-2	2500
#3-7	235-2	12912-2	2200
#4-7	235-2	12912-2	2000
#5-7	235-2	12912-2	1800
#6-7	235-2	12912-2	1800
#7-7	235-2	12912-2	1500
#8-7	235-2	12912-2	1300
#9-7	235-2	12912-2	1000
#10-7	235-2	12912-2	800
#11-7	235-2	12912-2	75
#12-7	235-2	12912-2	75
#1-8	238-2	12914-3	800
#2-8	238-2	12914-3	600
#3-8	238-2	12914-3	400
#4-8	238-2	12914-3	300
#5-8	238-2	12914-3	200
#6-8	238-2	12914-3	150
#7-8	238-2	12914-3	100
#8-8	238-2	12914-3	75
#9-8	238-2	12914-3	50
#10-8	238-2	12914-3	25
#11-8	238-2	12914-3	10

Preliminary results of the measurement on board

The PROPS sensor was working well during the whole cruise. The sensor performed very well also on the ROV. The recorded spectra were very stable and with a low noise. We could see a yellow-substance signal in some depth. Figure 10 shows the spectra recorded at 50 m, 100 m, 250 m and 400 m. The absorption at 220 nm increases with the depth, the lowest absorption is at 50 m.

The algorithm for the determination of the nitrate concentration did not work very well on board. One problem could be that the distilled water reference spectra was not at zero value after a series of measurements. Thus, we sometimes obtained negative nitrate concentrations. This problem has to be analyzed with the instruments manufacturer TriOS GmbH.




Fig. 10. Absorption amplitude vs. wavelength of a series of sample measurements with the PROPS sensor.

2.2. Marine Microbiology

2.2.1 Particle composition and size-related settling velocity (*Morten Iversen*)

At earlier cruises (POS-344 and MSM04b), we observed a sub-surface peak in the particle abundance. Such a peak has previously been described as an up-concentration of particles at density discontinuities. This is due to decreased settling velocity of the aggregates when they enter water masses with increasing density (McIntyre et al., 1995). However, when we compared the particle peak with the density discontinuities we did not find a strong correlation. The particle peak could form because particles with low density are laterally advected from the coastal water and into the open water (Nowald et al., 2006). To enable a better understanding of these observations, we made roller tank incubations to form aggregates. The incubations were done with water from the fluorescence maximum, from the sub-surface particle peak, and from the deep water below the particle peak. Roller tank incubation imitates the formation of marine snow aggregates, as it happens *in situ* via differential settling. Once the aggregates were formed, we measured size and settling velocity of individual aggregates. If the particle peak was due to lateral advection of particles from the coast, we expected to observe heterogeneity among the particles formed from water collected in the peak.

We only observed a clear sub-surface peak at station GeoB 12912. We formed aggregates from the water above, in, and below the peak and measured the settling velocity and size of the aggregates from the different depths (Fig. 11).




Fig. 11. Settling velocity versus size (equivalent spherical diameter) of the aggregates formed from water collected at GeoB 12912-1. Black circles are aggregates formed in water collected at 75 m (fluorescence maximum above the peak), red squares are aggregates formed in water from 1800 m (particle peak), and green triangles are aggregates formed in water collected at 2500 m depth (below the peak). The equations for the regression are also shown; from left to right for 75 m, 1800 m, and 2500 m.

We found a high heterogeneity in the settling velocity for the aggregates formed from the water collected within the particle peak (Fig. 11). This indicates that there is a different composition of material at this depth compared to the water above and below. Hence, our preliminary results point towards a lateral advection of particles which form the sub-surface particle peak observed off Cape Blanc. Though we did not observe a particle peak at all stations, we still made roller tank incubations and measurements of the aggregates formed at different depth (Table 6).

Table. 6. List of water samples taken for microbiological investigations.

Station	Sample depth	Sample volume	Analysis
GeoB #	(m)	(L)	
12907-4	35	20	POC content in water
	500	20	Aggregate formation
	2000	20	Aggregate size and settling velocity
12909-1	35	20	POC content in water
	400	20	Aggregate formation
	700	20	Aggregate size and settling velocity
12912-2	75	20	POC content in water
	1800	20	Aggregate formation
	2500	20	Aggregate size and settling velocity
12914 (pump)	3	50	POC content in water
			Aggregate formation
			Aggregate size and settling velocity

2.4. Oceanography and Marine Geology

2.4.1 CTD-O₂-chlorophyll-fluorescence probes (SBE-19 and SBE-3)

(*N. Nowald, C. Reuter, M. Klann, G. Fischer*)

Five CTD/O₂/chlorophyll-fluorescence profiles were taken with a self-contained SBE 19 profiler equipped with a conductivity-temperature-depth probe plus oxygen sensor and a CHELSEA-fluorometer. This CTD was attached to the frame of the ParCa system and was deployed five times during the cruise (see Table 7). The data were removed immediately after recovery of the system and standard downcast plots were made. Data were compared to the measurements with the shipboard SBE 3-CTD (10 profiles were acquired, see station list) which was equipped with a chlorophyll fluorescence and two oxygen sensors. Salinity and temperature data fitted perfectly between the two CTD systems (Fig. 12). The oxygen values of the SBE-3 were slightly higher than of the SBE-19 profiler and fitted almost perfectly to measured oxygen values (Winkler titration performed by the ICCM, see above) (Fig. 12).

Table 7. List of CTD-O₂-chlorophyll-fluorescence profiles taken with the SBE-19 profiler.

GeoB station #	Date 2008	Time (UTC)	LAT N	LONG W	Water depth (m)	Deployment depth (m)
12907-3	20.04	21:49	21°16.9'	20°47.8'	4154	2000
12910-1	23.04.	07:43	20°43.3'	18°42.3'	Ca. 2700	2500
12911-3	23.04	20:59	20°04.7'	18°58.7'	Ca. 3012	2500
12913-1	24.04.	20:03	20°35.0'	18°00.0'	766	750
12914-4	25.04.	20:38	21°19.9'	18°49.7'	3062	2000




Fig. 12. Downcast CTD-profiles taken at the eutrophic mooring site CBi with the the SBE-19 (a: attached to the ParCa frame, see below) and the shipboard instrument SBE-3 (b). Oxygen from Winkler titration (ICCM, see above) is also shown in b (black dots) and cooresponds nicely to the measured values. Note the good correspondance between the two instruments in S and T, chlorophyll fluoerescence, however, is too low in b (SBE-3).

2.4.2. Sampling for chlorophyll-a measurements (*M. Klann, G. Fischer*)

For the determination of chlorophyll-a concentrations in the surface waters, seawater from the shipboard installed seawater pump ("Aquariumpumpe") was sampled when sailing (Table 8). Half a liter or one liter of seawater were filtered onto a glass microfibre filter (Whatman, GF/F, 25 mm diameter). The samples were frozen at dark and will be analyzed by means of photometry at the laboratory in Bremen. The results from the samples from the shipboard pump will be compared with satellite-derived chlorophyll concentration maps (SeaWiFs, MODIS) and may serve as calibration of these data.

Table 8. Sampling for chlorophyll-a measurements (shipboard aquarium pump). T-S-data were taken from the ships' thermosalinograph.


Sample #	Date 2008	Time (UTC)	Latitude N	Longitude W	Water depth (m)	Salinity (‰)	SST (°C)	Volume (l)
1	20.04.	10:42	21°15.4262	20°49.2300	4154	35,13	20,77	2
2	20.04.	10:42	21°15.4262	20°49.2300	4154	35,13	20,77	2
3	22.04.	17:20	20°34.6920	18°00.102	770,3	34,67	17,16	2
4	22.04.	17:20	20°34.6920	18°00.102	770,3	34,67	17,16	2
5	23.04.	07:15	20°43.2708	18°42.3558	no data	35,07	20,31	2
6	23.04.	07:15	20°43.2708	18°42.3558	no data	35,07	20,31	2
7	23.04.	18:07	20°04.6860	18°58.7352	no data	34,62	19,12	1,5
8	23.04.	18:07	20°04.6860	18°58.7352	no data	34,62	19,12	1,5
9	24.04.	19:36	20°34.9920	18°00.0282	770	34,73	17,58	1,5
10	24.04.	19:36	20°34.9920	18°00.0282	770	34,73	17,58	1,5
11	25.04.	19:56	21°19.9212	18°48.6680	3060	35,15	19,81	1,5
12	25.04.	00:00	21°19.9212	18°48.6680	3060	35,15	19,81	1,2

2.4.3. Particle fluxes measured with sediment traps

(N. Nowald, C. Reuter, M. Klann, G. Fischer, G. Ruhland)

One aim of this part of the cruise was to recover and redeploy the moorings CI-21/22 (=ESTOC) located north of the Canary Islands in a mesotrophic/oligotrophic setting where sediment traps have been deployed since 1991. It was also planned to recover and redeploy mooring CB18/19 which is located about 200 nm off Cape Blanc (Mauritania, Fig. 1). This mesotrophic study site operated since 1988 is located at the edge of the Cape Blanc filament in about 4100 m water depth. It is used to monitor the long-term change of particle fluxes in the Mauritanian offshore upwelling zone. An additional mooring named CBi-5 was deployed during MERIAN MSM 04b cruise around 80 nm further to the east and was also planned to be exchanged (CBi-6, Fig. 1). The data of deployments and recoveries of the moorings are listed in Table 8 alongside with the sampling data of the traps.

On April 15th the mooring CI-21 equipped with 3 particle traps had been recovered successfully. All traps had worked perfectly, delivering 20 samples each. The mooring was redeployed in a comparable configuration in the afternoon of the same day as CI-22 at the same location (Fig. 13).



Mooring: CI-22

Position: 29°02.83'N 015° 15.94'W
Wasserdepth: 3589 m
Deployment: 15.04.2008
Voyage: Poseidon Pos-365

Releaser codes #878: Enable 5141 / Disable 5143
 Rel 5149
 #:901 Enable 5134 / Disable 5126
 Rel 5140



Fig. 13. Drawing of the ESTOC mooring CI-22 deployed north of the Canary Islands.

On April 20th, we successfully recovered mooring CB-18 ca. 200 nm off Cape Blanc which had worked perfectly. It was redeployed as CB-19 with a similar configuration on April 21. In the morning of April 23, we released the 1500 m long mooring array CBI-5 in the coastal part of the Cape Blanc filament which was equipped with two traps including one MSD trap with 40 cups and one current meter. We received one complete set of 40 samples of CBI-5 from the upper water column, the lower trap had failed. On April 24th, we could redeploy the mooring array CBI-6 (with a MSD trap of 40 cups), now equipped with a newly developed MSD platform with a FSI-CTD and a video camera system. The latter should monitor particle distribution and transport as well as

particle sizes over an annual cycle. It is planned to recover and redeploy the mooring arrays with RV MERIAN in spring 2009.

Table 9. Data for recoveries and redeployments of the sediment trap mooring arrays.

Mooring	Position	Water Depth (m)	Interval	Instr. (m)	Depth (no x days)	Intervals
<u>Mooring recoveries</u>						
CI-21/ESTOC:	29°02,6' N 15°15,7. W	3605	22.03.07- 05.04.08	SMT 234 SMT 234 SMT 234	702 1109 3066	20 x 19 20 x 19 20 x 19
<u>Cape Blanc mesotrophic:</u>						
CB-18	21°16,9' N 20°48,0' W	4168	25.03.07- 05.04.08	SMT 243 SMT 234	1222 3629	20 x 19 20 x 19
<u>Cape Blanc eutrophic:</u>						
CBi-5	20°44,9' N 18°41,9' W	2709	28.03.07- 05.04.08	MSD SMT 234 RCM8	1263 1872 1317	38 x 9,5, 2 x 6,5 20 x 19
<u>Mooring deployments:</u>						
CI-22/ESTOC:	29°02,8' N 15°15,9. W	3605	16.04.08- 22.03.09	SMT 234 NE SMT 234 NE SMT 234	702 1109 3066	20 x 17 20 x 17 20 x 17
<u>Cape Blanc mesotrophic:</u>						
CB-19	21°16,2' N 20°48,7' W	4155	22.04.08- 22.03.09	SMT 230Ti NE SMT 238	1209 3617	1 x 11, 19 x 17 20 x 17
<u>Cape Blanc eutrophic:</u>						
CBi-6	20°45,1' N 18°41,9' W	2699	25.04.08- 22.03.09	MSD platfrom MSD trap SMT 243 Ti NE	1189 1299 1869	2 x 4, 38 x 8,5 1 x 8, 19 x 17

Instruments used:

- SMT 230/241/243/234 = Sediment trap (Ti=Titanium trap), Aquatec Meerestechnik, Kiel
 RCM 8 = Aanderaa current meter, RCM 8
 SMT 230/234/238 NE = new electronics
 MSD = Mult-Sensor-Device trap with 40 cups
 MSD platfrom = platform with FSI-CTD and video camera

2.4.4. Particle distribution measured with optical systems (ParCa)

(N. Nowald, C. Reuter, M. Klann)

System description

The photographic particle camera system ParCa was deployed on 5 locations for the in-situ measurement of the vertical size distribution and concentration of particulate matter in the ocean ('marine snow'). ParCa consists of a modified NIKON CoolPix 995 digital camera. A strobe, mounted perpendicular to the optical axis of the camera, provides a collimated light beam of 12 cm width, illuminating a defined sample volume. Power Source is a 24V/38 Ah rechargeable lead battery designed for the use to full ocean depth. ParCa can operate in depths up to 3000 m. All devices are mounted in a 200 kg galvanised frame. Communication with the ship is done by a microcontroller and adapted software. An additionally installed SeaBird PDIM telemetry provides full control of the entire system, via the ships coaxial wire. Pictures were exposed while lowering

the system at a speed of 0.5 m/sec at approx. each 12 m of depth. A detailed station list is given below (Table 10).

Table 10. List of stations done with the ParCa system combined with the SBE 19-CTD.

GeoB #	Date	Time seafloor/ max. wire –length (UTC)	Latitude	Longitude	Water depth (m)	Deployment depth
2008						
12907	20.04.	09:45	21°16.9 N	20°47.8 W	4152	Down to 2000
12910	23.04.	06:30	20°43.3 N	18°42.3 W	~2650	Down to 2500m
12911	23.04.	19:30	20°04.7 N	18°58.7 W	3016	Down to 2500m
12913	24.04.	?	20°35.0 N	18°00.0 W	766	Down to 750m
12914	25.04.	19:40	21°19.9 N	18°49.6 W	3062	Down to 2000m

Preliminary results

Although a distinct surface maximum is found everywhere, concentrations vary in the upper ocean surface between the different stations (around 100 and 300 particles per image) (Fig. 14). Below the euphotic zone, the distribution of particulate matter is highly variable. Some characteristic distribution patterns, repeatedly observed in this area, were also found during this cruise. Station GeoB 12911, shows an extensive subsurface particle maximum around 300 m water depth. This kind of particle distribution is common off Cape Blanc and is found in many particle profiles acquired in this area. The same is true or the distinct deep water maximum (1500 m – 2200 m) at station GeoB 12910 (CBi trap mooring site). This characteristic maximum was tracked for three years up to now at this specific site and appears to be a common feature.




Fig. 14. Abundances of particles as number per image in the water column off Cape Blanc as derived from the ParCa system.

2.4.5. Particle investigations with the ROV *Cherokee*

(*N. Nowald, C. Reuter, G. Ruhland, M. Klann, E. Kopiske*)

System description

The Remotely Operated Vehicle (ROV) *Cherokee* is a commercially available, midsize inspection class ROV, manufactured by *Sub-Atlantic*, Aberdeen. It is operated by MARUM since 2001 and

was adapted and enhanced for scientific purposes. The vehicle was involved in many different research programs and cruises, performing more than 75 dives up to now.

The *Cherokee* ROV system consists of three major components: a spooling winch, the Surface Control Racks and the vehicle itself. Vehicle dimensions are 120x80x100 cm (LxBxH) and weight in air is around 450 kg. The *Cherokee* has a payload for scientific equipment of approx. 20 kg. The ROV is electrically propelled by 4 axial thrusters and total power of the system is 12 kW. *Cherokee* is 1000m depth rated, but due to several “cut offs” and terminations of the umbilical supply cable, only a diving depth of 850 m is guaranteed. Four video cameras are mounted on the ROV for observation and navigational purposes. A color video zoom camera (720x576 lines), a modified digital Nikon still camera (3.2 Megapixel) with associated flash light and two mini video cameras for the overview to the front and back area of the vehicle. The video-, and the still image camera are mounted onto a Pan & Tilt unit, which enhances the observation capabilities of the vehicle. The Pan & Tilt unit also carries 3 lasers for object size measurements on the seafloor. Underwater light is provided by 3x230W DSPL dimmable spots. For long or close range obstacle detection and measurement, a *Tritech Seaking* dual frequency sonar is mounted on the port side of the vehicle. It displays an acoustical realtime image on the topside sonar PC. The sonar operates at 325/675 Hz with a maximum scanning range of 300 m. Sonar screenshots and recordings are possible. Navigational devices such as compass, altimeter and depth sensor, are part of the basic sensor package onboard the ROV.

For scientific sampling and experiments, a small hydraulic manipulator system is used. The *Hydro-Lek* HLK-EH-5 is a non-proportional, 5 function manipulator, powered and controlled by a combined pressure pump and 6 station valve pack. Operating pressure is 130 bar and lifting capacity is 25 kg. Part of the hydraulical system is the toolbox, which is used for storing samples and/or mounting sampling tools. It can be hydraulically opened and closed. Standard sampling tools during RV PELAGIA cruise 64PE284 were a 1 l Niskin bottle and a sampling net with a diameter of 140 mm and a mesh size of 1 mm. The supply cable (Umbilical) contains 20 electrical conductors providing electrical power and basic telemetry. In addition, 4 optical multimode fibres provide 4 video and 4xRS232 channels.

Control over the system is given by three 19” racks, equipped with several display and recording devices. The *Surface Control Unit* contains the remote control for the vehicle and the manipulator. It also contains the sonar PC and screen, two small 10” color monitors for video camera display as well as an ampere-, and voltmeter. The *Pilot Rack*’s PC monitor shows navigational data from the ship and the ROV, such as heading, position, depth and so on. A large analogue monitor inside the rack displays the live video of the *Tritech* color zoom camera. The *Video Rack* includes 2 DV

recorders to record two cameras simultaneously. The rack also has a PC with software running to control the *Nikon* still image camera, the *Ifremer* framegrab utility *Adelie* and the USBL positioning system GAPS (*IXSEA*).

Scientific payload

The vehicle was equipped with a *Settling Chamber* in order to measure sinking velocities of marine particles in different water depths (Fig. 15). The box is made of *Plexiglas* and its dimensions are 30x60x30 cm (LxBxH). It can be opened and closed sideways, with aid of the ROV's manipulator. For accurate measurements, a collimated light source providing a slab of light of 12 cm width, is mounted on the port side of the vehicle. A caliper fixed in the middle of the slab of light, helps in measuring the distance a particle travelled through time within the closed box and calculating the total sample volume. During the cruise, the vehicle was deployed at station GeoB 12907. Recordings of sinking material were acquired in four different depths (50 m, 150 m, 250 m, 400 m). The video recordings will be analysed at the University of Bremen.



Fig. 15. The *Settling Chamber* mounted in front of the ROV *Cherokee*.

2.5. Marine Techology

2.5.1. Recovery of the DOLAN buoy

(G. Ruhland, E. Kopiske, S. Klar, K. Dehning, C. Reuter)

The Surface Buoy Unit (SBU) operates since 1997 and was formerly part of the DOMEST project, later it was used in the MERSEA project. The unit can carry several meteorological sensors, satellite telemetry links and sub-sea telemetry links like an ORCA acoustic modem as well as a cable telemetry down to 100 m. Due to the end of the project, the buoy was recovered but not redeployed. The recovery has been done on April 14th without any problems and all sensors and the buoy itself were in a good condition. The last configuration of the system is shown in Figure 16.




Fig. 16. Configuration of the Surface Buoy Unit (SBU) in the last deployment interval.

Additionally to the usually used equipment an underwater modem was installed to establish an hourly acoustic communication link to the nearby placed PACT mooring.

2.5.2. Recovery of the PACT mooring

(*G. Ruhland, E. Kopiske, S. Klar, K. Dehning, C. Reuter*)

In a distance of about one nautical mile to the SBU, a mooring was deployed on POS360 which was used for trials of PACT (Pressure Acoustically Coupled Tsunameter) units. The bottom unit of the PACT system which monitored changes in pressure was installed in a water depth of 3600 m. The HAM.Node receiver modem was mounted to the SBU mooring in 12 m depth and connected to an IRIDIUM satellite transceiver residing on top of the buoy. Data received from the PACT bottom unit was relayed to the AWI/Bremerhaven via a satellite link. The satellite link also provides basic diagnostic functionality (like pinging the bottom unit, information about battery status etc.). Figure 17 shows the complete series of messages which were sent during the test interval. Usually one message per hour was transmitted to save battery power. In between simulated tsunami alarms triggered a higher rate of transmissions.




Fig. 17. Pressure data telegrams as hourly routine and simulated tsunami alarm messages.

2.6. References

- Garside, C. (1993): Nitrate reductor efficiency as an error source in seawater analysis. *Mar. Chem.*, 4(1), 25-30.
- Koroleff, F., 1983. Determination of dissolved inorganic phosphate. In Methods of Seawater Analysis. K. Grasshoff, A. Ehrhardt and K. Kremling (eds), Verlag Chemie, 126-129.
- _____, 1983b. Determination of dissolved inorganic silicate. In Methods of Seawater Analysis. K. Grasshoff, A. Ehrhardt and K. Kremling (eds), Verlag Chemie, 175-180.
- Kremling, K. & A. Wenck, 1986. On the storage of dissolved inorganic phosphate, nitrate and reactive silicate in Atlantic Ocean water samples. *Meeresforsch.*, 31, 69-74.
- Llinás O.,M.J. Rueda, J.Pérez-Marrero,E.Pérez-Martell, R.Santana, M.G.Villagarcía, A.Cianca, J.Godoy and L.Maroto, 2002. Variability of the Antarctic Intermediate Waters in the Northern Canary Box. *Deep Sea Research. II*,49(17):3441-3453.
- McDonald, R.W. & F.A. McLaughlin, 1982. The effect of storage by freezing of dissolved inorganic phosphate, nitrate and reactive silicate for samples from coastal and internal water. *Water Research*, 16, 95-104.
- Murphy, J. & J.P. Riley, 1962. A modified single solution method for the determination of phosphate in natural waters. *Anal. Chim. Acta*, 27, 31-36.
- Nydahl, F., 1976. On the optimum conditions for the reduction of nitrate by cadmium. *Talanta*, 23, 349-357.
- Riley, J.P. & J.P. Skirpow, 1975. The Micronutrient Element. *Chemical Oceanography*, 2, 245-297.
- Strickland, J.D.H & Parsons, 1972. A practical handbook of seawater analysis. *Fisheries Research Board of Canada*, 167 pp.
- UNESCO (1978): Technical Papers in Marine Science, 28, 35pp.
- UNESCO (1984): La escala de salinidades practicas de 1978 y la ecuacion internacional de estado del agua de mar de 1980. Documentos tecnicos de la Unesco sobre Ciencias del Mar, no. 36.
- Welschmeyer, N.A., 1994. Fluorometric Analysis of Chlorophyll "a" in presence of Chlorophyll "b" and Phaeopigments. *Limnol. Oceanog.* 39 (8), 1985-1992.
- Woods, E.D., F.A.J. Armstrong & F.A. Richards (1967). Determination of nitrate in seawater by cadmium-cooper reduction to nitrate. *J. Mar. Biol. Ass. UK.*,47, 31-43.
- Wright S., Jeffrey S. (2006) Hbd Env Chem 2:71-104
- Wright S, Jeffrey S., Mantoura R., Llewellyn C., Bjornland T., Repeta D. and Welschmeyer N. (1991). Mar Ecol Prog Series, 77:183-196
- Scientific Committee on Oceanic Research. Commission, I.O. (ed.). (1994). Measurement of Algal Chlorophylls and Carotenoids by HPLC. Protocols for the Joint Global Ocean Flux Study (JGOFS) core measurements. Chapter 13, pp. 91-100.
- MacIntyre et al. 1995 – Limnology and Oceanography 40:449-468
- Nowald et al. 2006 – Ocean Sci. Discuss 3:903-938
- Welschmeyer, 1994 (ICCM)

2.7. Acknowledgements

This cruise was funded by the DFG Research Center and the Excellence Cluster at Marum and the Europrox PhD programm. The RV POSEIDON was an excellent platform to perfom our studies during the cruise. We thank Captain M. Schneider and his entire crew for their professional and friendly support that was the basis for a successful cruise. We also like to thank Tom Müller (IfM-Geomar) for his support during the planning phase of the cruise.

3. Station List (POS 365 Leg 1 and 2)

GeoB #	Ships #	Date 2008	Device	Time seafloor/ max. wire -length [UTC]	Latitude N	Longitude W	Water depth [m]	Recovery/Remarks
365-1								
12901-1	212	13.04.	PACT	16:35	29°11.48'	15°54.99'	3630	Recovery of PACT mooring, release
12901-2	212-2		ROS-CTD	19:12	29°11.48'	15°56.02'	3630	Down to 1000m: casts at 1000,200,150,100,90,80,70,55,40,25,10 m
12902-1	213	14.04.	DOLAN	07:14	29°10.89'	15°55.01'	3628	Recovery of DOLAN buoy started
12902-2	214		XBT	12:26	29°13.97'	15°53.69'	3630	Down to 1830m
12903-1	215		ROS-CTD	17:30	29°10.06'	15°20.00'	3595	Down to 3500m: casts at 3500,3000,2800,2500,2000,1800,1500,1300, 1200,1100,1000,800 m
12904-1	216	15.04	ESTOC-CI	07:06	29°02.55'	15°15.71'	3589	Recovery of mooring, 3 traps worked
12905-1	217		ROS-CTD	11:54	29°09.99'	15°20.03'	3594	Down to 800m: casts at: 800,600,400,300,200,150,125,100,75,50,25,10 m
12905-2	217-2		NOAA	12:23	29°10.01'	15°20.03'	3595	Deployment of NOAA drifter
12906-1	218		ESTOC-CI-22	17:00	29°02.83'	15°15.94'	3589	Deployment of ESTOC-CI-22 mooring with 3 traps
365-2								
12907-1	226	20.04.	CB-18	14:31	21°16.50'	20°48.39'	4156	Both traps worked
12907-2	227		ROS-CTD	19:23	21°16.87'	20°48.11'	4156	Down to 800m: casts at: 800,600,400,300,200,150,125,100,75,50,25,10 m
12907-3	227-2		Parca-CTD	21:49	21°16.86'	20°47.83'	4154	Down to 2000m
12907-4	228	21.04.	ROS-CTD	05:51	21°16.79'	20°47.97'	4159	Down to 2000m casts at: 2000(2x),1800,1600,1400,1200, 1000,800,500(2x),35(2x) m
12907-5	229	.	CB-19	07:54	21°14.04'	20°50.55'	4238	Start deployment
12908-1	230		XBT	19:52	20°59.97'	19°40.47'	3639	Down to 1800m
12909-1	231	22.04.	ROS-CTD	17:48	20°34.91'	18°00.11'	768	Down to 700m Casts at: 700(2x),600,400(2x),300,200,100,50, 35(2x), 10 m
12910-1	232	23.04.	Parca-CTD	07:43	20°43.25'	18°42.31'	Ca.270 0	Down to 2500m
12910-2	233		CBi-5	09:05	20°44.59'	18°41.99'	Ca.270 0	mooring released, MSD at #40, lower trap failed

GeoB #	Ships #	Date 2008	Device	Time seafloor/ max. wire -length [UTC]	Latitude N	Longitude W	Water depth [m]	Recovery/Remarks
12911-1	234		XBT	17:40	20°03.62'	18°59.39'	3013	Down to 1800m
12911-2	234-2		ROS-CTD	18:20	20°04.66'	18°58.74'	3012	Down to 800m Casts at: 00,600,400,300,200,150,125,100,75,50,25,10 m
12911-3	234-3		Parca-CTD	20:59	20°04.70'	18°58.71'	Ca.301 2	Down to 2500m
12912-1	235	24.04.	ROS-CTD	07:24	20°44.97'	18°41.95'	2721	Down to 800m Casts at:800,600,400,300,200,150,125,100,75,50,25,10 m
12912-2	235-2		ROS-CTD	09:09	20°44.98'	18°41.98'	Ca.272 1	Down to 2500m Casts at: 2500(2x),2200,2000,1800(2x),1500,1300,1000 800,75(2x) m
12912-3	235-3		CBi-6	12:00	20°44.13'	18°42.16'	2700	Start of deployment
12913-1	236		Parca-CTD	20:03	20°34.99'	18°00.01'	766	Down to 750m
12914-1	237	25.04.	XBT	11:53	21°18.63'	18°49.61'	3054	Down to 1800m
12914-2	238		ROV	12.23	21°20.11'	18°49.89'	3061	Deployment, down to 400m; with nutrient sensor
12914-3	238-2		ROS-CTD	18:59	21°19.82'	18°49.59'	3059	Down to 800m casts at: 800,600,400,300,200,150,125,100,75,50,25,10 m
12914-4	238-3		Parca-CTD	20:38	21°19.91'	18°49.66'	3062	Down to 2000m

CI (ESTOC), CB, CBi - Sediment trap mooring sites north of Gran Canaria (CI/ESTOC) and off Cape Blanc (Mauritania)

ROS-CTD – Multi-water sampler (rosette) with 12x10l bottles and CTD-SBE 3 (IFM-GEOMAR)

PARCA-CTD– Particle Camera System with CTD SBE 19 inside frame (CTD-O₂- chlorophyll-fluorescence (SBE-19, #2069)

ROV – Cherokee, with nitrate sensor and settling chamber

PACT – tsunameter mooring

DOLAN – instrument bouy

XBT – extended bathythermograph

Publications of this series:

- No. 1 Wefer, G., E. Suess and cruise participants**
Bericht über die POLARSTERN-Fahrt ANT IV/2, Rio de Janeiro - Punta Arenas, 6.11. - 1.12.1985.
60 pages, Bremen, 1986.
- No. 2 Hoffmann, G.**
Holozänstratigraphie und Küstenlinienverlagerung an der andalusischen Mittelmeerküste.
173 pages, Bremen, 1988. (out of print)
- No. 3 Wefer, G. and cruise participants**
Bericht über die METEOR-Fahrt M 6/6, Libreville - Las Palmas, 18.2. - 23.3.1988.
97 pages, Bremen, 1988.
- No. 4 Wefer, G., G.F. Lutze, T.J. Müller, O. Pfannkuche, W. Schenke, G. Siedler, W. Zenk**
Kurzbericht über die METEOR-Expedition No. 6, Hamburg - Hamburg, 28.10.1987 - 19.5.1988.
29 pages, Bremen, 1988. (out of print)
- No. 5 Fischer, G.**
Stabile Kohlenstoff-Isotope in partikulärer organischer Substanz aus dem Südpolarmeer
(Atlantischer Sektor). 161 pages, Bremen, 1989.
- No. 6 Berger, W.H. and G. Wefer**
Partikelfluß und Kohlenstoffkreislauf im Ozean.
Bericht und Kurzfassungen über den Workshop vom 3.-4. Juli 1989 in Bremen.
57 pages, Bremen, 1989.
- No. 7 Wefer, G. and cruise participants**
Bericht über die METEOR - Fahrt M 9/4, Dakar - Santa Cruz, 19.2. - 16.3.1989.
103 pages, Bremen, 1989.
- No. 8 Kölling, M.**
Modellierung geochemischer Prozesse im Sickerwasser und Grundwasser.
135 pages, Bremen, 1990.
- No. 9 Heinze, P.-M.**
Das Auftriebsgeschehen vor Peru im Spätquartär. 204 pages, Bremen, 1990. (out of print)
- No. 10 Willems, H., G. Wefer, M. Rinski, B. Donner, H.-J. Bellmann, L. Eißmann, A. Müller,
B.W. Flemming, H.-C. Höfle, J. Merkt, H. Streif, G. Hertweck, H. Kuntze, J. Schwaar,
W. Schäfer, M.-G. Schulz, F. Grube, B. Menke**
Beiträge zur Geologie und Paläontologie Norddeutschlands: Exkursionsführer.
202 pages, Bremen, 1990.
- No. 11 Wefer, G. and cruise participants**
Bericht über die METEOR-Fahrt M 12/1, Kapstadt - Funchal, 13.3.1990 - 14.4.1990.
66 pages, Bremen, 1990.
- No. 12 Dahmke, A., H.D. Schulz, A. Kölling, F. Kracht, A. Lücke**
Schwermetallspuren und geochemische Gleichgewichte zwischen Porenlösung und Sediment
im Wesermündungsgebiet. BMFT-Projekt MFU 0562, Abschlußbericht. 121 pages, Bremen, 1991.
- No. 13 Rostek, F.**
Physikalische Strukturen von Tiefseesedimenten des Südatlantiks und ihre Erfassung in
Echolotregistrierungen. 209 pages, Bremen, 1991.
- No. 14 Baumann, M.**
Die Ablagerung von Tschernobyl-Radiocäsium in der Norwegischen See und in der Nordsee.
133 pages, Bremen, 1991. (out of print)
- No. 15 Kölling, A.**
Frühdiagenetische Prozesse und Stoff-Flüsse in marinen und ästuarienen Sedimenten.
140 pages, Bremen, 1991.
- No. 16 SFB 261 (ed.)**
1. Kolloquium des Sonderforschungsbereichs 261 der Universität Bremen (14.Juni 1991):
Der Südatlantik im Spätquartär: Rekonstruktion von Stoffhaushalt und Stromsystemen.
Kurzfassungen der Vorträge und Poster. 66 pages, Bremen, 1991.
- No. 17 Pätzold, J. and cruise participants**
Bericht und erste Ergebnisse über die METEOR-Fahrt M 15/2, Rio de Janeiro - Vitoria,
18.1. - 7.2.1991. 46 pages, Bremen, 1993.
- No. 18 Wefer, G. and cruise participants**
Bericht und erste Ergebnisse über die METEOR-Fahrt M 16/1, Pointe Noire - Recife,
27.3. - 25.4.1991. 120 pages, Bremen, 1991.
- No. 19 Schulz, H.D. and cruise participants**
Bericht und erste Ergebnisse über die METEOR-Fahrt M 16/2, Recife - Belem, 28.4. - 20.5.1991.
149 pages, Bremen, 1991.

- No. 20** **Berner, H.**
Mechanismen der Sedimentbildung in der Fram-Straße, im Arktischen Ozean und in der Norwegischen See. 167 pages, Bremen, 1991.
- No. 21** **Schneider, R.**
Spätquartäre Produktivitätsänderungen im östlichen Angola-Becken: Reaktion auf Variationen im Passat-Monsun-Windsystem und in der Advektion des Benguela-Küstenstroms.
198 pages, Bremen, 1991. (out of print)
- No. 22** **Hebbeln, D.**
Spätquartäre Stratigraphie und Paläozeanographie in der Fram-Straße. 174 pages, Bremen, 1991.
- No. 23** **Lücke, A.**
Umsetzungsprozesse organischer Substanz während der Frühdiagenese in ästuarienen Sedimenten. 137 pages, Bremen, 1991.
- No. 24** **Wefer, G. and cruise participants**
Bericht und erste Ergebnisse der METEOR-Fahrt M 20/1, Bremen - Abidjan, 18.11.- 22.12.1991.
74 pages, Bremen, 1992.
- No. 25** **Schulz, H.D. and cruise participants**
Bericht und erste Ergebnisse der METEOR-Fahrt M 20/2, Abidjan - Dakar, 27.12.1991 - 3.2.1992.
173 pages, Bremen, 1992.
- No. 26** **Gingele, F.**
Zur klimabedingten Bildung biogener und terrigener Sedimente und ihrer Veränderung durch die Frühdiagenese im zentralen und östlichen Südatlantik. 202 pages, Bremen, 1992.
- No. 27** **Bickert, T.**
Rekonstruktion der spätquartären Bodenwasserzirkulation im östlichen Südatlantik über stabile Isotope benthischer Foraminiferen. 205 pages, Bremen, 1992. (out of print)
- No. 28** **Schmidt, H.**
Der Benguela-Strom im Bereich des Walfisch-Rückens im Spätquartär. 172 pages, Bremen, 1992.
- No. 29** **Meinecke, G.**
Spätquartäre Oberflächenwassertemperaturen im östlichen äquatorialen Atlantik.
181 pages, Bremen, 1992.
- No. 30** **Bathmann, U., U. Bleil, A. Dahmke, P. Müller, A. Nehrkorn, E.-M. Nöthig, M. Olesch, J. Pätzold, H.D. Schulz, V. Smetacek, V. Spieß, G. Wefer, H. Willems**
Bericht des Graduierten Kollegs. Stoff-Flüsse in marin Geosystemen.
Berichtszeitraum Oktober 1990 - Dezember 1992. 396 pages, Bremen, 1992.
- No. 31** **Damm, E.**
Frühdiagenetische Verteilung von Schwermetallen in Schlicksedimenten der westlichen Ostsee.
115 pages, Bremen, 1992.
- No. 32** **Antia, E.E.**
Sedimentology, Morphodynamics and Facies Association of a mesotidal Barrier Island Shoreface (Spiekeroog, Southern North Sea). 370 pages, Bremen, 1993.
- No. 33** **Duinker, J. and G. Wefer (ed.)**
Bericht über den 1. JGOFS-Workshop. 1./2. Dezember 1992 in Bremen. 83 pages, Bremen, 1993.
- No. 34** **Kasten, S.**
Die Verteilung von Schwermetallen in den Sedimenten eines stadtbrémischen Hafenbeckens.
103 pages, Bremen, 1993.
- No. 35** **Spieß, V.**
Digitale Sedimentographie. Neue Wege zu einer hochauflösenden Akustostratigraphie.
199 pages, Bremen, 1993.
- No. 36** **Schinzel, U.**
Laborversuche zu frühdiagenetischen Reaktionen von Eisen (III) - Oxidhydraten in marin Sedimenten. 189 pages, Bremen, 1993.
- No. 37** **Sieger, R.**
CoTAM - ein Modell zur Modellierung des Schwermetalltransports in Grundwasserleitern.
56 pages, Bremen, 1993. (out of print)
- No. 38** **Willems, H. (ed.)**
Geoscientific Investigations in the Tethyan Himalayas. 183 pages, Bremen, 1993.
- No. 39** **Hamer, K.**
Entwicklung von Laborversuchen als Grundlage für die Modellierung des Transportverhaltens von Arsenat, Blei, Cadmium und Kupfer in wassergesättigten Säulen. 147 pages, Bremen, 1993.
- No. 40** **Sieger, R.**
Modellierung des Stofftransports in porösen Medien unter Ankopplung kinetisch gesteuerter Sorptions- und Redoxprozesse sowie thermischer Gleichgewichte. 158 pages, Bremen, 1993.

- No. 41** **Thießen, W.**
Magnetische Eigenschaften von Sedimenten des östlichen Südatlantiks und ihre paläoceanographische Relevanz. 170 pages, Bremen, 1993.
- No. 42** **Spieß, V. and cruise participants**
Report and preliminary results of METEOR-Cruise M 23/1, Kapstadt - Rio de Janeiro, 4.-25.2.1993.
139 pages, Bremen, 1994.
- No. 43** **Bleil, U. and cruise participants**
Report and preliminary results of METEOR-Cruise M 23/2, Rio de Janeiro - Recife, 27.2.-19.3.1993
133 pages, Bremen, 1994.
- No. 44** **Wefer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 23/3, Recife - Las Palmas, 21.3. - 12.4.1993
71 pages, Bremen, 1994.
- No. 45** **Giese, M. and G. Wefer (ed.)**
Bericht über den 2. JGOFS-Workshop. 18./19. November 1993 in Bremen.
93 pages, Bremen, 1994.
- No. 46** **Balzer, W. and cruise participants**
Report and preliminary results of METEOR-Cruise M 22/1, Hamburg - Recife, 22.9. - 21.10.1992.
24 pages, Bremen, 1994.
- No. 47** **Stax, R.**
Zyklische Sedimentation von organischem Kohlenstoff in der Japan See: Anzeiger für Änderungen von Paläooceanographie und Paläoklima im Spätkänozoikum.
150 pages, Bremen, 1994.
- No. 48** **Skowronek, F.**
Frühdiagenetische Stoff-Flüsse gelöster Schwermetalle an der Oberfläche von Sedimenten des Weser Ästuars. 107 pages, Bremen, 1994.
- No. 49** **Dersch-Hansmann, M.**
Zur Klimaentwicklung in Ostasien während der letzten 5 Millionen Jahre:
Terrigener Sedimenteneintrag in die Japan See (ODP Ausfahrt 128). 149 pages, Bremen, 1994.
- No. 50** **Zabel, M.**
Frühdiagenetische Stoff-Flüsse in Oberflächen-Sedimenten des äquatorialen und östlichen Südatlantik. 129 pages, Bremen, 1994.
- No. 51** **Bleil, U. and cruise participants**
Report and preliminary results of SONNE-Cruise SO 86, Buenos Aires - Capetown, 22.4. - 31.5.93
116 pages, Bremen, 1994.
- No. 52** **Symposium: The South Atlantic: Present and Past Circulation.**
Bremen, Germany, 15 - 19 August 1994. Abstracts. 167 pages, Bremen, 1994.
- No. 53** **Kretzmann, U.B.**
57Fe-Mössbauer-Spektroskopie an Sedimenten - Möglichkeiten und Grenzen.
183 pages, Bremen, 1994.
- No. 54** **Bachmann, M.**
Die Karbonatrampe von Organya im oberen Oberapt und unteren Unteralb (NE-Spanien, Prov. Lerida): Fazies, Zyklo- und Sequenzstratigraphie. 147 pages, Bremen, 1994. (out of print)
- No. 55** **Kemle-von Mücke, S.**
Oberflächenwasserstruktur und -zirkulation des Südostatlantiks im Spätquartär.
151 pages, Bremen, 1994.
- No. 56** **Petermann, H.**
Magnetotaktische Bakterien und ihre Magnetosome in Oberflächensedimenten des Südatlantiks.
134 pages, Bremen, 1994.
- No. 57** **Mulitza, S.**
Spätquartäre Variationen der oberflächennahen Hydrographie im westlichen äquatorialen Atlantik.
97 pages, Bremen, 1994.
- No. 58** **Segl, M. and cruise participants**
Report and preliminary results of METEOR-Cruise M 29/1, Buenos-Aires - Montevideo,
17.6. - 13.7.1994
94 pages, Bremen, 1994.
- No. 59** **Bleil, U. and cruise participants**
Report and preliminary results of METEOR-Cruise M 29/2, Montevideo - Rio de Janeiro
15.7. - 8.8.1994. 153 pages, Bremen, 1994.
- No. 60** **Henrich, R. and cruise participants**
Report and preliminary results of METEOR-Cruise M 29/3, Rio de Janeiro - Las Palmas
11.8. - 5.9.1994. Bremen, 1994. (out of print)

- No. 61** **Sagemann, J.**
Saisonale Variationen von Porenwasserprofilen, Nährstoff-Flüssen und Reaktionen
in intertidalen Sedimenten des Weser-Ästuars. 110 pages, Bremen, 1994. (out of print)
- No. 62** **Giese, M. and G. Wefer**
Bericht über den 3. JGOFS-Workshop. 5./6. Dezember 1994 in Bremen.
84 pages, Bremen, 1995.
- No. 63** **Mann, U.**
Genese kretazischer Schwarzschiefer in Kolumbien: Globale vs. regionale/lokale Prozesse.
153 pages, Bremen, 1995. (out of print)
- No. 64** **Willems, H., Wan X., Yin J., Dongdui L., Liu G., S. Dürr, K.-U. Gräfe**
The Mesozoic development of the N-Indian passive margin and of the Xigaze Forearc Basin in
southern Tibet, China. – Excursion Guide to IGCP 362 Working-Group Meeting
"Integrated Stratigraphy". 113 pages, Bremen, 1995. (out of print)
- No. 65** **Hünken, U.**
Liefergebiets - Charakterisierung proterozoischer Goldseifen in Ghana anhand von
Fluideinschluß - Untersuchungen. 270 pages, Bremen, 1995.
- No. 66** **Nyandwi, N.**
The Nature of the Sediment Distribution Patterns in the Spiekeroog Backbarrier Area,
the East Frisian Islands. 162 pages, Bremen, 1995.
- No. 67** **Isenbeck-Schröter, M.**
Transportverhalten von Schwermetallkationen und Oxoanionen in wassergesättigten Sanden.
- Laborversuche in Säulen und ihre Modellierung -. 182 pages, Bremen, 1995.
- No. 68** **Hebbeln, D. and cruise participants**
Report and preliminary results of SONNE-Cruise SO 102, Valparaiso - Valparaiso, 95.
134 pages, Bremen, 1995.
- No. 69** **Willems, H. (Sprecher), U.Bathmann, U. Bleil, T. v. Dobeneck, K. Herterich, B.B. Jorgensen,
E.-M. Nöthig, M. Olesch, J. Pätzold, H.D. Schulz, V. Smetacek, V. Speiß, G. Wefer**
Bericht des Graduierten-Kollegs Stoff-Flüsse in marine Geosystemen.
Berichtszeitraum Januar 1993 - Dezember 1995. 45 & 468 pages, Bremen, 1995.
- No. 70** **Giese, M. and G. Wefer**
Bericht über den 4. JGOFS-Workshop. 20./21. November 1995 in Bremen. 60 pages, Bremen, 1996.
(out of print)
- No. 71** **Meggers, H.**
Pliozän-quartäre Karbonatsedimentation und Paläozeanographie des Nordatlantiks und
des Europäischen Nordmeeres - Hinweise aus planktischen Foraminiferengemeinschaften.
143 pages, Bremen, 1996. (out of print)
- No. 72** **Teske, A.**
Phylogenetische und ökologische Untersuchungen an Bakterien des oxidativen und reduktiven
marinen Schwefelkreislaufs mittels ribosomaler RNA. 220 pages, Bremen, 1996. (out of print)
- No. 73** **Andersen, N.**
Biogeochemische Charakterisierung von Sinkstoffen und Sedimenten aus ostatlantischen
Produktions-Systemen mit Hilfe von Biomarkern. 215 pages, Bremen, 1996.
- No. 74** **Treppke, U.**
Saisonaltät im Diatomeen- und Silikoflagellatenfluß im östlichen tropischen und subtropischen
Atlantik. 200 pages, Bremen, 1996.
- No. 75** **Schüring, J.**
Die Verwendung von Steinkohlebergematerialien im Deponiebau im Hinblick auf die
Pyritverwitterung und die Eignung als geochemische Barriere. 110 pages, Bremen, 1996.
- No. 76** **Pätzold, J. and cruise participants**
Report and preliminary results of VICTOR HENSEN cruise JOPS II, Leg 6,
Fortaleza - Recife, 10.3. - 26.3. 1995 and Leg 8, Vitoria - Vitoria, 10.4. - 23.4.1995.
87 pages, Bremen, 1996.
- No. 77** **Bleil, U. and cruise participants**
Report and preliminary results of METEOR-Cruise M 34/1, Cape Town - Walvis Bay, 3.-26.1.1996.
129 pages, Bremen, 1996.
- No. 78** **Schulz, H.D. and cruise participants**
Report and preliminary results of METEOR-Cruise M 34/2, Walvis Bay - Walvis Bay, 29.1.-18.2.96
133 pages, Bremen, 1996.
- No. 79** **Wefer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 34/3, Walvis Bay - Recife, 21.2.-17.3.1996.
168 pages, Bremen, 1996.

- No. 80** **Fischer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 34/4, Recife - Bridgetown, 19.3.-15.4.1996.
105 pages, Bremen, 1996.
- No. 81** **Kulbrok, F.**
Biostratigraphie, Fazies und Sequenzstratigraphie einer Karbonatrampe in den Schichten der Oberkreide und des Alttertiärs Nordost-Ägyptens (Eastern Desert, N'Golf von Suez, Sinai).
153 pages, Bremen, 1996.
- No. 82** **Kasten, S.**
Early Diagenetic Metal Enrichments in Marine Sediments as Documents of Nonsteady-State Depositional Conditions. Bremen, 1996.
- No. 83** **Holmes, M.E.**
Reconstruction of Surface Ocean Nitrate Utilization in the Southeast Atlantic Ocean Based on Stable Nitrogen Isotopes. 113 pages, Bremen, 1996.
- No. 84** **Rühlemann, C.**
Akkumulation von Carbonat und organischem Kohlenstoff im tropischen Atlantik:
Spätquartäre Produktivitäts-Variationen und ihre Steuerungsmechanismen. 139 pages, Bremen, 1996.
- No. 85** **Ratmeyer, V.**
Untersuchungen zum Eintrag und Transport lithogener und organischer partikulärer Substanz im östlichen subtropischen Nordatlantik. 154 pages, Bremen, 1996.
- No. 86** **Cepek, M.**
Zeitliche und räumliche Variationen von Coccolithophoriden-Gemeinschaften im subtropischen Ost-Atlantik: Untersuchungen an Plankton, Sinkstoffen und Sedimenten. 156 pages, Bremen, 1996.
- No. 87** **Otto, S.**
Die Bedeutung von gelöstem organischen Kohlenstoff (DOC) für den Kohlenstofffluß im Ozean. 150 pages, Bremen, 1996.
- No. 88** **Hensen, C.**
Frühdiagenetische Prozesse und Quantifizierung benthischer Stoff-Flüsse in Oberflächensedimenten des Südatlantiks. 132 pages, Bremen, 1996.
- No. 89** **Giese, M. and G. Wefer**
Bericht über den 5. JGOFS-Workshop. 27./28. November 1996 in Bremen. 73 pages, Bremen, 1997.
- No. 90** **Wefer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 37/1, Lisbon - Las Palmas, 4.-23.12.1996.
79 pages, Bremen, 1997.
- No. 91** **Isenbeck-Schröter, M., E. Bedbur, M. Kofod, B. König, T. Schramm & G. Mattheß**
Occurrence of Pesticide Residues in Water - Assessment of the Current Situation in Selected EU Countries. 65 pages, Bremen 1997.
- No. 92** **Kühn, M.**
Geochemische Folgereaktionen bei der hydrogeothermalen Energiegewinnung. 129 pages, Bremen 1997.
- No. 93** **Determin, S. & K. Herterich**
JGOFS-A6 "Daten und Modelle": Sammlung JGOFS-relevanten Modelle in Deutschland.
26 pages, Bremen, 1997.
- No. 94** **Fischer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 38/1, Las Palmas - Recife, 25.1.-1.3.1997,
with Appendix: Core Descriptions from METEOR Cruise M 37/1. Bremen, 1997.
- No. 95** **Bleil, U. and cruise participants**
Report and preliminary results of METEOR-Cruise M 38/2, Recife - Las Palmas, 4.3.-14.4.1997.
126 pages, Bremen, 1997.
- No. 96** **Neuer, S. and cruise participants**
Report and preliminary results of VICTOR HENSEN-Cruise 96/1. Bremen, 1997.
- No. 97** **Villinger, H. and cruise participants**
Fahrtbericht SO 111, 20.8. - 16.9.1996. 115 pages, Bremen, 1997.
- No. 98** **Lüning, S.**
Late Cretaceous - Early Tertiary sequence stratigraphy, paleoecology and geodynamics of Eastern Sinai, Egypt. 218 pages, Bremen, 1997.
- No. 99** **Haese, R.R.**
Beschreibung und Quantifizierung frühdiagenetischer Reaktionen des Eisens in Sedimenten des Südatlantiks. 118 pages, Bremen, 1997.

- No. 100 Lührte, R. von**
Verwertung von Bremer Baggergut als Material zur Oberflächenabdichtung von Deponien - Geochemisches Langzeitverhalten und Schwermetall-Mobilität (Cd, Cu, Ni, Pb, Zn). Bremen, 1997.
- No. 101 Ebert, M.**
Der Einfluß des Redoxmilieus auf die Mobilität von Chrom im durchströmten Aquifer.
135 pages, Bremen, 1997.
- No. 102 Krögel, F.**
Einfluß von Viskosität und Dichte des Seewassers auf Transport und Ablagerung von Wattsedimenten (Langeooger Rückseitenwatt, südliche Nordsee). 168 pages, Bremen, 1997.
- No. 103 Kerntopf, B.**
Dinoflagellate Distribution Patterns and Preservation in the Equatorial Atlantic and Offshore North-West Africa. 137 pages, Bremen, 1997.
- No. 104 Breitzke, M.**
Elastische Wellenausbreitung in marinen Sedimenten - Neue Entwicklungen der Ultraschall Sedimentphysik und Sedimentechographie. 298 pages, Bremen, 1997.
- No. 105 Marchant, M.**
Rezente und spätquartäre Sedimentation planktischer Foraminiferen im Peru-Chile Strom.
115 pages, Bremen, 1997.
- No. 106 Habicht, K.S.**
Sulfur isotope fractionation in marine sediments and bacterial cultures.
125 pages, Bremen, 1997.
- No. 107 Hamer, K., R.v. Lührte, G. Becker, T. Felis, S. Keffel, B. Strotmann, C. Waschkowitz, M. Kölling, M. Isenbeck-Schröter, H.D. Schulz**
Endbericht zum Forschungsvorhaben 060 des Landes Bremen: Baggergut der Hafengruppe Bremen-Stadt: Modelluntersuchungen zur Schwermetallmobilität und Möglichkeiten der Verwertung von Hafenschlick aus Bremischen Häfen. 98 pages, Bremen, 1997.
- No. 108 Greeff, O.W.**
Entwicklung und Erprobung eines benthischen Landersystems zur in situ-Bestimmung von Sulfatreduktionsraten mariner Sedimente. 121 pages, Bremen, 1997.
- No. 109 Pätzold, M. und G. Wefer**
Bericht über den 6. JGOFS-Workshop am 4./5.12.1997 in Bremen. Im Anhang:
Publikationen zum deutschen Beitrag zur Joint Global Ocean Flux Study (JGOFS), Stand 1/1998.
122 pages, Bremen, 1998.
- No. 110 Landenberger, H.**
CoTReM, ein Multi-Komponenten Transport- und Reaktions-Modell. 142 pages, Bremen, 1998.
- No. 111 Villinger, H. und Fahrteilnehmer**
Fahrtbericht SO 124, 4.10. - 16.10.199. 90 pages, Bremen, 1997.
- No. 112 Gietl, R.**
Biostratigraphie und Sedimentationsmuster einer nordostägyptischen Karbonatrampe unter Berücksichtigung der Alveolinien-Faunen. 142 pages, Bremen, 1998.
- No. 113 Ziebis, W.**
The Impact of the Thalassinidean Shrimp Callianassa truncata on the Geochemistry of permeable, coastal Sediments. 158 pages, Bremen 1998.
- No. 114 Schulz, H.D. and cruise participants**
Report and preliminary results of METEOR-Cruise M 41/1, Málaga - Libreville, 13.2.-15.3.1998.
Bremen, 1998.
- No. 115 Völker, D.J.**
Untersuchungen an strömungsbeeinflußten Sedimentationsmustern im Südozean. Interpretation sedimentechographischer Daten und numerische Modellierung. 152 pages, Bremen, 1998.
- No. 116 Schlünz, B.**
Riverine Organic Carbon Input into the Ocean in Relation to Late Quaternary Climate Change.
136 pages, Bremen, 1998.
- No. 117 Kuhnert, H.**
Aufzeichnug des Klimas vor Westaustralien in stabilen Isotopen in Korallenskeletten.
109 pages, Bremen, 1998.
- No. 118 Kirst, G.**
Rekonstruktion von Oberflächenwassertemperaturen im östlichen Südatlantik anhand von Alkenonen. 130 pages, Bremen, 1998.
- No. 119 Dürkoop, A.**
Der Brasil-Strom im Spätquartär: Rekonstruktion der oberflächennahen Hydrographie während der letzten 400 000 Jahre. 121 pages, Bremen, 1998.

- No. 120 Lamy, F.**
Spätquartäre Variationen des terrigenen Sedimenteintrags entlang des chilenischen Kontinentalhangs als Abbild von Klimavariabilität im Milanković- und Sub-Milanković-Zeitbereich.
141 pages, Bremen, 1998.
- No. 121 Neuer, S. and cruise participants**
Report and preliminary results of POSEIDON-Cruise Pos 237/2, Vigo – Las Palmas, 18.3.-31.3.1998. 39 pages, Bremen, 1998
- No. 122 Romero, O.E.**
Marine planktonic diatoms from the tropical and equatorial Atlantic: temporal flux patterns and the sediment record. 205 pages, Bremen, 1998.
- No. 123 Spiess, V. und Fahrtteilnehmer**
Report and preliminary results of RV SONNE Cruise 125, Cochin – Chittagong, 17.10.-17.11.1997. 128 pages, Bremen, 1998.
- No. 124 Arz, H.W.**
Dokumentation von kurzfristigen Klimaschwankungen des Spätquartärs in Sedimenten des westlichen äquatorialen Atlantiks. 96 pages, Bremen, 1998.
- No. 125 Wolff, T.**
Mixed layer characteristics in the equatorial Atlantic during the late Quaternary as deduced from planktonic foraminifera. 132 pages, Bremen, 1998.
- No. 126 Dittert, N.**
Late Quaternary Planktic Foraminifera Assemblages in the South Atlantic Ocean: Quantitative Determination and Preservational Aspects. 165 pages, Bremen, 1998.
- No. 127 Höll, C.**
Kalkige und organisch-wandige Dinoflagellaten-Zysten in Spätquartären Sedimenten des tropischen Atlantiks und ihre palökologische Auswertbarkeit. 121 pages, Bremen, 1998.
- No. 128 Hencke, J.**
Redoxreaktionen im Grundwasser: Etablierung und Verlagerung von Reaktionsfronten und ihre Bedeutung für die Spurenelement-Mobilität. 122 pages, Bremen 1998.
- No. 129 Pätzold, J. and cruise participants**
Report and preliminary results of METEOR-Cruise M 41/3, Víctoria, Brasil – Salvador de Bahia, Brasil, 18.4. - 15.5.1998. Bremen, 1999.
- No. 130 Fischer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 41/4, Salvador de Bahia, Brasil – Las Palmas, Spain, 18.5. – 13.6.1998. Bremen, 1999.
- No. 131 Schlünz, B. und G. Wefer**
Bericht über den 7. JGOFS-Workshop am 3. und 4.12.1998 in Bremen. Im Anhang: Publikationen zum deutschen Beitrag zur Joint Global Ocean Flux Study (JGOFS), Stand 1/ 1999. 100 pages, Bremen, 1999.
- No. 132 Wefer, G. and cruise participants**
Report and preliminary results of METEOR-Cruise M 42/4, Las Palmas - Las Palmas - Viena do Castelo; 26.09.1998 - 26.10.1998. 104 pages, Bremen, 1999.
- No. 133 Felis, T.**
Climate and ocean variability reconstructed from stable isotope records of modern subtropical corals (Northern Red Sea). 111 pages, Bremen, 1999.
- No. 134 Draschba , S.**
North Atlantic climate variability recorded in reef corals from Bermuda. 108 pages, Bremen, 1999.
- No. 135 Schmieder, F.**
Magnetic Cyclostratigraphy of South Atlantic Sediments. 82 pages, Bremen, 1999.
- No. 136 Rieß, W.**
In situ measurements of respiration and mineralisation processes – Interaction between fauna and geochemical fluxes at active interfaces. 68 pages, Bremen, 1999.
- No. 137 Devey, C.W. and cruise participants**
Report and shipboard results from METEOR-cruise M 41/2, Libreville – Vitoria, 18.3. – 15.4.98. 59 pages, Bremen, 1999.
- No. 138 Wenzhöfer, F.**
Biogeochemical processes at the sediment water interface and quantification of metabolically driven calcite dissolution in deep sea sediments. 103 pages, Bremen, 1999.
- No. 139 Klump, J.**
Biogenic barite as a proxy of paleoproductivity variations in the Southern Peru-Chile Current. 107 pages, Bremen, 1999.

- No. 140** **Huber, R.**
Carbonate sedimentation in the northern Northatlantic since the late pliocene. 103 pages, Bremen, 1999.
- No. 141** **Schulz, H.**
Nitrate-storing sulfur bacteria in sediments of coastal upwelling. 94 pages, Bremen, 1999.
- No. 142** **Mai, S.**
Die Sedimentverteilung im Wattenmeer: ein Simulationsmodell. 114 pages, Bremen, 1999.
- No. 143** **Neuer, S. and cruise participants**
Report and preliminary results of Poseidon Cruise 248, Las Palmas - Las Palmas, 15.2.-26.2.1999. 45 pages, Bremen, 1999.
- No. 144** **Weber, A.**
Schwefelkreislauf in marinen Sedimenten und Messung von in situ Sulfatreduktionsraten. 122 pages, Bremen, 1999.
- No. 145** **Hadeler, A.**
Sorptionsreaktionen im Grundwasser: Unterschiedliche Aspekte bei der Modellierung des Transportverhaltens von Zink. 122 pages, 1999.
- No. 146** **Dierßen, H.**
Zum Kreislauf ausgewählter Spurenmetalle im Südatlantik: Vertikaltransport und Wechselwirkung zwischen Partikeln und Lösung. 167 pages, Bremen, 1999.
- No. 147** **Zühlsdorff, L.**
High resolution multi-frequency seismic surveys at the Eastern Juan de Fuca Ridge Flank and the Cascadia Margin – Evidence for thermally and tectonically driven fluid upflow in marine sediments. 118 pages, Bremen 1999.
- No. 148** **Kinkel, H.**
Living and late Quaternary Coccolithophores in the equatorial Atlantic Ocean: response of distribution and productivity patterns to changing surface water circulation. 183 pages, Bremen, 2000.
- No. 149** **Pätzold, J. and cruise participants**
Report and preliminary results of METEOR Cruise M 44/3, Aqaba (Jordan) - Safaga (Egypt) – Dubá (Saudi Arabia) – Suez (Egypt) - Haifa (Israel), 12.3.-26.3.-2.4.-4.4.1999. 135 pages, Bremen, 2000.
- No. 150** **Schlünz, B. and G. Wefer**
Bericht über den 8. JGOFS-Workshop am 2. und 3.12.1999 in Bremen. Im Anhang: Publikationen zum deutschen Beitrag zur Joint Global Ocean Flux Study (JGOFS), Stand 1/ 2000. 95 pages, Bremen, 2000.
- No. 151** **Schnack, K.**
Biostratigraphie und fazielle Entwicklung in der Oberkreide und im Alttertiär im Bereich der Kharga Schwelle, Westliche Wüste, SW-Ägypten. 142 pages, Bremen, 2000.
- No. 152** **Karwath, B.**
Ecological studies on living and fossil calcareous dinoflagellates of the equatorial and tropical Atlantic Ocean. 175 pages, Bremen, 2000.
- No. 153** **Moustafa, Y.**
Paleoclimatic reconstructions of the Northern Red Sea during the Holocene inferred from stable isotope records of modern and fossil corals and molluscs. 102 pages, Bremen, 2000.
- No. 154** **Villinger, H. and cruise participants**
Report and preliminary results of SONNE-cruise 145-1 Balboa – Talcahuana, 21.12.1999 – 28.01.2000. 147 pages, Bremen, 2000.
- No. 155** **Rusch, A.**
Dynamik der Feinfraktion im Oberflächenhorizont permeabler Schelfsedimente. 102 pages, Bremen, 2000.
- No. 156** **Moos, C.**
Reconstruction of upwelling intensity and paleo-nutrient gradients in the northwest Arabian Sea derived from stable carbon and oxygen isotopes of planktic foraminifera. 103 pages, Bremen, 2000.
- No. 157** **Xu, W.**
Mass physical sediment properties and trends in a Wadden Sea tidal basin. 127 pages, Bremen, 2000.
- No. 158** **Meinecke, G. and cruise participants**
Report and preliminary results of METEOR Cruise M 45/1, Malaga (Spain) - Lissabon (Portugal), 19.05. - 08.06.1999. 39 pages, Bremen, 2000.
- No. 159** **Vink, A.**
Reconstruction of recent and late Quaternary surface water masses of the western subtropical Atlantic Ocean based on calcareous and organic-walled dinoflagellate cysts. 160 pages, Bremen, 2000.
- No. 160** **Willems, H. (Sprecher), U. Bleil, R. Henrich, K. Herterich, B.B. Jørgensen, H.-J. Kuß, M. Olesch, H.D. Schulz, V. Spieß, G. Wefer**
Abschlußbericht des Graduierten-Kollegs Stoff-Flüsse in marine Geosystemen.
Zusammenfassung und Berichtszeitraum Januar 1996 - Dezember 2000. 340 pages, Bremen, 2000.

- No. 161** **Sprengel, C.**
Untersuchungen zur Sedimentation und Ökologie von Coccolithophoriden im Bereich der Kanarischen Inseln: Saisonale Flussmuster und Karbonatelexport. 165 pages, Bremen, 2000.
- No. 162** **Donner, B. and G. Wefer**
Bericht über den JGOFS-Workshop am 18.-21.9.2000 in Bremen: Biogeochemical Cycles: German Contributions to the International Joint Global Ocean Flux Study. 87 pages, Bremen, 2000.
- No. 163** **Neuer, S. and cruise participants**
Report and preliminary results of Meteor Cruise M 45/5, Bremen – Las Palmas, October 1 – November 3, 1999. 93 pages, Bremen, 2000.
- No. 164** **Devey, C. and cruise participants**
Report and preliminary results of Sonne Cruise SO 145/2, Talcahuano (Chile) - Arica (Chile), February 4 – February 29, 2000. 63 pages, Bremen, 2000.
- No. 165** **Freudenthal, T.**
Reconstruction of productivity gradients in the Canary Islands region off Morocco by means of sinking particles and sediments. 147 pages, Bremen, 2000.
- No. 166** **Adler, M.**
Modeling of one-dimensional transport in porous media with respect to simultaneous geochemical reactions in CoTReM. 147 pages, Bremen, 2000.
- No. 167** **Santamarina Cuneo, P.**
Fluxes of suspended particulate matter through a tidal inlet of the East Frisian Wadden Sea (southern North Sea). 91 pages, Bremen, 2000.
- No. 168** **Benthien, A.**
Effects of CO₂ and nutrient concentration on the stable carbon isotope composition of C37:2 alkenones in sediments of the South Atlantic Ocean. 104 pages, Bremen, 2001.
- No. 169** **Lavik, G.**
Nitrogen isotopes of sinking matter and sediments in the South Atlantic. 140 pages, Bremen, 2001.
- No. 170** **Budziak, D.**
Late Quaternary monsoonal climate and related variations in paleoproductivity and alkenone-derived sea-surface temperatures in the western Arabian Sea. 114 pages, Bremen, 2001.
- No. 171** **Gerhardt, S.**
Late Quaternary water mass variability derived from the pteropod preservation state in sediments of the western South Atlantic Ocean and the Caribbean Sea. 109 pages, Bremen, 2001.
- No. 172** **Bleil, U. and cruise participants**
Report and preliminary results of Meteor Cruise M 46/3, Montevideo (Uruguay) – Mar del Plata (Argentina), January 4 – February 7, 2000. Bremen, 2001.
- No. 173** **Wefer, G. and cruise participants**
Report and preliminary results of Meteor Cruise M 46/4, Mar del Plata (Argentina) – Salvador da Bahia (Brazil), February 10 – March 13, 2000. With partial results of METEOR cruise M 46/2. 136 pages, Bremen, 2001.
- No. 174** **Schulz, H.D. and cruise participants**
Report and preliminary results of Meteor Cruise M 46/2, Recife (Brazil) – Montevideo (Uruguay), December 2 – December 29, 1999. 107 pages, Bremen, 2001.
- No. 175** **Schmidt, A.**
Magnetic mineral fluxes in the Quaternary South Atlantic: Implications for the paleoenvironment. 97 pages, Bremen, 2001.
- No. 176** **Bruhns, P.**
Crystal chemical characterization of heavy metal incorporation in brick burning processes. 93 pages, Bremen, 2001.
- No. 177** **Karius, V.**
Baggergut der Hafengruppe Bremen-Stadt in der Ziegelherstellung. 131 pages, Bremen, 2001.
- No. 178** **Adegbie, A. T.**
Reconstruction of paleoenvironmental conditions in Equatorial Atlantic and the Gulf of Guinea Basins for the last 245,000 years. 113 pages, Bremen, 2001.
- No. 179** **Spieß, V. and cruise participants**
Report and preliminary results of R/V Sonne Cruise SO 149, Victoria - Victoria, 16.8. - 16.9.2000. 100 pages, Bremen, 2001.
- No. 180** **Kim, J.-H.**
Reconstruction of past sea-surface temperatures in the eastern South Atlantic and the eastern South Pacific across Termination I based on the Alkenone Method. 114 pages, Bremen, 2001.

- No. 181 von Lom-Keil, H.**
Sedimentary waves on the Namibian continental margin and in the Argentine Basin – Bottom flow reconstructions based on high resolution echosounder data. 126 pages, Bremen, 2001.
- No. 182 Hebbeln, D. and cruise participants**
PUCK: Report and preliminary results of R/V Sonne Cruise SO 156, Valparaiso (Chile) - Talcahuano (Chile), March 29 - May 14, 2001. 195 pages, Bremen, 2001.
- No. 183 Wendler, J.**
Reconstruction of astronomically-forced cyclic and abrupt paleoecological changes in the Upper Cretaceous Boreal Realm based on calcareous dinoflagellate cysts. 149 pages, Bremen, 2001.
- No. 184 Volbers, A.**
Planktic foraminifera as paleoceanographic indicators: production, preservation, and reconstruction of upwelling intensity. Implications from late Quaternary South Atlantic sediments. 122 pages, Bremen, 2001.
- No. 185 Bleil, U. and cruise participants**
Report and preliminary results of R/V METEOR Cruise M 49/3, Montevideo (Uruguay) - Salvador (Brasil), March 9 - April 1, 2001. 99 pages, Bremen, 2001.
- No. 186 Scheibner, C.**
Architecture of a carbonate platform-to-basin transition on a structural high (Campanian-early Eocene, Eastern Desert, Egypt) – classical and modelling approaches combined. 173 pages, Bremen, 2001.
- No. 187 Schneider, S.**
Quartäre Schwankungen in Strömungsintensität und Produktivität als Abbild der Wassermassen-Variabilität im äquatorialen Atlantik (ODP Sites 959 und 663): Ergebnisse aus Siltkorn-Analysen. 134 pages, Bremen, 2001.
- No. 188 Uliana, E.**
Late Quaternary biogenic opal sedimentation in diatom assemblages in Kongo Fan sediments. 96 pages, Bremen, 2002.
- No. 189 Esper, O.**
Reconstruction of Recent and Late Quaternary oceanographic conditions in the eastern South Atlantic Ocean based on calcareous- and organic-walled dinoflagellate cysts. 130 pages, Bremen, 2001.
- No. 190 Wendler, I.**
Production and preservation of calcareous dinoflagellate cysts in the modern Arabian Sea. 117 pages, Bremen, 2002.
- No. 191 Bauer, J.**
Late Cenomanian – Santonian carbonate platform evolution of Sinai (Egypt): stratigraphy, facies, and sequence architecture. 178 pages, Bremen, 2002.
- No. 192 Hildebrand-Habel, T.**
Die Entwicklung kalkiger Dinoflagellaten im Südatlantik seit der höheren Oberkreide. 152 pages, Bremen, 2002.
- No. 193 Hecht, H.**
Sauerstoff-Optopoden zur Quantifizierung von Pyritverwitterungsprozessen im Labor- und Langzeit-in-situ-Einsatz. Entwicklung - Anwendung – Modellierung. 130 pages, Bremen, 2002.
- No. 194 Fischer, G. and cruise participants**
Report and Preliminary Results of RV METEOR-Cruise M49/4, Salvador da Bahia – Halifax, 4.4.-5.5.2001. 84 pages, Bremen, 2002.
- No. 195 Gröger, M.**
Deep-water circulation in the western equatorial Atlantic: inferences from carbonate preservation studies and silt grain-size analysis. 95 pages, Bremen, 2002.
- No. 196 Meinecke, G. and cruise participants**
Report of RV POSEIDON Cruise POS 271, Las Palmas - Las Palmas, 19.3.-29.3.2001. 19 pages, Bremen, 2002.
- No. 197 Meggers, H. and cruise participants**
Report of RV POSEIDON Cruise POS 272, Las Palmas - Las Palmas, 1.4.-14.4.2001. 19 pages, Bremen, 2002.
- No. 198 Gräfe, K.-U.**
Stratigraphische Korrelation und Steuerungsfaktoren Sedimentärer Zyklen in ausgewählten Borealen und Tethyalen Becken des Cenoman/Turon (Oberkreide) Europas und Nordwestafrikas. 197 pages, Bremen, 2002.
- No. 199 Jahn, B.**
Mid to Late Pleistocene Variations of Marine Productivity in and Terrigenous Input to the Southeast Atlantic. 97 pages, Bremen, 2002.
- No. 200 Al-Rousan, S.**
Ocean and climate history recorded in stable isotopes of coral and foraminifers from the northern Gulf of Aqaba. 116 pages, Bremen, 2002.

- No. 201** **Azouzi, B.**
Regionalisierung hydraulischer und hydrogeochemischer Daten mit geostatistischen Methoden.
108 pages, Bremen, 2002.
- No. 202** **Spieß, V. and cruise participants**
Report and preliminary results of METEOR Cruise M 47/3, Libreville (Gabun) - Walvis Bay
(Namibia), 01.06 - 03.07.2000. 70 pages, Bremen 2002.
- No. 203** **Spieß, V. and cruise participants**
Report and preliminary results of METEOR Cruise M 49/2, Montevideo (Uruguay) - Montevideo,
13.02 - 07.03.2001. 84 pages, Bremen 2002.
- No. 204** **Mollenhauer, G.**
Organic carbon accumulation in the South Atlantic Ocean: Sedimentary processes and
glacial/interglacial Budgets. 139 pages, Bremen 2002.
- No. 205** **Spieß, V. and cruise participants**
Report and preliminary results of METEOR Cruise M49/1, Cape Town (South Africa) - Montevideo
(Uruguay), 04.01.2001 - 10.02.2001. 57 pages, Bremen, 2003.
- No. 206** **Meier, K.J.S.**
Calcareous dinoflagellates from the Mediterranean Sea: taxonomy, ecology and palaeoenvironmental
application. 126 pages, Bremen, 2003.
- No. 207** **Rakic, S.**
Untersuchungen zur Polymorphie und Kristallchemie von Silikaten der Zusammensetzung $\text{Me}_2\text{Si}_2\text{O}_5$
($\text{Me}:\text{Na}, \text{K}$). 139 pages, Bremen, 2003.
- No. 208** **Pfeifer, K.**
Auswirkungen frühdiagenetischer Prozesse auf Calcit- und Barytgehalte in marinen Oberflächen-
sedimenten. 110 pages, Bremen, 2003.
- No. 209** **Heuer, V.**
Spurenelemente in Sedimenten des Südatlantik. Primärer Eintrag und frühdiagenetische Überprägung.
136 pages, Bremen, 2003.
- No. 210** **Streng, M.**
Phylogenetic Aspects and Taxonomy of Calcareous Dinoflagellates. 157 pages, Bremen 2003.
- No. 211** **Boeckel, B.**
Present and past coccolith assemblages in the South Atlantic: implications for species ecology,
carbonate contribution and palaeoceanographic applicability. 157 pages, Bremen, 2003.
- No. 212** **Precht, E.**
Advectional interfacial exchange in permeable sediments driven by surface gravity waves and its
ecological consequences. 131 pages, Bremen, 2003.
- No. 213** **Frenz, M.**
Grain-size composition of Quaternary South Atlantic sediments and its paleoceanographic significance.
123 pages, Bremen, 2003.
- No. 214** **Meggers, H. and cruise participants**
Report and preliminary results of METEOR Cruise M 53/1, Limassol - Las Palmas – Mindelo,
30.03.2002 - 03.05.2002. 81 pages, Bremen, 2003.
- No. 215** **Schulz, H.D. and cruise participants**
Report and preliminary results of METEOR Cruise M 58/1, Dakar – Las Palmas, 15.04..2003 –
12.05.2003. Bremen, 2003.
- No. 216** **Schneider, R. and cruise participants**
Report and preliminary results of METEOR Cruise M 57/1, Cape Town – Walvis Bay, 20.01. –
08.02.2003. 123 pages, Bremen, 2003.
- No. 217** **Kallmeyer, J.**
Sulfate reduction in the deep Biosphere. 157 pages, Bremen, 2003.
- No. 218** **Røy, H.**
Dynamic Structure and Function of the Diffusive Boundary Layer at the Seafloor. 149 pages,
Bremen, 2003.
- No. 219** **Pätzold, J., C. Hübscher and cruise participants**
Report and preliminary results of METEOR Cruise M 52/2&3, Istanbul – Limassol – Limassol,
04.02. – 27.03.2002. Bremen, 2003.
- No. 220** **Zabel, M. and cruise participants**
Report and preliminary results of METEOR Cruise M 57/2, Walvis Bay – Walvis Bay, 11.02. –
12.03.2003. 136 pages, Bremen 2003.
- No. 221** **Salem, M.**
Geophysical investigations of submarine prolongations of alluvial fans on the western side of the Gulf
of Aqaba-Red Sea. 100 pages, Bremen, 2003.

- No. 222** **Tilch, E.**
Oszillation von Wattflächen und deren fossiles Erhaltungspotential (Spiekerooger Rückseitenwatt, südliche Nordsee). 137 pages, Bremen, 2003.
- No. 223** **Frisch, U. and F. Kockel**
Der Bremen-Knoten im Strukturnetz Nordwest-Deutschlands. Stratigraphie, Paläogeographie, Strukturgeologie. 379 pages, Bremen, 2004.
- No. 224** **Kolonic, S.**
Mechanisms and biogeochemical implications of Cenomanian/Turonian black shale formation in North Africa: An integrated geochemical, millennial-scale study from the Tarfaya-LaAyoune Basin in SW Morocco. 174 pages, Bremen, 2004. Report online available only.
- No. 225** **Panteleit, B.**
Geochemische Prozesse in der Salz- Süßwasser Übergangszone. 106 pages, Bremen, 2004.
- No. 226** **Seiter, K.**
Regionalisierung und Quantifizierung benthischer Mineralisationsprozesse. 135 pages, Bremen, 2004.
- No. 227** **Bleil, U. and cruise participants**
Report and preliminary results of METEOR Cruise M 58/2, Las Palmas – Las Palmas (Canary Islands, Spain), 15.05. – 08.06.2003. 123 pages, Bremen, 2004.
- No. 228** **Kopf, A. and cruise participants**
Report and preliminary results of SONNE Cruise SO175, Miami - Bremerhaven, 12.11 - 30.12.2003. 218 pages, Bremen, 2004.
- No. 229** **Fabian, M.**
Near Surface Tilt and Pore Pressure Changes Induced by Pumping in Multi-Layered Poroelastic Half-Spaces. 121 pages, Bremen, 2004.
- No. 230** **Segl, M. , and cruise participants**
Report and preliminary results of POSEIDON cruise 304 Galway – Lisbon, 5. – 22. Oct. 2004. 27 pages, Bremen 2004
- No. 231** **Meinecke, G. and cruise participants**
Report and preliminary results of POSEIDON Cruise 296, Las Palmas – Las Palmas, 04.04 – 14.04.2003. 42 pages, Bremen 2005.
- No. 232** **Meinecke, G. and cruise participants**
Report and preliminary results of POSEIDON Cruise 310, Las Palmas – Las Palmas, 12.04 – 26.04.2004. 49 pages, Bremen 2005.
- No. 233** **Meinecke, G. and cruise participants**
Report and preliminary results of METEOR Cruise 58/3, Las Palmas - Ponta Delgada, 11.06 - 24.06.2003. 50 pages, Bremen 2005.
- No. 234** **Feseker, T.**
Numerical Studies on Groundwater Flow in Coastal Aquifers. 219 pages. Bremen 2004.
- No. 235** **Sahling, H. and cruise participants**
Report and preliminary results of R/V POSEIDON Cruise P317/4, Istanbul-Istanbul , 16 October - 4 November 2004. 92 pages, Bremen 2004.
- No. 236** **Meinecke, G. und Fahrteilnehmer**
Report and preliminary results of POSEIDON Cruise 305, Las Palmas (Spain) - Lisbon (Portugal), October 28th – November 6th, 2004. 43 pages, Bremen 2005.
- No. 237** **Ruhland, G. and cruise participants**
Report and preliminary results of POSEIDON Cruise 319, Las Palmas (Spain) - Las Palmas (Spain), December 6th – December 17th, 2004. 50 pages, Bremen 2005.
- No. 238** **Chang, T.S.**
Dynamics of fine-grained sediments and stratigraphic evolution of a back-barrier tidal basin of the German Wadden Sea (southern North Sea). 102 pages, Bremen 2005.
- No. 239** **Lager, T.**
Predicting the source strength of recycling materials within the scope of a seepage water prognosis by means of standardized laboratory methods. 141 pages, Bremen 2005.
- No. 240** **Meinecke, G.**
DOLAN - Operationelle Datenübertragung im Ozean und Laterales Akustisches Netzwerk in der Tiefsee. Abschlußbericht. 42 pages, Bremen 2005.
- No. 241** **Guasti, E.**
Early Paleogene environmental turnover in the southern Tethys as recorded by foraminiferal and organic-walled dinoflagellate cysts assemblages. 203 pages, Bremen 2005.
- No. 242** **Riedinger, N.**
Preservation and diagenetic overprint of geochemical and geophysical signals in ocean margin sediments related to depositional dynamics. 91 pages, Bremen 2005.

- No. 243 Ruhland, G. and cruise participants**
Report and preliminary results of POSEIDON cruise 320, Las Palmas (Spain) - Las Palmas (Spain), March 08th - March 18th, 2005. 57 pages, Bremen 2005.
- No. 244 Inthorn, M.**
Lateral particle transport in nepheloid layers – a key factor for organic matter distribution and quality in the Benguela high-productivity area. 127 pages, Bremen, 2006.
- No. 245 Aspetsberger, F.**
Benthic carbon turnover in continental slope and deep sea sediments: importance of organic matter quality at different time scales. 136 pages, Bremen, 2006.
- No. 246 Hebbeln, D. and cruise participants**
Report and preliminary results of RV SONNE Cruise SO-184, PABESIA, Durban (South Africa) – Cilacap (Indonesia) – Darwin (Australia), July 08th - September 13th, 2005. 142 pages, Bremen 2006.
- No. 247 Ratmeyer, V. and cruise participants**
Report and preliminary results of RV METEOR Cruise M61/3. Development of Carbonate Mounds on the Celtic Continental Margin, Northeast Atlantic. Cork (Ireland) – Ponta Delgada (Portugal), 04.06. – 21.06.2004. 64 pages, Bremen 2006.
- No. 248 Wien, K.**
Element Stratigraphy and Age Models for Pelagites and Gravity Mass Flow Deposits based on Shipboard XRF Analysis. 100 pages, Bremen 2006.
- No. 249 Krastel, S. and cruise participants**
Report and preliminary results of RV METEOR Cruise M65/2, Dakar - Las Palmas, 04.07. – 26.07.2005. 185 pages, Bremen 2006.
- No. 250 Heil, G.M.N.**
Abrupt Climate Shifts in the Western Tropical to Subtropical Atlantic Region during the Last Glacial. 121 pages, Bremen 2006.
- No. 251 Ruhland, G. and cruise participants**
Report and preliminary results of POSEIDON Cruise 330, Las Palmas – Las Palmas, November 21th – December 03rd, 2005. 48 pages, Bremen 2006.
- No. 252 Mulitza , S. and cruise participants**
Report and preliminary results of METEOR Cruise M65/1, Dakar – Dakar, 11.06.- 1.07.2005. 149 pages, Bremen 2006.
- No. 253 Kopf, A. and cruise participants**
Report and preliminary results of POSEIDON Cruise P336, Heraklion - Heraklion, 28.04. – 17.05.2006. 127 pages, Bremen, 2006.
- No. 254 Wefer, G. and cruise participants**
Report and preliminary results of R/V METEOR Cruise M65/3, Las Palmas - Las Palmas (Spain), July 31st - August 10th, 2005. 24 pages, Bremen 2006.
- No. 255 Hanebuth, T.J.J. and cruise participants**
Report and first results of the POSEIDON Cruise P342 GALIOMAR, Vigo – Lisboa (Portugal), August 19th – September 06th, 2006. Distribution Pattern, Residence Times and Export of Sediments on the Pleistocene/Holocene Galician Shelf (NW Iberian Peninsula). 203 pages, Bremen, 2007.
- No. 256 Ahke, A.**
Composition of molecular organic matter pools, pigments and proteins, in Benguela upwelling and Arctic Sediments. 192 pages, Bremen 2007.
- No. 257 Becker, V.**
Seeper - Ein Modell für die Praxis der Sickerwasserprognose. 170 pages, Bremen 2007.
- No. 258 Ruhland, G. and cruise participants**
Report and preliminary results of Poseidon cruise 333, Las Palmas (Spain) – Las Palmas (Spain), March 1st – March 10th, 2006. 32 pages, Bremen 2007.
- No. 259 Fischer, G., G. Ruhland and cruise participants**
Report and preliminary results of Poseidon cruise 344, leg 1 and leg 2, Las Palmas (Spain) – Las Palmas (Spain), Oct. 20th – Nov 2nd & Nov. 4th – Nov 13th, 2006. 46 pages, Bremen 2007.
- No. 260 Westphal, H. and cruise participants**
Report and preliminary results of Poseidon cruise 346, MACUMA. Las Palmas (Spain) – Las Palmas (Spain), 28.12.2006 – 15.1.2007. 49 pages, Bremen 2007.
- No. 261 Bohrmann, G., T. Pape, and cruise participants**
Report and preliminary results of R/V METEOR Cruise M72/3, Istanbul – Trabzon – Istanbul, March 17th – April 23rd, 2007. Marine gas hydrates of the Eastern Black Sea. 130 pages, Bremen 2007.
- No. 262 Bohrmann, G., and cruise participants**
Report and preliminary results of R/V METEOR Cruise M70/3, Iraklion – Iraklion, 21 November – 8 December 2006. Cold Seeps of the Anaximander Mountains / Eastern Mediterranean. 75 pages, Bremen 2008.

- No. 263 Bohrmann, G., Spiess, V., and cruise participants**
Report and preliminary results of R/V Meteor Cruise M67/2a and 2b, Balboa -- Tampico -- Bridgetown, 15 March -- 24 April, 2006. Fluid seepage in the Gulf of Mexico. Bremen 2008.
- No. 264 Kopf, A., and cruise participants**
Report and preliminary results of Meteor Cruise M73/1: LIMA-LAMO (Ligurian Margin Landslide Measurements & Observatory), Cadiz, 22.07.2007 – Genoa, 11.08.2007. 170 pages, Bremen 2008.
- No. 265 Hebbeln, D., and cruise participants**
Report and preliminary results of RV Pelagia Cruise 64PE284. Cold-water Corals in the Gulf of Cádiz and on Coral Patch Seamount (NE Atlantic). Portimão - Portimão, 18.02. - 09.03.2008. 90 pages, Bremen 2008.
- No. 266 Bohrmann, G. and cruise participants**
Report and preliminary results of R/V Meteor Cruise M74/3, Fujairah – Male, 30 October - 28 November, 2007. Cold Seeps of the Makran subduction zone (Continental margin of Pakistan). 161 pages, Bremen 2008.
- No. 267 Sachs, O.**
Benthic organic carbon fluxes in the Southern Ocean: Regional differences and links to surface primary production and carbon export. 143 pages, Bremen, 2008.
- No. 268 Zonneveld, K. and cruise participants**
Report and preliminary results of R/V POSEIDON Cruise P339, Piräus - Messina, 16 June - 2 July 2006. CAPPUCINO - Calabrian and Adriatic palaeoproductivity and climatic variability in the last two millenia. 61 pages, Bremen, 2008.
- No. 269 Ruhland, G. and cruise participants**
Report and preliminary results of R/V POSEIDON Cruise P360, Las Palmas (Spain) - Las Palmas (Spain), Oct. 29th - Nov. 6th, 2007. 27 pages, Bremen, 2008.
- No. 270 Ruhland, G. , G. Fischer and cruise participants**
Report and preliminary results of R/V POSEIDON Cruise 365 (Leg 1+2). Leg 1: Las Palmas - Las Palmas, 13.4. - 16.4.2008. Leg 2: Las Palmas - Las Palmas, 18.4. - 29.4.2008. 40 pages, Bremen, 2009.