

Supporting Information

Molecular Networking-Guided Isolation of New Etzionin Type Diketopiperazine Hydroxamates from the Persian Gulf Sponge *Cliona celata*

Reza Mohsenian Kouchaksaraee¹, Fengjie Li¹, Melika Nazemi,² Mahdi Moridi Farimani³ and Deniz Tasdemir^{1,4,*}

¹ GEOMAR Centre for Marine Biotechnology (GEOMAR-Biotech), Research Unit Marine Natural Products Chemistry, GEOMAR Helmholtz Centre for Ocean Research Kiel, Am Kiel-Kanal 44, 24106 Kiel, Germany; rmohsenian@geomar.de; fli@geomar.de

² Persian Gulf and Oman Sea Ecological Center, Iranian Fisheries Science Research Institute, Agricultural Research, Education and Extension Organization (AREEO), 7916793165 Bandar Abbas, Iran; melikanazemi@pgoseri.ac.ir

³ Department of Phytochemistry, Medicinal Plants and Drugs Research Institute, Shahid Beheshti University, G. C., Evin, 1983969411 Tehran, Iran; m_moridi@sbu.ac.ir

⁴ Faculty of Mathematics and Natural Sciences, Kiel University, Christian-Albrechts-Platz 4, 24118 Kiel, Germany

* Correspondence: dtasdemir@geomar.de; Tel.: +49-431-6004430

Content	Page No.
Table S1. Putative identification of the compounds (clusters A and B) in the global molecular network of the crude extract of <i>Cliona celata</i>	3
Figure S1. Global molecular network of the crude extract of <i>Cliona celata</i>	4
Figure S2. UPLC-HRMS chromatogram of the crude extract from <i>Cliona celata</i>	5
Figure S3. ^1H NMR spectrum of the fraction F3 from <i>Cliona celata</i> (600 MHz, CD ₃ OD)	5
Figure S4. ^1H NMR spectrum of compound 1 (TFA salt, 600 MHz, CD ₃ OD)	6
Figure S5. ^{13}C NMR spectrum of compound 1 (TFA salt, 150 MHz, CD ₃ OD)	6
Figure S6. HSQC spectrum of compound 1 (TFA salt, 600 MHz, CD ₃ OD, non-uniform sampling)	7
Figure S7. HSQC spectrum of compound 1 (TFA salt, 600 MHz, CD ₃ OD, traditional planes)	7
Figure S8. HMBC spectrum of compound 1 (TFA salt, 600 MHz, CD ₃ OD, non-uniform sampling)	8
Figure S9. HMBC spectrum of compound 1 (TFA salt, 600 MHz, CD ₃ OD, traditional planes)	8
Figure S10. COSY spectrum of compound 1 (TFA salt, 600 MHz, CD ₃ OD)	9
Figure S11. NOESY spectrum of compound 1 (TFA salt, 600 MHz, CD ₃ OD)	9
Figure S12. HR-ESIMS spectrum of compound 1	10
Figure S13. ^1H NMR spectrum of compound 2 (TFA salt, 600 MHz, CD ₃ OD)	11
Figure S14. ^{13}C NMR spectrum of compound 2 (TFA salt, 150 MHz, CD ₃ OD)	11
Figure S15. HSQC spectrum of compound 2 (TFA salt, 600 MHz, CD ₃ OD)	12
Figure S16. HMBC spectrum of compound 2 (TFA salt, 600 MHz, CD ₃ OD)	12
Figure S17. COSY spectrum of compound 2 (TFA salt, 600 MHz, CD ₃ OD)	13
Figure S18. NOESY spectrum of compound 2 (TFA salt, 600 MHz, CD ₃ OD)	13
Figure S19. HR-ESIMS spectrum of compound 2	14
Figure S20. ^1H NMR spectrum of compound 3 (TFA salt, 600 MHz, CD ₃ OD)	15
Figure S21. ^{13}C NMR spectrum of compound 3 (TFA salt, 150 MHz, CD ₃ OD)	15
Figure S22. HSQC spectrum of compound 3 (TFA salt, 600 MHz, CD ₃ OD)	16
Figure S23. HMBC spectrum of compound 3 (TFA salt, 600 MHz, CD ₃ OD)	16
Figure S24. COSY spectrum of compound 3 (TFA salt, 600 MHz, CD ₃ OD)	17
Figure S25. NOESY spectrum of compound 3 (TFA salt, 600 MHz, CD ₃ OD)	17
Figure S26. HR-ESIMS spectrum of compound 3	18

Table S1. Putative identification of the compounds (cluster A and B) in the global molecular network of the crude extract of *Cliona celata*.

No.	t _R (min)	Parent mass m/z	Cluster	Molecular formula of the m/z [M + H] ⁺	Δ ppm	Key MS ² fragments	LoA*
1	5.43	459.332	A	C ₂₆ H ₄₃ N ₄ O ₃	0.0	205.097; 238.217; 343.238; 385.249; 441.297	3
2	4.97	473.293	A	C ₂₆ H ₄₁ N ₄ O ₄	0.4	221.093; 236.202; 357.213; 399.229; 456.887	3
3	5.42	475.315	A	C ₂₆ H ₄₃ N ₄ O ₄	0.1	221.093; 252.233; 359.233; 401.244; 458.302	5
4	5.44	489.332	A	C ₂₇ H ₄₅ N ₄ O ₄	0.2	221.093; 259.233; 359.233; 401.224; 471.333	5
5	6.03	491.305	A	C ₂₆ H ₄₃ N ₄ O ₅	0.4	214.181; 238.218; 375.228; 417.240; 474.229	3
6	5.54	503.359	A	C ₂₈ H ₄₇ N ₄ O ₄	0.4	221.093; 266.248; 359.233; 401.243; 485.347	3
7	5.86	503.359	A	C ₂₈ H ₄₇ N ₄ O ₄	1.2	221.093; 266.249; 373.249; 415.259; 485.347	3
8	5.35	505.338	A	C ₂₇ H ₄₅ N ₄ O ₅	-0.1	221.090; 238.216; 359.233; 401.240; 487.332	3
9	6.04	505.339	A	C ₂₇ H ₄₅ N ₄ O ₅	0.6	214.180; 252.232; 375.229; 417.246; 487.237	3
10	5.28	637.373	A	C ₃₂ H ₅₃ N ₄ O ₉	0.8	221.093; 238.277; 359.232; 401.245; 553.339	3
11	7.04	419.231	B	C ₂₃ H ₃₅ N ₂ O ₅	0.6	181.159; 196.170; 221.093; 359.233; 401.244	5
12	7.15	403.235	B	C ₂₃ H ₃₃ N ₂ O ₄	0.6	181.159; 205.096; 315.242; 343.237; 385.247	3

*: LoA (Level of Assignment); 1: Accurate mass matched to database-Tentative assignment, 2: Accurate mass matched to database and tandem MS spectrum matched to *in silico* fragmentation pattern, 3: Tandem MS spectrum matched to database or literature, 4: RT matched to standard compound, 5: MS/MS spectrum matched to standard compound.

Manual dereplication was performed considering the parent mass, biological source, retention time, elemental composition analysis, and predicated fragmentation patterns.

The MS² fragmentation pattern of a molecule was predicated on the Competitive Fragmentation Modeling for Metabolite Identification (CFM-ID) platform (<http://cfmid.wishartlab.com>) and compared with our experimental data.

Figure S1. Global molecular network of the crude extract of *Cliona celata*.




Figure S2. UPLC-HRMS chromatogram of the crude extract from *Cliona celata*.




Figure S3. ^1H NMR spectrum of the fraction F3 from *Cliona celata* (600 MHz, CD_3OD)




Figure S4. ^1H NMR spectrum of compound **1** (TFA salt, 600 MHz, CD_3OD).




Figure S5. ^{13}C NMR spectrum of compound **1** (TFA salt, 150 MHz, CD_3OD)




Figure S6. HSQC spectrum of compound 1 (TFA salt, 600 MHz, CD₃OD, non-uniform sampling).




Figure S7. HSQC spectrum of compound 1 (TFA salt, 600 MHz, CD₃OD, traditional planes).




Figure S8. HMBC spectrum of compound **1** (TFA salt, 600 MHz, CD₃OD, non-uniform sampling).




Figure S9. HMBC spectrum of compound **1** (TFA salt, 600 MHz, CD₃OD, traditional planes).




Figure S10. COSY spectrum of compound **1** (TFA salt, 600 MHz, CD₃OD).




Figure S11. NOESY spectrum of compound **1** (TFA salt, 600 MHz, CD₃OD).




Figure S12. HR-ESIMS spectrum of compound 1.




Figure S13. ^1H NMR spectrum of compound **2** (TFA salt, 600 MHz, CD_3OD).




Figure S14. ^{13}C NMR spectrum of compound **2** (TFA salt, 150 MHz, CD_3OD).




Figure S15. HSQC spectrum of compound **2** (TFA salt, 600 MHz, CD₃OD).




Figure S16. HMBC spectrum of compound **2** (TFA salt, 600 MHz, CD₃OD).




Figure S17. COSY spectrum of compound **2** (TFA salt, 600 MHz, CD₃OD).




Figure S18. NOESY spectrum of compound **2** (TFA salt, 600 MHz, CD₃OD).




Figure S19. HR-ESIMS spectrum of compound 2.




Figure S20. ^1H NMR spectrum of compound 3 (TFA salt, 600 MHz, CD_3OD).




Figure S21. ^{13}C NMR spectrum of compound 3 (TFA salt, 150 MHz, CD_3OD).




Figure S22. HSQC spectrum of compound 3 (TFA salt, 600 MHz, CD₃OD).




Figure S23. HMBC spectrum of compound 3 (TFA salt, 600 MHz, CD₃OD).




Figure S24. COSY spectrum of compound 3 (TFA salt, 600 MHz, CD₃OD).




Figure S25. NOESY spectrum of compound 3 (TFA salt, 600 MHz, CD₃OD).




Figure S26. HR-ESIMS spectrum of compound 3.

