





Geophysical Research Letters®

RESEARCH LETTER

10.1029/2023GL107741

Slow Slip Events in New Zealand: Irregular, yet Predictable?

S. Truttmann¹ , T. Poulet² , L. Wallace^{3,4,5} , M. Herwegh¹ , and M. Veveakis⁶

Key Points:

- Nonlinear analysis of GNSS displacement time series unveils evidence for deterministic chaos in slow slip events in New Zealand
- Our theoretical findings imply that irregularly occurring slow slip events could potentially be forecasted a few weeks in advance

Supporting Information:

Supporting Information may be found in the online version of this article.

Correspondence to:

S. Truttmann,
sandro.truttmann@unibe.ch

Citation:

Truttmann, S., Poulet, T., Wallace, L., Herwegh, M., & Veveakis, M. (2024). Slow slip events in New Zealand: Irregular, yet predictable? *Geophysical Research Letters*, *51*, e2023GL107741. <https://doi.org/10.1029/2023GL107741>

Received 7 DEC 2023
Accepted 26 FEB 2024

Author Contributions:

Conceptualization: S. Truttmann, T. Poulet, M. Veveakis
Formal analysis: S. Truttmann, T. Poulet, L. Wallace
Methodology: S. Truttmann, T. Poulet, M. Veveakis
Project administration: M. Herwegh
Software: S. Truttmann, T. Poulet
Supervision: T. Poulet, M. Herwegh
Visualization: S. Truttmann, T. Poulet
Writing – original draft: S. Truttmann
Writing – review & editing: S. Truttmann, T. Poulet, L. Wallace, M. Herwegh, M. Veveakis

© 2024. The Authors.

This is an open access article under the terms of the [Creative Commons Attribution License](https://creativecommons.org/licenses/by/4.0/), which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

¹Institute of Geological Sciences, University of Bern, Bern, Switzerland, ²CSIRO Mineral Resources, Kensington, WA, Australia, ³GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany, ⁴Institute of Geosciences, Christian-Albrechts-Universität zu Kiel, Kiel, Germany, ⁵University of Texas Institute for Geophysics, Austin, TX, USA, ⁶Civil and Environmental Engineering, Duke University, Durham, NC, USA

Abstract Current earthquake forecasting approaches are mainly based on probabilistic assumptions, as earthquakes seem to occur randomly. Such apparent randomness can however be caused by deterministic chaos, rendering deterministic short-term forecasts possible. Due to the short historical and instrumental record of earthquakes, chaos detection has proven challenging, but more frequently occurring slow slip events (SSE) are promising candidates to probe for determinism. Here, we characterize the SSE signatures obtained from GNSS position time series in the Hikurangi Subduction Zone (New Zealand) to investigate whether the seemingly random SSE occurrence is governed by chaotic determinism. We find evidence for deterministic chaos for stations recording shallow SSEs, suggesting that short-term deterministic forecasting of SSEs, similar to weather forecasts, might indeed be possible over timescales of a few weeks. We anticipate that our findings could open the door for next-generation SSE forecasting, adding new tools to existing probabilistic approaches.

Plain Language Summary Since earthquakes appear to occur randomly, the currently available probabilistic predictions are based on past earthquake records. These predictions estimate the likelihood of an earthquake of a given magnitude occurring within a defined time period. In contrast to such probabilistic approaches, deterministic systems are fully predictable, albeit often confined to short time scales due to their potential chaotic behavior. Probing for deterministic predictability in the earthquake cycle is intractable due to the limited historical instrumental record. However, frequently occurring slow slip events - captured by transient GNSS displacements that can last several weeks - provide a unique opportunity to explore deterministic predictability in these types of slow earthquakes. By studying GNSS time series from various stations on New Zealand's North Island, we have discovered evidence suggesting that these irregularly occurring slow slip events might be governed by chaotic determinism. This implies the potential to forecast both timing and magnitude of slow slip events a few weeks in advance using deterministic methods, much like we predict weather patterns. Consequently, our theoretical findings could therefore pave the way for innovative approaches to short-term slow slip forecasting.

1. Introduction

Tectonic plate movements can lead to various modes of deformation, ranging from slow slip to fast earthquakes. Accurately forecasting such slip events remains a significant challenge, with current methods primarily relying on probabilistic considerations such as frequency-magnitude distributions (Jordan et al., 2011). These types of forecasts are based on the recordings of past events, and likelihood of future earthquakes occurrence is expressed as a probability that an event with a certain magnitude will occur within a certain time and area, with the assumption of a time-independent, linear stochastic (or random) process (e.g., Gerstenberger et al., 2020), with recent models also incorporating time-dependent characteristics (e.g., Field et al., 2015; Gerstenberger et al., 2022).

A fundamentally different approach is to consider deterministic forecasts which yield a single possible outcome only. The determinism could stem from the understanding of underlying physical processes, such as assuming that earthquakes originate from frictional sliding, which is a nonlinear process (Urbakh et al., 2004). In nonlinear systems, small changes in the input can lead to disproportionate, and often non-intuitive outputs. Such deterministic nonlinearity can result in chaotic systems, which are characterized by their irregularity and apparent randomness, yet are still deterministic and predictable at short timescales (Kantz & Schreiber, 2003). If analyzed with conventional linear tools (e.g., Fourier transform), chaotic time series appear random, but are in fact full of patterns only revealed with nonlinear analysis techniques (Abarbanel, 1996). Previous studies have suggested that both




Figure 1. (a) Map of the GNSS stations of the GeoNet monitoring network (www.geonet.org.nz; blue dots) and cumulative slow slip (2002–2014; red contours in mm) recorded in the Hikurangi subduction zone (after Wallace, 2020). Gray dashed lines represent depth contours of the subduction interface (in km below sea level; data from Williams et al., 2013). (b) Time series of the daily displacement signals (blue: data; red: low-pass filtered data) measured at four different GNSS stations (all E component) (GNS Science, 2000). SSEs are identified as distinctive jumps in the GNSS position timeseries, followed by slow westward drifting Inter-SSE periods. Black lines in the bottom of each subfigure show the derived daily velocities.

slow and fast fault slip might be governed by deterministic chaos (Barbot, 2019; Gualandi et al., 2020; Huang & Turcotte, 1990a, 1990b; Kato, 2016; Nie & Barbot, 2021; Poulet et al., 2014; Shelly, 2010; Veveakis et al., 2017).

Due to the long recurrence times of major earthquakes, data to identify deterministic behavior for seismic slip is currently insufficient. However, the frequently occurring slow slip events (SSE) with long rise times and slow rupture velocities do not suffer from this limitation and provide an opportunity for the detection of determinism. Land surface displacement during SSEs has been detected over the last two decades with continuously operating GNSS at many plate boundaries. In some cases, such as Cascadia and New Zealand, these networks have been operating for 20 years or more, and have recorded more than a dozen slow slip cycles (Schmidt & Gao, 2010; Szeliga et al., 2008; Wallace, 2020; Wallace & Beavan, 2010).

In the North Island of New Zealand, SSEs occur frequently but irregularly on the Hikurangi subduction zone (SZ) interface (Figure 1a). A large variety of events have been detected by the continuously operating GNSS monitoring network of GeoNet (www.geonet.org.nz; Wallace, 2020; Wallace & Beavan, 2010). Shallow SSEs (<15 km depth) that are short in duration (<1 month) and occur frequently (every 1–2 years) are observed offshore the northern and central East Coast. At the southern Hikurangi subduction zone, where most of the interface appears to be interseismically locked to ~25–30 km depth, SSEs occur at greater depths (>30 km), last longer (>1 year), and seem less frequent (ca. every 5 years).

In this study, we aim to explore the presence of deterministic chaos in the continuous GNSS time series recording SSEs in the Hikurangi SZ using nonlinear analysis techniques. If SSEs indeed exhibit characteristics of deterministic chaos, it would enable to go beyond probabilistic approaches and predict both timing and magnitude of SSEs at the short term.

2. Methods and Results

2.1. Phase Space Reconstruction

We use the daily position timeseries data of various GNSS stations as the surface manifestation of the spatio-temporal evolution of slip events at depth, to investigate the dynamical properties of the SSEs (Figure 1) (GNS




Figure 2. (a) Time series of measured displacements (blue; filtered signal in red) and daily velocities (black). (b) The three first singular vectors (SV) derived with the singular system analysis (SSA) of the daily velocities for the station GISB. (c) Reconstruction of the phase space for data from the station GISB, using the first three SVs as coordinates. (d)–(f) Examples of the phase space for shallow (station PAWA), deep (station TAKP), and no obvious SSE signals (station RIPA).

Science, 2000). We therefore employ low-pass Finite Impulse Response filtering using a Hamming window with a cut-off frequency f_c of $1/50$ and a window length L of 60 days to reduce measurement noise (see Text S1 in Supporting Information S1). We then derive the position differences, yielding daily velocities $v(t)$ (Figure 2a), and use singular system analysis (SSA) with an SSA window width M of 60, to decompose the signal into different singular vectors (SV) (Figure 2b). The values t_i of these vectors at a specific time can then be used as new coordinates to reconstruct the multidimensional phase space of the SSE system (Broomhead & King, 1986; King et al., 1987) (see Text S2 in Supporting Information S1) (Figure 2c). The phase space describes all possible states of a dynamical system, with each state t_i defined as a point, and each degree of freedom represented by an axis. The temporal sequence of all states forms a trajectory, along which the system evolves in time. If the trajectories are structured in phase space, they form an attractor that is an invariant of the system and can reveal otherwise undetected patterns (Abarbanel, 1996). Since the reconstructed attractor represents a proxy of the original system,

it provides the basis for the detection of deterministic chaos and forecasting with nonlinear analysis tools (Kantz & Schreiber, 2003).

Figure 2b displays the three first left singular vectors (SV) obtained through the SSA of $v(t)$ from GNSS station GISB. The phase space, reconstructed from these SVs, reveals a structured attractor (Figure 2c). This attractor comprises well-defined, loop-shaped trajectories with upward-folded sides, where each loop corresponds to an SSE. The size of each trajectory is directly proportional to the SSE's magnitude. During Inter-SSE periods, the system occupies a small subspace near the attractor node only, while significant changes in the SVs occur during SSE activity. Analyzing data from multiple GNSS stations, we found that stations clearly recording shallow SSEs have well-defined attractors with similar structures, such as GISB and PAWA (Figures 2c and 2d). The attractors of deep SSEs, exemplified by the station TAKP, exhibit a similar dynamical fingerprint as well, but with significantly fewer loops (Figure 2e). Due to their longer recurrence intervals of 4–5 years and fewer SSE cycles observed, the attractors of deep SSEs are more sparsely mapped. Therefore, we do not attempt to further quantify properties of the deep SSEs. At stations which were not significantly affected by SSEs, we were not able to obtain well-defined attractors (see RIPA, Figure 2f).

The phase space's dimensionality approximates the degrees of freedom of the underlying physical problem (Theiler, 1990), and low-dimensional spaces are characteristic for deterministic chaos (see Section 2.3). The dimensionality of the phase space, called embedding dimension d , can be estimated directly from the spectrum of singular values (Broomhead & King, 1986; Ghil et al., 2002; King et al., 1987) (see Text S2 in Supporting Information S1). We find dimensionalities d of approximately six for shallow and 35 for deep SSEs (Figure S2 in Supporting Information S1). The significantly larger dimensionality for the deep SSEs is likely due to noise, since larger values of M decrease the significance of the individual SVs and smooth the signal (Figures S3–S5 in Supporting Information S1) (Vautard & Ghil, 1989; Walwer et al., 2016). Since d approximates a theoretical upper bound of the number of degrees of freedom (Vautard & Ghil, 1989), it provides a first-order indication of the presence of a low-dimensional system, at least for the shallow SSEs. It is important to note that d is not an invariant of the system, but depends on data quality, sampling rate, and window width (Broomhead & King, 1986; Vautard & Ghil, 1989). Varying both low-pass filtering and M does however not significantly change the values of d (Figures S6–S8 in Supporting Information S1).

2.2. Nonlinearity

The observation of irregular recurrence times and the unraveling of structure in phase space with geometrically complex attractors, consisting of many different-sized trajectories, arguably point to a chaotic system. Nonlinearity is a necessary but not sufficient condition for deterministic chaos (Kantz & Schreiber, 2003), and fortunately, the detection of nonlinearity is relatively insensitive to noise (Theiler, 1992). A popular method to test for nonlinearity in a time series from an unknown dynamical system is surrogate data hypothesis testing (Kantz & Schreiber, 2003; Lancaster et al., 2018; Theiler, 1992) (see Text S3 in Supporting Information S1). We use wavelet iterative amplitude adjusted Fourier transform (WIAAFT) surrogates to test the null hypothesis that the signals can be explained by a non-stationary linear stochastic process, potentially with a nonlinear measurement transformation (Keylock, 2006). If the values of a discriminatory statistic, here Takens' best estimate of the correlation dimension $D_2^{(T)}$, for 100 realization of surrogates differ significantly from the statistics computed from the measured time series, we reject the null hypothesis (Lancaster et al., 2018; Theiler, 1992). Since a deterministic nonlinear system supposedly leads to a lower dimension of the attractor, we reject the null hypothesis if $D_2^{(T)}$ of the measured time series is smaller than the fifth percentile of the surrogates (Lancaster et al., 2018; Theiler, 1992).

Using this approach, we can reject the null hypothesis for 11 different GNSS time series, all recording shallow East Coast SSEs (Figure 3). These measurements thus cannot be explained by a linear stochastic process, and we can assume nonlinearity. Interestingly, the detected nonlinear signals appear in two spatial clusters, both located in close proximity to the regions with maximum cumulative slow slip (Figure 3a). This suggests that the shallow SSEs in the nearby source regions are nonlinear systems that might even be chaotic (Theiler, 1992). For all other stations recording only a few SSEs, including stations recording deep SSEs, we could not reject the null hypothesis, which does however not explicitly exclude nonlinearity.




Figure 3. (a) Map of the GNSS stations, colored after the results of the surrogate data hypothesis tests. Stations with detected nonlinearity (red squares) cluster in close proximity to locations of largest onshore displacement during shallow SSEs offshore the east coast. We note that the stations recording deep SSEs only incorporate a few SSE cycles (due to their longer recurrence intervals), so they do not record a sufficient number of SSEs to assess nonlinearity (though this cannot be ruled-out). (b) Box-whisker plots of the surrogate data test for a selection of stations along the east coast. The whiskers indicate the fifth, and 95th percentiles of the surrogate distributions. Blue diamonds show Takens' best estimate of the correlation dimension $D_2^{(T)}$ of the measured time series.

2.3. Indications of Deterministic Chaos

In contrast to finding evidence for nonlinearity, detecting deterministic chaos in short and noisy time series has proven more challenging (Casdagli, 1992; Kantz & Schreiber, 2003; Theiler, 1992). The revealed structure in phase space (Figure 2), as well as the detected nonlinearity (Figure 3), gives a first hint for the presence of chaos (Abarbanel, 1996). Further invariants that are indicative of chaotic systems are finite-dimensional fractal (or strange) attractors and exponential divergence of nearby states. However, due to the noisy and limited data available, quantifying these invariants has proven difficult (Eckmann & Ruelle, 1992; Theiler, 1992). Therefore, we further analyze the data from the station GISB only, which records shallow SSEs exceptionally well and is, in our view, the most suitable site in the Hikurangi SZ for this analysis.

The qualitatively self-similar attractor shown in Figure 2c suggests a geometrically complex, potentially fractal attractor (Casdagli, 1992). Estimating the correlation dimension D_2 with an entropy-based approach (see Text S4), we were unable to identify a scaling region or derive a value of D_2 , regardless of the filters used (Figures S9–S12 in Supporting Information S1). This is not surprising, given that the measurement noise obscures the small-scale features of the attractor (Datseris & Parlitz, 2022; Kantz & Schreiber, 2003). A stochastic process should however result in a constant increase of D_2 with increasing d (Casdagli, 1992; Kantz & Schreiber, 2003). We thus argue that the saturation of D_2 with increasing d (Figures S9–S12 in Supporting Information S1) can be viewed as a first-order indication of a low-dimensional system.

Chaotic systems are governed by exponential divergence of nearby states, expressed by a positive Lyapunov exponent λ_1 (see Text S5 in Supporting Information S1) (Kantz & Schreiber, 2003). Directly deriving λ_1 from the attractors (Skokos et al., 2016), we found positive λ_1 values in the range of 8–12, dependent on the filtering parameters (Figures S13–S16 in Supporting Information S1). Due to the exponential divergence of nearby states, small variations in the initial conditions lead to vast differences in the outcome, rendering chaotic systems unpredictable at longer time-scales despite their deterministic nature. The Lyapunov time t_λ , defined as the inverse of λ_1 , serves as a measure of the characteristic timescale on which a system is chaotic and provides a theoretical




Figure 4. Conceptual example of local modeling-based short-term SSE forecast for the station GISB, using the GNSS position time series from 2002 to September 2020 as training data. (a) Training data (green to yellow) and forecast (orange) in phase space. (b) Forecast converted to the GNSS position time series (orange line), compared to the measured displacement (blue; filtered signal in red) and the derived daily velocities (black). The vertical solid line marks the end of observations and start of the forecast (current state), and the dashed white line indicates the theoretical forecast limit defined by the Lyapunov time t_λ .

limit of the predictability. Values of λ_1 between 8 and 12 suggest that the shallow SSEs recorded at GISB should be predictable to within 1/8 to 1/12 of the approximate recurrence period of the shallow SSEs of about one year, implying that the onset of shallow SSEs might be forecast about 30–45 days in advance.

3. Discussion

Compared to the Cascadia SZ, SSEs in New Zealand occur quite irregularly. This is not surprising, given the more heterogeneous structure (Barnes et al., 2020; Gase et al., 2022) and slip behavior of the Hikurangi SZ, with substantial spatial variation in the distribution of SSEs and interseismic coupling (Wallace, 2020). Our findings imply that, despite appearing random, the SSEs in New Zealand might still be deterministic and their onset could be forecast a few weeks in advance, remarkably similar to observations from the Cascadia SZ (Gualandi et al., 2020). We argue that the irregular recurrence of SSEs in New Zealand might be governed by stress interactions between different SSE source regions, since deterministic chaos can arise from the coupling of different systems (Huang & Turcotte, 1990b).

As the amount of currently available data is limited, only a subset of the SSE attractors is mapped, and the attractors likely consist of an infinite number of different-sized trajectories, explaining the system's variability in magnitudes. This self-similarity observed at larger scales suggests that SSEs with very small magnitudes and short durations of a few days might occur during Inter-SSE periods. Such hidden SSEs are however obscured by measurement noise, and stronger filtering for noise reduction would remove the signals of such tiny SSEs as well, making their identification difficult (Frank, 2016; Jolivet & Frank, 2020; Kato & Nakagawa, 2020). We suggest that SSEs in New Zealand occur more frequently than observed with the currently available data.

The existence of deterministic chaos opens the door for next-generation SSE forecasting, since the system's short-term behavior can directly be derived from the phase space, for example, using local modeling (Abarbanel, 1996; Engster & Parlitz, 2006) (see Text S6 in Supporting Information S1). To illustrate this conceptually, we use the GNSS data from 2002 to September 2020 of station GISB as training data to reconstruct the phase space (Figure 4a). The trajectories closest to the most recent measurement then approximate the future evolution of the system in phase space (Figure 4a), which can be converted back to the measured GNSS position time series (Figure 4b). In the case of the example in Figure 4, the short-term forecast of the daily positions appears relatively accurate. Beyond the theoretical forecast limit, the chaotic nature of the system renders long-term forecasts impossible, which is illustrated by the mismatch of the SSE forecast in late 2021. However, the theoretical limit

derived from λ_1 is a rather conservative estimate of the system's predictability (Datseris & Parlitz, 2022). For accurate short-term forecasts, a good description of the phase space is required, which is not currently fulfilled with the limited amount of data available, which maps the phase space sparsely (Figures 2c and 4a, and Figure S17 in Supporting Information S1). Potential ways to overcome these limitations could be enhanced data quality (i.e., higher signal-to-noise ratio), longer time series, or the incorporation of physics-based models. In our view, the well-recorded shallow SSEs represent the most promising systems for effective short-term SSE forecasting in New Zealand.

4. Conclusion

In summary, indications for nonlinearity, sensitivity to initial conditions, and self-similarity suggest that the irregularly occurring SSEs in New Zealand may exhibit deterministic chaos. We demonstrate that, despite the apparent randomness, with long and precise enough datasets, there appears to be some degree of predictability in SSEs. Similar to forecasts of chaotic weather systems, both timing and magnitude of SSEs have potential to be predicted a few weeks in advance. As SSEs lead to stress changes on adjacent parts of the subduction interface which can trigger large earthquakes (e.g., Kato et al., 2012; Koulali et al., 2017; Radiguet et al., 2016; Segou & Parsons, 2020), short-term SSE forecasting might also allow to identify phases of enhanced earthquake hazard (Obara & Kato, 2016). Evidence that also seismic slip might be chaotic (e.g., Barbot, 2019; Kato, 2016; Shelly, 2010) suggests that similar deterministic earthquake forecasts could one day be possible when long enough time series of the seismic cycle become available.

Data Availability Statement

The data that support the findings of this study are openly available from GNS Science at <http://doi.org/10.21420/30F4-1A55> (GNS Science, 2000).

Acknowledgments

S. Truttmann thankfully acknowledges funding from the Swiss Geophysical Commission (SGPK) and an UniBE Doc. Mobility travel grant. L. Wallace acknowledges funding from NSF grant EAR-212166. M. Veveakis acknowledges the support of the U.S. National Science Foundation through project CMMI-2042325.

References

- Abarbanel, H. D. I. (1996). *Analysis of observed chaotic data*. Springer New York. <https://doi.org/10.1007/978-1-4612-0763-4>
- Barbot, S. (2019). Slow-slip, slow earthquakes, period-two cycles, full and partial ruptures, and deterministic chaos in a single asperity fault. *Tectonophysics*, 768, 228171. <https://doi.org/10.1016/j.tecto.2019.228171>
- Barnes, P. M., Wallace, L. M., Saffer, D. M., Bell, R. E., Underwood, M. B., Fagereng, A., et al. (2020). Slow slip source characterized by lithological and geometric heterogeneity. *Science Advances*, 6(13), eaay3314. <https://doi.org/10.1126/sciadv.aay3314>
- Broomhead, D. S., & King, G. P. (1986). Extracting qualitative dynamics from experimental data. *Physica D: Nonlinear Phenomena*, 20(2–3), 217–236. [https://doi.org/10.1016/0167-2789\(86\)90031-X](https://doi.org/10.1016/0167-2789(86)90031-X)
- Casdagli, M. (1992). Chaos and deterministic versus stochastic non-linear modelling. *Journal of the Royal Statistical Society: Series B (Methodological)*, 54(2), 303–328. <https://doi.org/10.1111/j.2517-6161.1992.tb01884.x>
- Datseris, G., & Parlitz, U. (2022). *Nonlinear dynamics: A concise introduction interlaced with code*. Springer International Publishing. <https://doi.org/10.1007/978-3-030-91032-7>
- Eckmann, J.-P., & Ruelle, D. (1992). Fundamental limitations for estimating dimensions and Lyapunov exponents in dynamical systems. *Physica D*, 56(2–3), 185–187. [https://doi.org/10.1016/0167-2789\(92\)90023-G](https://doi.org/10.1016/0167-2789(92)90023-G)
- Engster, D., & Parlitz, U. (2006). Local and cluster weighted modeling for time series prediction. In *Handbook of Time Series Analysis* (pp. 39–66).
- Field, E. H., Biasi, G. P., Bird, P., Dawson, T. E., Felzer, K. R., Jackson, D. D., et al. (2015). Long-term time-dependent probabilities for the Third Uniform California Earthquake Rupture Forecast (UCERF3). *Bulletin of the Seismological Society of America*, 105(2A), 511–543. <https://doi.org/10.1785/0120140093>
- Frank, W. B. (2016). Slow slip hidden in the noise: The intermittence of tectonic release. *Geophysical Research Letters*, 43(19). <https://doi.org/10.1002/2016GL069537>
- Gase, A. C., Bangs, N. L., Van Avendonk, H. J. A., Bassett, D., & Henrys, S. A. (2022). Hikurangi megathrust slip behavior influenced by lateral variability in sediment subduction. *Geology*, 50(10), 1145–1149. <https://doi.org/10.1130/G50261.1>
- Gerstenberger, M. C., Marzocchi, W., Allen, T., Pagani, M., Adams, J., Danciu, L., et al. (2020). Probabilistic seismic hazard analysis at regional and national scales: State of the art and future challenges. *Reviews of Geophysics*, 58(2). <https://doi.org/10.1029/2019RG000653>
- Gerstenberger, M. C., Van Dissen, R. J., Rollins, C., DiCaprio, C., Chamberlain, C., Christophersen, A., et al. (2022). The seismicity rate model for the 2022 New Zealand National Seismic Hazard Model. In *GNS Science Report, 2022/47* (p. 156). <https://doi.org/10.21420/2EXG-NP48>
- Ghil, M., Allen, M. R., Dettinger, M. D., Ide, K., Kondrashov, D., Mann, M. E., et al. (2002). Advanced spectral methods for climatic time series: Climatic time series analysis. *Reviews of Geophysics*, 40(1), 3–3–41. <https://doi.org/10.1029/2000RG000092>
- GNS Science (2000). GeoNet Aotearoa New Zealand Continuous GNSS Network Time Series Dataset. [Dataset]. GNS Science, GeoNet. <https://doi.org/10.21420/30F4-1A55>
- Gualandi, A., Avouac, J.-P., Michel, S., & Faranda, D. (2020). The predictable chaos of slow earthquakes. *Science Advances*, 6(27), eaaz5548. <https://doi.org/10.1126/sciadv.aaz5548>
- Huang, J., & Turcotte, D. L. (1990a). Are earthquakes an example of deterministic chaos? *Geophysical Research Letters*, 17(3), 223–226. <https://doi.org/10.1029/GL017i003p00223>
- Huang, J., & Turcotte, D. L. (1990b). Evidence for chaotic fault interactions in the seismicity of the San Andreas fault and Nankai trough. *Nature*, 348(6298), 234–236. <https://doi.org/10.1038/348234a0>

- Jolivet, R., & Frank, W. B. (2020). The transient and intermittent nature of slow slip. *AGU Advances*, 1(1). <https://doi.org/10.1029/2019AV000126>
- Jordan, T. H., Chen, Y.-T., Gasparini, P., Madariaga, R., Main, I., Marzocchi, W., et al. (2011). Operational earthquake forecasting: State of knowledge and guidelines for implementation. *Annals of Geophysics*, 4(54). <https://doi.org/10.4401/ag-5350>
- Kantz, H., & Schreiber, T. (2003). *Nonlinear time series analysis* (2nd ed.). Cambridge University Press. <https://doi.org/10.1017/CBO9780511755798>
- Kato, A., & Nakagawa, S. (2020). Detection of deep low-frequency earthquakes in the Nankai subduction zone over 11 years using a matched filter technique. *Earth Planets and Space*, 72(1), 128. <https://doi.org/10.1186/s40623-020-01257-4>
- Kato, A., Obara, K., Igarashi, T., Tsuruoka, H., Nakagawa, S., & Hirata, N. (2012). Propagation of slow slip leading up to the 2011 M_w 9.0 Tohoku-Oki earthquake. *Science*, 335(6069), 705–708. <https://doi.org/10.1126/science.1215141>
- Kato, N. (2016). Earthquake cycles in a model of interacting fault patches: Complex behavior at transition from seismic to aseismic slip. *Bulletin of the Seismological Society of America*, 106(4), 1772–1787. <https://doi.org/10.1785/0120150185>
- Keylock, C. J. (2006). Constrained surrogate time series with preservation of the mean and variance structure. *Physical Review E*, 73(3), 036707. <https://doi.org/10.1103/PhysRevE.73.036707>
- King, G. P., Jones, R., & Broomhead, D. S. (1987). Phase portraits from a time series: A singular system approach. *Nuclear Physics B - Proceedings Supplements*, 2, 379–390. [https://doi.org/10.1016/0920-5632\(87\)90029-6](https://doi.org/10.1016/0920-5632(87)90029-6)
- Koulali, A., McClusky, S., Wallace, L., Allgeyer, S., Tregoning, P., D'Anastasio, E., & Benavente, R. (2017). Slow slip events and the 2016 Te Araroa M_w 7.1 earthquake interaction: Northern Hikurangi subduction, New Zealand. *Geophysical Research Letters*, 44(16), 8336–8344. <https://doi.org/10.1002/2017GL074776>
- Lancaster, G., Iatsenko, D., Pidde, A., Ticcinelli, V., & Stefanovska, A. (2018). Surrogate data for hypothesis testing of physical systems. *Physics Reports*, 748, 1–60. <https://doi.org/10.1016/j.physrep.2018.06.001>
- Nie, S., & Barbot, S. (2021). Seismogenic and tremorgenic slow slip near the stability transition of frictional sliding. *Earth and Planetary Science Letters*, 569, 117037. <https://doi.org/10.1016/j.epsl.2021.117037>
- Obara, K., & Kato, A. (2016). Connecting slow earthquakes to huge earthquakes. *Science*, 353(6269), 253–257. <https://doi.org/10.1126/science.aaf1512>
- Poulet, T., Veveakis, E., Regenauer-Lieb, K., & Yuen, D. A. (2014). Thermo-poro-mechanics of chemically active creeping faults: 3. The role of serpentinite in episodic tremor and slip sequences, and transition to chaos. *Journal of Geophysical Research: Solid Earth*, 119(6), 4606–4625. <https://doi.org/10.1002/2014JB011004>
- Radiguet, M., Perfettini, H., Cotte, N., Gualandi, A., Valette, B., Kostoglodov, V., et al. (2016). Triggering of the 2014 M_w 7.3 Papanoa earthquake by a slow slip event in Guerrero, Mexico. *Nature Geoscience*, 9(11), 829–833. <https://doi.org/10.1038/ngeo2817>
- Schmidt, D. A., & Gao, H. (2010). Source parameters and time-dependent slip distributions of slow slip events on the Cascadia subduction zone from 1998 to 2008. *Journal of Geophysical Research*, 115(B4), B00A18. <https://doi.org/10.1029/2008JB006045>
- Segou, M., & Parsons, T. (2020). The role of seismic and slip slip events in triggering the 2018 $M7.1$ Anchorage earthquake in the Southcentral Alaska subduction zone. *Geophysical Research Letters*, 47(10), e2019GL086640. <https://doi.org/10.1029/2019GL086640>
- Shelly, D. R. (2010). Periodic, chaotic, and doubled earthquake recurrence intervals on the deep san Andreas fault. *Science*, 328(5984), 1385–1388. <https://doi.org/10.1126/science.1189741>
- Skokos, C., Gottwald, G. A., & Laskar, J. (Eds.). (2016). *Chaos detection and predictability* (Vol. 915). Springer Berlin Heidelberg. <https://doi.org/10.1007/978-3-662-48410-4>
- Szeliga, W., Melbourne, T., Santillan, M., & Miller, M. (2008). GPS constraints on 34 slow slip events within the Cascadia subduction zone, 1997–2005. *Journal of Geophysical Research*, 113(B4), B04404. <https://doi.org/10.1029/2007JB004948>
- Theiler, J. (1990). Estimating fractal dimension. *Journal of the Optical Society of America A*, 7(6), 1055–1073. <https://doi.org/10.1364/JOSAA.7.001055>
- Theiler, J., Eubank, S., Longtin, A., Galdrikian, B., & Doynne Farmer, J. (1992). Testing for nonlinearity in time series: The method of surrogate data. *Physica D: Nonlinear Phenomena*, 58(1–4), 77–94. [https://doi.org/10.1016/0167-2789\(92\)90102-S](https://doi.org/10.1016/0167-2789(92)90102-S)
- Urbakh, M., Klafter, J., Gourdon, D., & Israealachvili, J. (2004). The nonlinear nature of friction. *Nature*, 430(6999), 525–528. <https://doi.org/10.1038/nature02750>
- Vautard, R., & Ghil, M. (1989). Singular spectrum analysis in nonlinear dynamics, with applications to paleoclimatic time series. *Physica D: Nonlinear Phenomena*, 35(3), 395–424. [https://doi.org/10.1016/0167-2789\(89\)90077-8](https://doi.org/10.1016/0167-2789(89)90077-8)
- Veveakis, E., Alevizos, S., & Poulet, T. (2017). Episodic Tremor and Slip (ETS) as a chaotic multiphysics spring. *Physics of the Earth and Planetary Interiors*, 264, 20–34. <https://doi.org/10.1016/j.pepi.2016.10.002>
- Wallace, L. M. (2020). Slow slip events in New Zealand. *Annual Review of Earth and Planetary Sciences*, 48(1), 175–203. <https://doi.org/10.1146/annurev-earth-071719-055104>
- Wallace, L. M., & Beavan, J. (2010). Diverse slow slip behavior at the Hikurangi subduction margin, New Zealand. *Journal of Geophysical Research*, 115(B12), B12402. <https://doi.org/10.1029/2010JB007717>
- Walwer, D., Calais, E., & Ghil, M. (2016). Data-adaptive detection of transient deformation in geodetic networks. *Journal of Geophysical Research: Solid Earth*, 121(3), 2129–2152. <https://doi.org/10.1002/2015JB012424>
- Williams, C. A., Eberhart-Phillips, D., Bannister, S., Barker, D. H. N., Henrys, S., Reyners, M., & Sutherland, R. (2013). Revised interface geometry for the Hikurangi Subduction Zone, New Zealand. *Seismological Research Letters*, 84(6), 1066–1073. <https://doi.org/10.1785/0220130035>

References From the Supporting Information

- Ashkenazy, Y. (1999). The use of generalized information dimension in measuring fractal dimension of time series. *Physica A: Statistical Mechanics and its Applications*, 271(3–4), 427–447. [https://doi.org/10.1016/S0378-4371\(99\)00192-2](https://doi.org/10.1016/S0378-4371(99)00192-2)
- Broomhead, D. S., Huke, J. P., & Muldoon, M. R. (1992). Linear filters and non-linear systems. *Journal of the Royal Statistical Society: Series B (Methodological)*, 54(2), 373–382. <https://doi.org/10.1111/j.2517-6161.1992.tb01887.x>
- Casaleggio, A., & Marchesi, M. L. (1993). *Some results on the effect of digital filtering on the estimation of the Correlation Dimension*. In *IEEE Winter Workshop on Nonlinear Digital Signal Processing* (pp. 6.3_2.1–6.3_2.6). IEEE. <https://doi.org/10.1109/NDSP.1993.767747>
- Datseris, G. (2018). DynamicalSystems.jl: A Julia software library for chaos and nonlinear dynamics. *Journal of Open Source Software*, 3(23), 598. <https://doi.org/10.21105/joss.00598>

- Grassberger, P. (1983). Generalized dimensions of strange attractors. *Physics Letters*, 97A(6), 227–230. [https://doi.org/10.1016/0375-9601\(83\)90753-3](https://doi.org/10.1016/0375-9601(83)90753-3)
- Haaga, K. A., & Datsis, G. (2022). TimeseriesSurrogates.jl: A Julia package for generating surrogate data. *Journal of Open Source Software*, 7(77), 4414. <https://doi.org/10.21105/joss.04414>
- Hentschel, H. G. E., & Procaccia, I. (1983). The infinite number of generalized dimensions of fractals and strange attractors. *Physica D: Nonlinear Phenomena*, 8(3), 435–444. [https://doi.org/10.1016/0167-2789\(83\)90235-X](https://doi.org/10.1016/0167-2789(83)90235-X)
- Kantz, H. (1994). A robust method to estimate the maximal Lyapunov exponent of a time series. *Physics Letters A*, 185(1), 77–87. [https://doi.org/10.1016/0375-9601\(94\)90991-1](https://doi.org/10.1016/0375-9601(94)90991-1)
- Packard, N. H., Crutchfield, J. P., Farmer, J. D., & Shaw, R. S. (1980). Geometry from a time series. *Physical Review Letters*, 45(9), 712–716. <https://doi.org/10.1103/physrevlett.45.712>
- Rényi, A. (1959). On the dimension and entropy of probability distributions. *Acta Mathematica Academiae Scientiarum Hungaricae*, 10(1–2), 193–215. <https://doi.org/10.1007/BF02063299>
- Sauer, T., & Yorke, J. A. (1993). How many delay coordinates do you need? *International Journal of Bifurcation and Chaos*, 3(3), 737–744. <https://doi.org/10.1142/S0218127493000647>
- Takens, F. (1985). On the numerical determination of the dimension of an attractor. In B. L. J. Braaksma, H. W. Broer, & F. Takens (Eds.), *Dynamical systems and bifurcations* (Vol. 1125, pp. 99–106). Springer Berlin Heidelberg. <https://doi.org/10.1007/BFb0075637>
- Takens, F. (1981). *Detecting strange attractors in turbulence*. In D. Rand & L.-S. Young (Eds.), *Dynamical systems and turbulence, Warwick 1980* (Vol. 898, pp. 366–381). Springer Berlin Heidelberg. <https://doi.org/10.1007/BFb0091924>
- Theiler, J. (1987). Efficient algorithm for estimating the correlation dimension from a set of discrete points. *Physical Review A*, 36(9), 4456–4462. <https://doi.org/10.1103/PhysRevA.36.4456>
- Walwer, D., Ghil, M., & Calais, E. (2022). A data-based minimal model of episodic inflation events at volcanoes. *Frontiers in Earth Science*, 10, 759475. <https://doi.org/10.3389/feart.2022.759475>