

About Duval Extensions*

Tero Harju Dirk Nowotka
Turku Centre for Computer Science, TUCS
Department of Mathematics, University of Turku

June 2003

Abstract

A word $v = wu$ is a (nontrivial) Duval extension of the unbordered word w , if (u is not a prefix of v and) w is an unbordered factor of v of maximum length. A survey of the state of the art of research on Duval extensions is given in this paper.

1 Introduction

A survey about the research on the relationship between the length of a word and its unbordered factors is given in this paper. This line of research was introduced by Ehrenfeucht and Silberger [3] and Assous and Pouzet [1] in 1979. It was carried further and culminated in a strong conjecture by Duval [2] in 1982. Only recently this conjecture was proved in [6].

We will give a historical overview on this line of research, its main results and conjectures so far, in Section 2. This will lead to the concept of Duval extensions which are introduced in Section 3. The main result known so far on Duval extensions, Theorem 8, is presented there. The subsections 3.1, 3.2, and 3.3 will introduce further research lines on Duval extensions. We conclude with Section 4.

We shall now introduce the main notations of this paper. We refer the reader to [7, 8] for more basic and general definitions.

Consider a finite alphabet A of letters. Let A^* denote the monoid of all finite words over A including the empty word, denoted by ε . Let $w \in A^*$. Then we can express w as a sequence of letters $w_{(1)}w_{(2)} \cdots w_{(n)}$ where $w_{(i)} \in A$ is a letter, for every $1 \leq i \leq n$. We denote the length n of w by $|w|$. Note, that $|\varepsilon| = 0$. A word w is called *primitive* if it cannot be factored such that $w = u^k$ for some $k \geq 2$. Let $w = uv$ for some words u and v . Then vu is called *conjugate* of w . Let $[w]$ denote the set of all conjugates of w . Note, that $w \in [w]$.

*A preliminary version of the given survey is contained in an essay written for the Bulletin of the EATCS.

A nonempty word u is called a *border* of a word w , if $w = uv = v'u$ for some words v and v' . We call w *bordered*, if it has a border that is shorter than w , otherwise w is called *unbordered*. Note, that every bordered word w has a minimum border u such that $w = uvu$, where u is unbordered. Suppose $w = uv$, then u is called a *prefix* of w , denoted by $u \leq w$, and v is called a *suffix* of w , denoted by $v \preceq w$.

Let \triangleleft_A be an ordering of $A = \{a_1, a_2, \dots, a_n\}$, say $a_1 \triangleleft_A a_2 \triangleleft_A \dots \triangleleft_A a_n$. Then \triangleleft_A induces a *lexicographic order*, also denoted by \triangleleft_A , on A^* such that

$$u \triangleleft_A v \iff u \leq v \quad \text{or} \quad u = xau' \text{ and } v = xbu' \text{ with } a \triangleleft_A b$$

where $a, b \in A$. We write \triangleleft for \triangleleft_A , for some alphabet A , if the context is clear.

Let us consider the following examples. Let $A = \{a, b\}$ and $u, v, w \in A^*$ such that $u = abaa$ and $v = baaba$ and $w = abaaba$. Then u and v are primitive, but w is not. Furthermore, $[u] = \{aaab, aaba, abaa, baaa\}$ is the set of all conjugates of u . Let $a \triangleleft b$. Then $u \triangleleft w \triangleleft v$. We have that a is the shortest border of u and w , whereas ba is the shortest border of v . The smallest unbordered factor of w has length three.

2 The Maximum Length of Unbordered Factors

When the length of unbordered factors of a word is investigated, that is usually done in terms of the length of the word and its minimum period.

Lets make our terminology more precise. Consider a word w over some alphabet A . An integer $1 \leq p \leq n$ is a *period* of w , if $w_{(i)} = w_{(i+p)}$ for all $1 \leq i \leq n - p$. The smallest period of w is called the *minimum period* (or simply, the period) of w , denoted by $\partial(w)$. Let $\mu(w)$ denote the maximum length of unbordered factors of w . For example, let $w = abaabbaaba$, then $\partial(w) = 7$ and $\mu(w) = 6$.

Clearly, the maximum length of unbordered factors $\mu(w)$ of w is bound by the period $\partial(w)$ of w . We have

$$\mu(w) \leq \partial(w)$$

since for every factor v of w , with $\partial(w) < |v|$, the prefix $v_{(1)}v_{(2)} \dots v_{(|v|-\partial(w))}$ of v is also a suffix of v by the definition of period.

It is a natural question to ask at what length of w is $\mu(w)$ necessarily maximal, that is, $\mu(w) = \partial(w)$. Of course, the length of w is considered with respect to either $\mu(w)$ or $\partial(w)$.

In 1979 Ehrenfeucht and Silberger [3], as well as, Assous and Pouzet [1] addressed this question first. Ehrenfeucht and Silberger [3] stated

Theorem 1. *If $2\partial(w) \leq |w|$ then $\mu(w) = \partial(w)$.*

They also established that every primitive word w has at least σ -many unbordered conjugates, where σ is the number of different letters occuring in w , which leads directly to

Theorem 2. *If $2\partial(w) - \sigma \leq |w|$ then $\mu(w) = \partial(w)$.*

However, this result was stated by Duval [2] only in 1981.

The real challenge, though, turned out to be giving a bound on the length of w with respect to $\mu(w)$. It was conjectured in [3] that $2\mu(w) \leq |w|$ implies $\mu(w) = \partial(w)$. However, Assous and Pouzet gave the following counter example contradicting that conjecture.

Example 3. *Let*

$$w = a^n b a^{n+1} b a^n b a^{n+2} b a^n b a^{n+1} b a^n$$

for which $|w| = 7n + 10$ and $\mu(w) = 3n + 6$ and $\partial(w) = 4n + 7$.

Assous and Pouzet themselves gave the following conjecture.

Conjecture 4. *Let $f: \mathbb{N} \rightarrow \mathbb{N}$ such that $f(\mu(w)) \leq |w|$ implies $\mu(w) = \partial(w)$. Then*

$$f(\mu(w)) \leq 3\mu(w) .$$

In 1982 Duval [2] established the following.

Theorem 5. *If $4\mu(w) - 6 \leq |w|$ then $\mu(w) = \partial(w)$.*

He also stated Conjecture 7 (see next section) about what was later called Duval extensions that would imply

$$\text{If } 3\mu(w) \leq |w| \text{ then } \mu(w) = \partial(w) .$$

3 Duval Extensions

In the previous section we recalled a question initially raised by Ehrenfeucht and Silberger [3]. The problem was to estimate a bound on the length of w , depending on $\mu(w)$, such that $\mu(w) = \partial(w)$. Duval [2] introduced a restricted version of that problem by assuming that w has an unbordered prefix of length $\mu(w)$. Let us first fix some more notations first.

Let w and u be nonempty words where w is also unbordered. We call wu a *Duval extension* of w , if every factor of wu longer than $|w|$ is bordered, that is, $\mu(wu) = |w|$. A Duval extension wu is called *trivial*, if $\partial(wu) = \mu(wu)$. A nontrivial Duval extension wu of w is called *minimal*, if u is of minimal length, that is, $u = u'a$ and $w = u'bw'$ where $a, b \in A$ and $a \neq b$.

Example 6. *Let $w = abaabbabaababb$ and $u = aaba$. Then*

$$w.u = abaabbabaababb.aaba$$

(for the sake of readability, we use a dot to mark where w ends) is a non-trivial Duval extension of w of length $|wu| = 18$, where $\mu(wu) = |w| = 14$ and $\partial(wu) = 15$. However, wu is not a minimal Duval extension, whereas

$$w.u' = abaabbabaababb.aa$$

is minimal, with $u' = aa \leq u$. Note, that wu is not the longest nontrivial Duval extension of w since

$$w.v = abaabbabaababb.abaaba$$

is longer, with $v = abaaba$ and $|wv| = 20$ and $\partial(wv) = 17$. One can check that wv is a nontrivial Duval extension of w of maximum length, and at the same time wv is also a minimal Duval extension of w .

In 1982 Duval [2] stated the following conjecture.

Conjecture 7. *Let wu be a nontrivial Duval extension of w . Then $|u| < |w|$.*

It follows directly from this conjecture that for any word w , we have that $3\mu(w) \leq |w|$ implies $\mu(w) = \partial(w)$. Duval's conjecture has remained popular throughout the years, see for example Chapter 8 in [8]. Only recently, an improved version of this conjecture was proved by the authors of this paper; see [6].

Theorem 8. *Let wu be a nontrivial Duval extension of w . Then $|u| < |w| - 1$.*

This bound is tight as can be seen from the following example.

Example 9. *Let $w = a^n ba^{n+m} bb$ and $u = a^{n+m} ba^n$ with $n, m \geq 1$. Then*

$$w.u = a^n ba^{n+m} bb.a^{n+m} ba^n$$

is a nontrivial Duval extension of w and $|u| = |w| - 2$.

We get the following corollary from Theorem 8.

Corollary 10. *If $3\mu(w) - 2 \leq |w|$ then $\mu(w) = \partial(w)$.*

This is the best bound in the general case known to us so far. However, Duval extensions have also become a subject of interest on their own. We will investigate them more closely in the next three subsections.

3.1 Words without Nontrivial Duval Extensions

The set of words having no nontrivial Duval extension has been investigated in [4] and [10]. We recall these results here.

Infinite words of minimal subword complexity are called *Sturmian* words, cf. [11, 8]. Minimal subword complexity means that a Sturmian word contains exactly $n + 1$ different factors of length n for every $n \geq 1$. Let us consider finite factors of Sturmian words in the following, and let's simply call them Sturmian words. Mignosi and Zamboni showed the following uniqueness result for Duval extensions in [10].

Theorem 11. *Unbordered Sturmian words have no nontrivial Duval extension.*

This result was improved by the authors of this paper in [4] to Lyndon words. Let a primitive word w be called *Lyndon* word if it is minimal among its conjugates, that is, if $w \triangleleft v$ for every $v \in [w]$ and some arbitrary order \triangleleft on A , cf. [9, 8]. Note, that Lyndon words are unbordered.

Theorem 12. *Lyndon words have no nontrivial Duval extension.*

Theorem 14 states that unbordered Sturmian words are indeed Lyndon words. The following lemma will be used to prove that result.

Let $\tau: A^* \rightarrow B^*$ be a morphism, and \triangleleft_A and \triangleleft_B be orders on A and B , respectively, such that

$$a_1 \triangleleft_A a_2 \implies \tau(a_1) \triangleleft_B \tau(a_2) \tag{1}$$

for every $a_1, a_2 \in A$, and $\tau(a)$ is a Lyndon word w.r.t. \triangleleft_B for every $a \in A$.

Lemma 13. *If $w \in A^*$ is a Lyndon word, then $\tau(w)$ is a Lyndon word.*

The following theorem shows that Theorem 12 implies Theorem 11.

Theorem 14. *Every unbordered Sturmian word is a Lyndon word.*

Proofs of Lemma 13 and Theorem 14 can be found in [5]. The converse of Theorem 14 is certainly not true. Indeed, consider the word *aabbab* which is a Lyndon word but not a Sturmian word since it contains four factors of length two.

Another property of Duval extensions will be introduced next.

3.2 Minimal Duval Extensions

The minimal Duval extension of a word w is the smallest prefix of a nontrivial Duval extension of w such that the prefix itself is a nontrivial Duval extension of w . The following theorem gives a rather surprising property of nontrivial Duval extensions. Its proof can be found in [5].

Theorem 15. *Let wu be a minimal Duval extension of w . Then u is a factor of w .*

Consider the following example.

Example 16. *Let $w = abaabbabaababb$ as in Example 6. Then*

$$w.u = abaabbabaababb.aaba$$

and

$$w.v = abaabbabaababb.abaaba$$

are both minimal Duval extensions of w , and u and v both occur in w .

3.3 Maximum Duval Extensions

The investigation of maximum Duval extensions has been motivated by the hope to estimate a precise bound on the relation between the length of a word w.r.t. the maximum length of its unbordered factors and its period. As we have seen with Corollary 10, there is the following upper bound: If $3\mu(w) - 2 \leq |w|$

then $\mu(w) = \partial(w)$. However, Example 3 is the best one known, showing that $\mu(w) < \partial(w)$ and $7/3\mu(w) - 4 = |w|$.

We have the following conjecture about the structure of maximum Duval extensions.

Conjecture 17. *Let $w = w'ab^k$ for some $k \geq 1$. If wu is a nontrivial Duval extension of w of length $2|w| - 2$, then b^k does not occur in w' .*

It has been shown in [4] that this conjecture would also imply Theorem 8.

4 Conclusions

We have recalled the problem of estimating the relationship between the length of a word and the maximum length of its unbordered factors. The final answer is still unknown. However, quite some progress has been made since the problem was raised in 1979. In particular the special case of Duval extensions raised attention and led to new results. For example, the long standing conjecture by Duval was just recently solved. However, open problems remain about the structure of Duval extensions and words that have no nontrivial Duval extension. Further research on those questions are likely to lead to a final answer to the general case.

Acknowledgements

The authors would like to thank Julien Cassaigne for pointing out that unbordered Sturmian words might be Lyndon words which led us to the effort of proving it.

References

- [1] R. Assous and M. Pouzet. Une caractérisation des mots périodiques. *Discrete Math.*, 25(1):1–5, 1979.
- [2] J.-P. Duval. Relationship between the period of a finite word and the length of its unbordered segments. *Discrete Math.*, 40(1):31–44, 1982.
- [3] A. Ehrenfeucht and D. M. Silberger. Periodicity and unbordered segments of words. *Discrete Math.*, 26(2):101–109, 1979.
- [4] T. Harju and D. Nowotka. Duval’s conjecture and Lyndon words. technical report 479, Turku Centre of Computer Science (TUCS), Turku, Finland, October 2002. submitted.
- [5] T. Harju and D. Nowotka. Minimal Duval extensions. technical report 520, Turku Centre of Computer Science (TUCS), Turku, Finland, April 2003. submitted.

- [6] T. Harju and D. Nowotka. Periodicity and unbordered words. technical report 523, Turku Centre of Computer Science (TUCS), Turku, Finland, April 2003. submitted.
- [7] M. Lothaire. *Combinatorics on Words*, volume 17 of *Encyclopedia of Mathematics*. Addison-Wesley, Reading, MA, 1983.
- [8] M. Lothaire. *Algebraic Combinatorics on Words*, volume 90 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, United Kingdom, 2002.
- [9] R. C. Lyndon. On Burnside's problem. *Trans. Amer. Math. Soc.*, 77:202–215, 1954.
- [10] F. Mignosi and L. Q. Zamboni. A note on a conjecture of Duval and Sturmian words. *Theor. Inform. Appl.*, 36(1):1–3, 2002.
- [11] M. Morse and G. A. Hedlund. Symbolic dynamics II: Sturmian trajectories. *Amer. J. Math.*, 61:1–42, 1940.