



The surface energy balance of early summer land-fast sea ice in Atka Bay, Antarctica

Stephan Paul (1), Clemens Drüe (1), Mario Hoppmann (2), Priska Hunkeler (2), Günther Heinemann (1), and Sascha Willmes (1)

(1) Environmental Meteorology, University of Trier, Behringstr. 21, 54296 Trier, Germany (paul@uni-trier.de), (2) Alfred Wegener Institute, Helmholtz Centre for Polar and Marine Research, Bussestr. 24, 27580 Bremerhaven, Germany

In-situ measurements of the land-fast sea ice energy balance are scarce. We present a data set that comprises eddy-covariance measurements of sensible and latent heat as well as measurements of the sea-ice temperature gradient, long-wave and short-wave radiation measurements over land-fast sea ice in Atka Bay, Antarctica. With this setup we are able to monitor all components of the sea-ice energy budget. Additionally, we also measured the turbulent flux of CO₂ over sea ice. This 37 day-long data set is evaluated for the transition period from austral winter to summer (November to December 2012) with regard to atmospheric stability and the general weather conditions. Results for the eddy-covariance measurements show an average sensible heat flux of $6.45 \pm 10.72 \text{ W/m}^2$ and a latent heat flux of $12.71 \pm 9.48 \text{ W/m}^2$ (with one standard deviation respectively) for low pressure/high wind-speed conditions. The average net radiation is $44.37 \pm 41.54 \text{ W/m}^2$ and for the CO₂ flux an average of $-3.35 \pm 3.37 \mu\text{mol/m}^2$ was measured. During high pressure/low wind-speed conditions an average of $-3.03 \pm 10.48 \text{ W/m}^2$ and $10.76 \pm 10.52 \text{ W/m}^2$ was recorded for the sensible and latent heat flux, while the average net radiation and the CO₂ flux are $35.63 \pm 56.70 \text{ W/m}^2$ and $-1.95 \pm 1.72 \mu\text{mol/m}^2$ respectively. The fast ice is therefore found as a sink of CO₂ for both situations.