

Intraseasonal variability in the tropical Atlantic: Observations vs. reduced gravity simulations

Robert Kopte, Peter Brandt, Richard J. Greatbatch, Martin Claus

1) Variability in the tropical Atlantic:

- ▶ Spectral peaks at annual and semi-annual, and 120-day periods associated with 4th, 2nd and 1st baroclinic modes
- ▶ Peaks correspond to resonant basin modes, composed of equatorial Kelvin and Rossby waves, as well as coastally trapped waves^[1]




Fig. 1: Periodogram of SLA in the tropical Atlantic and zonal velocity at 23°W, 0°N.

2) Reduced gravity simulations of the tropical Atlantic

- ▶ To study the intra-seasonal variability in particular, reduced gravity model (RGM) simulations are used:

$$u_t - fv = -g'_n \eta_x + \frac{\tau_x^x}{\rho_0}, \quad v_t + fu = -g'_n \eta_y + \frac{\tau_x^y}{\rho_0}, \quad \eta_t + H(u_x + v_y) = 0, \quad g'_n = \frac{c_n^2}{H}$$

- ▶ The model is run separately for the first five baroclinic modes ($c_1=2.47$ m/s, $c_2=1.32$ m/s, $c_3=0.94$ m/s, $c_4=0.74$ m/s, $c_5=0.57$ m/s), forced with interannually varying wind stress from NCEP (1990-2014)
- ▶ To allow for comparison of model and observations, the model output is fitted to AVISO sea level anomaly (SLA)^[2]




Fig. 2: Model domain with sponge layers at northern/southern boundaries, and area/sections of interest.

3) Comparison of reduced gravity simulations with AVISO SLA:

- ▶ Dominance of the annual and semi-annual cycle, well reproduced by the RGM (Fig. 3 and Fig.4)
- ▶ In AVISO, presence of continuous and recurrent eastward propagations^[3], with the intra-seasonal climatology representing ~25% of the seasonal cycle amplitude (Fig. 5)
- ▶ Only ~50% of the intra-seasonal signal amplitude is reproduced by the RGM, however the phase-lock of the propagations appears to be consistent

Fig. 3: Hovmöller plot of SLA along equatorial wave guide (green line in Fig. 2). Left: AVISO, right: Reconstruction from RGM modes 1-5.




Fig. 4: Climatologic (1995-2014) seasonal cycle of SLA along equatorial wave guide.

Fig. 5: Climatologic (1995-2014) seasonal cycle of band-passed (25-130d) SLA along equatorial wave guide.

4a) Basin mode of the 1st baroclinic mode




Fig. 6: Maps of amplitude (left) and phase (right) of 120-day harmonics fitted to SLA data. Top: AVISO, middle: RGM modes 1-5, bottom: RGM mode 1 only.

- ▶ Basin-wide structure of SLA associated with the 1st baroclinic mode with variability detectable as far as 10°S
- ▶ Consistent pattern in the RGM, although with considerable lower amplitude
- ▶ Mode 1 in the RGM simulations explains most of the structure seen in the “full” model

5) Summary and outlook

- ▶ Intra-seasonal SLA variability in the tropical Atlantic is essentially wind-driven, as it can be reproduced by reduced gravity simulations, although with weaker amplitudes
- ▶ Possible reasons for discrepancies to be tested:
 - ▶ Bad choice / spatial variability of phase speeds, which leads to the missing of resonance to a periodic forcing
 - ▶ Uncaptured (i.e. non-linear) effects of the North Equatorial Counter Current (NECC) on westwards propagating Rossby waves at ~4°N (Fig. 8)




Fig. 8: Climatologic (1995-2014), band-passed (25-130d) SLA along 4°N-Equ.-4°N (see Fig. 2). Top: AVISO, bottom: RGM modes 1-5.

4b) Equatorial wave analysis

- ▶ Averaged over the equatorial belt (5°S-5°N), zonal wavenumber-frequency diagrams of SLA exhibit spectral peaks near the theoretical dispersion curves of the first baroclinic mode equatorial Kelvin and Rossby waves
- ▶ In the RGM, total energy is lower, however there is considerable energy in the mixed Rossby-gravity wave range




Fig. 7: Mean (5°S-5°N, see horiz. black lines in Fig. 2) wave-number-frequency diagram. Left: AVISO, right: RGM (modes 1-5). The white lines represent the theoretical dispersion curves of the Kelvin wave and the first three meridional Rossby waves.

References

- [1] Cane MA, Moore DW (1981): A Note on Low-Frequency Equatorial Basin Modes. *J Phys Oceanogr* 11:1578-1584 doi:10.1175/1520-0485
- [2] The altimeter products were produced by Ssalto/Duacs and distributed by Aviso, with support from Cnes (<http://www.aviso.altimetry.fr/duacs/>)
- [3] Polo I, Lazar A, Rodriguez-Fonseca B, Arnault S (2008): Oceanic Kelvin waves and tropical Atlantic intraseasonal variability: 1. Kelvin wave characterization. *J Geophys Res-Oceans* 113:18 doi:10.1029/2007jco04495