

INTERACTION OF TEMPERATURE AND CO₂ ENRICHMENT ON SOYBEAN: PHOTOSYNTHESIS AND SEED YIELD

N. SIONIT¹, B. R. STRAIN, and E. P. FLINT

¹Department of Botany, Duke University, Durham, North Carolina 27706, U.S.A.
Received 24 July 1986, accepted 9 Jan. 1987.

SIONIT, N., STRAIN, B. R. AND FLINT, E. P. 1987. Interaction of temperature and CO₂ enrichment on soybean: Photosynthesis and seed yield. *Can. J. Plant Sci.* **67**: 629–636.

Seed yield and photosynthetic responses of soybean (*Glycine max* L. Merr. 'Ransom') were studied in growth chambers at day/night temperatures of 18/12, 22/16, and 26/20°C and atmospheric CO₂ concentrations of 350, 675 and 1000 µL L⁻¹. No seeds were produced at 18/12°C within any of the CO₂ concentrations. Numbers of pods and seeds increased with increasing temperature and CO₂ levels. Carbon dioxide enrichment increased seed yield of soybean grown at moderately cool temperatures. This increase was associated with an increase in net photosynthetic rate. Leaf photosynthesis in response to CO₂ enrichment increased more at 22/16°C than at 26/20°C. Increases in temperature and CO₂ levels enhanced total growth of plants but hastened senescence of leaves. The extended photosynthetic capacity at cool temperatures did not result in allocating more dry matter to developing pods. CO₂ enrichment at 26/20°C resulted in greater seed yield increases than CO₂ enrichment at lower temperatures.

Key words: Soybean yield, low temperature, CO₂ × temperature

[Effets d'une interaction entre la température et la teneur en CO₂ sur la photosynthèse et le rendement grainier du soja.]

Titre abrégé: Effets du CO₂ et de la température sur le soja.

Nous avons étudié les variations du rendement grainier et de la photosynthèse du soja (*Glycine max* L. Merr. 'Ransom') dans des chambres de croissance à des températures (jour/nuit) de 18/12, 22/16 et 26/20°C et à des teneurs en CO₂ de l'atmosphère de 350, 675 et 1 000 µL L⁻¹. Aucune semence n'a été produite à 18/12°C aux teneurs en CO₂ testées. Le nombre de gousses et de semences a augmenté en même temps que la température et les teneurs en CO₂. L'enrichissement en CO₂ a provoqué une hausse du rendement grainier du soja cultivé à des températures modérément fraîches. Cette hausse est attribuable à une augmentation du taux net de photosynthèse. L'augmentation de la photosynthèse des feuilles avec l'enrichissement en CO₂ a été plus rapide à 22/16°C qu'à 26/20°C. Les hausses de la température et de la teneur en CO₂ ont favorisé la croissance des plantes mais hâté la sénescence des feuilles. La capacité accrue de photosynthèse aux températures fraîches n'a pas provoqué d'augmentation de la proportion de la matière sèche contenue dans les gousses en développement. L'enrichissement en CO₂ à 26/20°C a donné une augmentation plus importante du rendement grainier que l'enrichissement en CO₂ aux températures plus basses.

Mots clés: Rendement du soja, basse température, CO₂ × température

¹Present address (N.S.): Systems Ecology Research Group, San Diego State University, San Diego, California 92182, U.S.A.

Productivity of seed crops is a function of total photosynthate produced by the source and proportion of the assimilate allocated to reproductive sinks (Kramer 1981). Variability in yield among crop species and cultivars of the same species is due to genetic potential and environmental conditions (Boyer 1982). Environmental factors have been shown to affect the processes of both photosynthesis and dry matter allocation in plants. For example, CO₂ concentration in the atmosphere (Kimball 1983; Strain 1985) and temperature regime (Patterson and Flint 1979) are two important factors that regulate gas exchange and growth rates in plants. Net photosynthesis increases while stomatal conductance decreases in leaves with increases in atmospheric CO₂ concentration (Allen et al. 1981; Strain and Armentano 1982; Rogers et al. 1983; Sionit et al. 1984; Jones et al. 1985a,b; Valle et al. 1985). Thomas and Raper (1978), and Seddigh and Jolliff (1984) have shown that temperature regime has a significant effect on net CO₂ exchange rate of leaves, carbohydrate allocation, and yield of soybean.

Most previous investigations on the effects of atmospheric CO₂ enrichment have given little attention to the possible interactions between CO₂ and temperature. The vigor and productivity of okra (*Abelmoschus esculentus*) at low temperatures, however, has been shown (Sionit et al. 1981) to be greater when atmospheric CO₂ is enriched. The present study was undertaken to analyze the combined effects of increasing atmospheric CO₂ concentration and temperature regimes on photosynthesis and production of soybean.

MATERIALS AND METHODS

Soybean (*Glycine max* (L.) Merr. 'Ransom') seeds were germinated in a 1:1:1 (by volume) mixture of gravel:vermiculite:Turface (Jaeger et al. 1981) in 3.5-L plastic pots and grown in controlled environment chambers of the Duke University Phytotron (Kramer et al. 1970). Ransom is a determinate, maturity group VII, cultivar that is widely planted in the southeastern U.S.A. There was one plant in each pot and the pots were watered to the drip point three times every day with

modified half-strength Hoagland's solution (Downs and Hellmers 1978). The photoperiod was 12 h. For the 3-wk flower induction period from 21 to 41 d after planting the photoperiod was 9 h. A combination of fluorescent and incandescent lamps at an input wattage ratio of 15:4 provided a photosynthetic photon flux density of $550 \pm 50 \mu\text{mol m}^{-2} \text{s}^{-1}$, as measured with a LiCor Quantum sensor at the top of the plant canopy in the chambers. The relative humidity was 70% during the day and close to saturation at night. The day/night temperature regimes were 18/12, 22/16, and 26/20°C. Carbon dioxide was injected into the chambers automatically and its concentration monitored continuously by an infrared gas analyzer (Hellmers and Giles 1979). The injection system provided CO₂ concentrations of 350 ± 40 , 675 ± 30 , or $1000 \pm 45 \mu\text{L L}^{-1}$ in three different chambers at each temperature regime.

Eight plants were harvested at each of the three CO₂ and three temperature environments at 20, 40, 67 and 115 (late senescence to maturity) days after planting for dry weight and leaf area determinations. Numbers of pods and seeds were determined and dry weights of seed per plant and per seed were obtained after drying at 65°C to constant weight. Net assimilation rate (NAR), which is the amount of dry matter produced per unit leaf area per unit time ($\text{g m}^{-2} \text{d}^{-1}$), was calculated for the harvest intervals of 20–40 and 40–67 d. Net photosynthetic rate and stomatal conductance measurements were made in the respective chambers on four plants on the middle leaflet of the third or fourth trifoliolate leaves from the top of the main stem with a steady state micucuvette system developed by Bingham et al. (1980). Vapor pressure deficit in the cuvette was controlled and maintained at $8.0 \pm 0.15 \text{ g m}^{-3}$ in all measurements. The plants were at early- to late-flowering stage, stages R₁ to R₄ (Fehr and Caviness 1977), during the measurements. No photosynthetic measurement was made at 18/12°C because of technical difficulties. A two-way analysis of variance was used for CO₂ and temperature to determine differences among treatments.

RESULTS AND DISCUSSION

The number of pods per plant increased with increasing day/night temperature regime from 18/12°C to 26/20°C and increased CO₂ concentration from 350 to 675 $\mu\text{L L}^{-1}$ (Table 1). Further increase in CO₂ concentration to 1000 $\mu\text{L L}^{-1}$ produced little or no

Table 1. Pods per plant of soybean grown at three CO₂ concentrations and three day/night temperature regimes

CO ₂ Concentration ($\mu\text{L L}^{-1}$)	Day/night temperature (°C)		
	18/12	22/16	26/20
350	43.9	86.4	121.6
675	55.6	95.4	170.0
1000	50.1	115.9	168.9
<i>Source of variation</i>			
CO ₂		**	
Temp.		**	
CO ₂ × Temp.		*	

*, ** $P < 0.05$ and $P < 0.01$, respectively.

additional increase in number of pods at all temperatures. Greater increases in number of pods per plant were observed in response to CO₂ enrichment from 350 $\mu\text{L L}^{-1}$ to 675 $\mu\text{L L}^{-1}$ at high temperatures than at low temperatures. The temperature × CO₂ interaction for pod production was statistically significant (Table 1). Seddigh and Jolliff (1984) reported that pod production in a field-grown indeterminate soybean cultivar was increased during the early reproductive stage by warmer night temperatures (24°C as compared to 16 or 10°C), but due to differences in rates of pod abortion the final number of pods at maturity was not affected by warm night temperatures. Thomas and Raper (1978) recorded a two-fold increase in pod number of a phytotron-grown determinate soybean cultivar by increasing night temperature from 14°C to 18°C at day temperature of 22°C. In another study, Thomas and Raper (1976) reported that increasing pod production at 30/26°C was dependent upon the number of consecutive short-day photoperiods and morphological development stage at floral induction. Carbon dioxide treatment complicated the temperature effect on pod production because it produced differences in the size of plants at flower induction (Sionit et al. 1987). This plus the direct effect of CO₂ enrichment on subsequent growth (Sionit et al. 1987) resulted in more branches and more nodes on the main stem, causing a greater potential for pod production.

All pods produced at 18/12°C in each of the

CO₂ treatments were small, parthenocarpic and abnormal in shape. No seeds were developed in these pods; however, CO₂ enrichment up to 675 $\mu\text{L L}^{-1}$ increased the number of pods. Such a cold-induced pod abnormality in soybean has been observed by Thomas and Raper (1981) at 18/18°C and 22/14°C and by Lawn and Hume (1985) at 20/12°C and at 20/8°C. The physiological basis for production of abnormal pods at cool temperatures has partly been investigated. Thomas and Raper (1981) proposed that the events occurring during floral development were responsible for abnormal pods because successive development after floral initiation seemed to have "increasingly stricter temperature requirements". Lawn and Hume (1985) found that absence of normal pod set at low night temperatures in several soybean lines of temperate and tropical origins was related to reduction in pollen formation and reduced pollen vigor resulting in curled and distorted pollen tubes.

Sionit et al. (1981), in a phytotron study, observed that CO₂ enrichment at suboptimal temperature of 20/14°C compensated for the adverse effect of cool temperature on pod number for okra.

Increasing CO₂ concentration from 350 to 675 $\mu\text{L L}^{-1}$ at high temperatures caused a greater percentage increase in seed yield than it did at 22/16°C (Fig. 1). The increase in total seed yield in response to increasing CO₂ concentration and temperature regime was mainly associated with the increased numbers of pods (Table 1) and seeds (Fig. 1) and not with the weight of individual seeds (Fig. 1). Seddigh and Jolliff (1984) observed that soybean grown under ambient day temperature in the field produced smaller seeds at night temperatures of 10 or 16 than at 24°C. These investigators concluded that production of assimilates was unaffected by cool nights but growth rate of seeds was limited by the direct effect of low night temperature. The growth rate of individual seeds appears to be insensitive to a range of variation in the supply of assimilate (Egli and Leggett 1976). Although we did not measure rates of trans-




Fig. 1. Seed yield, number of seeds/plant and seed weight of Ransom soybean at three CO₂ concentrations (L=350, M=675 and H=1000 $\mu\text{L L}^{-1}$) and three day/night temperature regimes (18/12, 22/16 and 26/20 °C).

Table 2. Mean and standard error for stomatal conductances and photosynthetic rates of leaves of soybean grown at three CO₂ concentrations and two day/night temperature regimes

CO ₂ concentration ($\mu\text{L L}^{-1}$)	Stomatal conductance (cm s^{-1})		Net photosynthetic rate ($\mu\text{mol m}^{-2} \text{s}^{-1}$)	
	Day/night temperature, °C			
	22/16	26/20	22/16	26/20
350	0.56+0.04	0.26+0.03	14.3±0.9	22.8±0.4
675	0.25+0.07	0.17+0.02	17.7±1.1	26.1±2.0
1000	0.27+0.02	0.14+0.02	19.8±1.2	21.8±1.1

location of assimilates to the seeds, data of Fig. 1 suggest that production and translocation of assimilates to individual seeds probably were not limited at 22/16°C at any CO₂ concentration. It might be that 22/16°C temperature regime is optimum, or close to optimum, for maximum seed size as indicated by no statistical differences obtained in seed size between 22/16°C and 26/20°C treatments. Atmospheric CO₂ enrichment had little or no effect on seed size at these temperature regimes. Therefore, the differences in seed yield per plant at different atmospheric CO₂ and temperature levels were due to differences in number of pods per plant and not due to the weight of individual seeds. Numbers of seeds per pod were unaffected by the CO₂ and temperature treatments. Sionit et al. (1987) reported increasing growth of different vegetative plant parts in response to increasing levels of atmospheric CO₂ and temperature. It appears that most of the yield differences obtained in the present experiment were due to morphological differences (Sionit et al. 1987) in growth of the plants. The overall influence of temperature and CO₂ treatments, therefore, was such that the sink size was probably partly adjusted in response to cool temperature and CO₂ enrichment. Thus, in most instances, when plants were grown in 1000 $\mu\text{L L}^{-1}$ CO₂ at 22/16°C the high CO₂ compensated for the deleterious effect of low temperature in producing pods and seed mass.

As expected, leaves of soybeans at high CO₂ concentrations or high temperatures had lower stomatal conductance than leaves of plants grown at low CO₂ or low temperature

levels (Table 2). While stomatal conductance decreased, leaf photosynthetic rates increased with increasing CO₂ and temperature levels except for 1000 $\mu\text{L L}^{-1}$ CO₂ at 26/20°C (Table 2). Carbohydrate accumulation in plant leaves has been suggested to be a cause of decreased photosynthesis in 1000 $\mu\text{L L}^{-1}$ compared to lower CO₂ concentrations (Clough et al. 1981; Wulff and Strain 1982). The decrease in photosynthesis in response to 1000 $\mu\text{L L}^{-1}$ CO₂ was not observed in our experiment at the lower temperature regime (22/16°C). This may partially be due to a reduced rate of production of assimilates at lower temperatures.

Temperature regime and CO₂ enrichment influenced net assimilation rate (NAR), but the interaction of these treatments did not produce statistically significant effects (Table 3). The NAR was always higher at high than at low CO₂ concentrations. Soybean plants grown at high CO₂ levels were larger, produced more branches and greater leaf area than plants grown at lower CO₂ levels (Sionit et al. 1987). Self shading did not influence the NAR at high CO₂ levels in our study as has been demonstrated by Flint and Patterson (1983). Ransom soybean had higher NAR values at 22/16°C than at either 18/12°C or 26/20°C. NAR reflects a long-term assimilation by the whole plant between two sampling dates but net photosynthesis is a measure of CO₂ uptake by a single, unshaded, and relatively young leaf. Therefore, NAR (Table 3) is dependent upon the phenology and size of the plants, whereas leaf photosynthesis (Table 2) is related to stomatal conductances of the measured leaves. These

Table 3. Net assimilation rates ($\text{g m}^{-2} \text{d}^{-1}$) of soybean during the 20- to 40-d and 40- to 67-d harvest intervals at three CO_2 concentrations and three day/night temperature regimes

CO ₂ concentration ($\mu\text{L L}^{-1}$)	20-40 day			40-67 day		
	Day/night temperature °C					
	18/12	22/16	26/20	18/12	22/16	26/20
350	7.7	9.3	6.8	6.3	10.7	6.6
675	10.8	11.7	8.9	8.4	13.7	9.5
1000	11.9	13.3	10.8	9.0	14.6	10.0
<i>Source of variation</i>	<i>20-40 day</i>			<i>40-67 day</i>		
CO ₂	**			**		
Temp.	**			**		
CO ₂ × Temp.	NS			NS		

*, ** $P < 0.05$ and $P < 0.01$, respectively.

factors may have contributed to different trends of changes observed in NAR and net photosynthesis in response to increasing temperature from 22/16°C to 26/20°C.

Although seed yield of Ransom soybean at 22/16°C and 1000 $\mu\text{L L}^{-1}$ CO₂ was equal to that at 26/20°C and ambient CO₂ (Fig. 1), the photosynthetic rates were different (Table 2). CO₂ enrichment increased net photosynthesis at low temperature but did not completely compensate for the low-temperature effects. Increasing the temperature from 22/16°C to 26/20°C increased net photosynthesis by 59, 46 and 10% at 350, 675 and 1000 $\mu\text{L L}^{-1}$ CO₂, respectively. Increasing CO₂ concentration from 350 to 675 $\mu\text{L L}^{-1}$ and from 675 to 1000 $\mu\text{L L}^{-1}$ increased net photosynthesis by 23.8 and 11.4%, respectively, at 22/16°C and by 14.1% and -16.5%, respectively, at 26/20°C. The data suggest that leaf photosynthesis in soybean benefited from CO₂ enrichment more at a moderately low temperature regime than at a high temperature regime.

The increase in temperature and CO₂ hastened the time of flowering, appearance of pods, and senescence of leaves. Plants grown at the combination of high CO₂ and high temperatures approached the full seed development stage about 19 d earlier than those grown at low levels of temperature and ambient CO₂. Although the low temperature regime at all CO₂ levels prolonged the

duration of efficient photosynthesis of plants by postponing senescence of the leaves, the extended photosynthetic capacity did not result in allocating more dry matter to developing pods. This may have been due to significantly lower rates of net photosynthesis and of subsequent translocation of substrates at low temperatures.

The data show that amelioration of low-temperature effects on soybean yield by atmospheric carbon dioxide enrichment is limited to moderately low temperatures and at a CO₂ level of 1000 $\mu\text{L L}^{-1}$. The projected global increase in carbon dioxide concentration for the next 50 yr and the resultant climatic warming are most likely to increase soybean yields in areas which will have moderately warm to warm climates.

ACKNOWLEDGMENTS

This study was supported by NSF Grants Nos. DEB 80-21312, BSR 82-15533, and BSR 83-14925. Helpful suggestions during the course of the experiment and critical reading of the manuscript by Drs. P. J. Kramer, D. T. Patterson and C. David Raper, Jr. are greatly appreciated. The authors are grateful to Drs. G. E. Bingham and J. H. Shinn for use of their steady state minicuvette system. The technical assistance of C. H. Jeager and the skillful assistance of the phytotron staff are acknowledged.

Allen, L. H., Jr., Boote, K. J., Jones, J. W., Mishoe, J. W., Jones, P. K., Valle, R., Vu, C. V. and Campbell, W. 1981. Response of vegetation

- to carbon dioxide: Effects of increased carbon dioxide on photosynthesis and agricultural productivity of soybeans. Report 003. U.S. Dept. of Energy, Carbon Dioxide Res. Div.; and U.S. Dept. of Agriculture, Res. Ser. 131 pp.
- Bingham, G. E., Coyne, P. I., Kennedy, R. B. and Jackson, W. L. 1980.** Design and fabrication of a portable minicuvette system for measuring leaf photosynthesis and stomatal conductance under controlled conditions. UCRL-52895. Lawrence Livermore Laboratory, Livermore, Calif.
- Boyer, J. S. 1982.** Plant productivity and environment. *Science* **218**: 443–448.
- Clough, J. M., Peet, M. M. and Kramer, P. J. 1981.** Effects of high atmospheric CO₂ and sink size on rates of photosynthesis of a soybean cultivar. *Plant Physiol.* **67**: 1007–1010.
- Downs, R. J. and Hellmers, H. 1978.** Controlled climate and plant research. Technical Note No. 148. World Meteorological Organization. Geneva, Switzerland.
- Egli, D. B. and Leggett, J. E. 1976.** Rate of dry matter accumulation in soybean seeds with varying source-sink ratios. *Agron. J.* **68**: 371–374.
- Fehr, W. R. and Caviness, C. E. 1977.** Stages of soybean development. Special Rep. 80, Iowa State University, Ames, Iowa.
- Flint, E. P. and Patterson, D. T. 1983.** Interference and temperature effects on growth in soybean (*Glycine max*) and associated C₃ and C₄ weeds. *Weed Sci.* **31**: 193–199.
- Hellmers, H. and Giles, L. J. 1979.** Carbon dioxide: Critique I. Pages 229–234. *In* T. W. Tibbitts and T. T. Kozlowski, ed. Controlled environment guidelines for plant research. Academic Press, New York.
- Jaeger, C. H., Hellmers, H. and Teare, I. D. 1981.** An improved medium for nutriculture incorporating arcillite. *Hortic. Sci.* **6**: 176–177.
- Jones, P., Allen, L. H., Jr. and Jones, J. W. 1985a.** Responses of soybean canopy photosynthesis and transpiration to whole-day temperature changes in different CO₂ environments. *Agron. J.* **77**: 242–249.
- Jones, P., Allen, L. H., Jr., Jones, J. W. and Valle, R. 1985b.** Photosynthesis and transpiration responses of soybean canopies to short- and long-term CO₂ treatments. *Agron. J.* **77**: 119–126.
- Kimball, B. A. 1983.** Carbon dioxide and agricultural yield: An assemblage and analysis of 430 prior observations. *Agron. J.* **75**: 779–788.
- Kramer, P. J. 1981.** Carbon dioxide concentration, photosynthesis, and dry matter production. *BioScience* **31**: 29–33.
- Kramer, P. J., Hellmers, H. and Downs, R. J. 1970.** SEPEL: New phytotrons for environmental research. *BioScience* **20**: 1201–1208.
- Lawn, R. J. and Hume, D. J. 1985.** Response of tropical and temperate soybean genotypes to temperature during early reproductive growth. *Crop Sci.* **25**: 137–142.
- Patterson, D. T. and Flint, E. P. 1979.** Effects of simulated field temperatures and chilling on itchgrass (*Rottboellia exaltata*), corn (*Zea mays*), and soybean (*Glycine max*). *Weed Sci.* **27**: 645–650.
- Rogers, H. H., Bingham, G. E., Cure, J. D., Smith, J. M. and Surano, K. A. 1983.** Responses of selected plant species to elevated carbon dioxide in the field. *J. Environ. Qual.* **12**: 569–574.
- Seddigh, M. and Jolliff, G. D. 1984.** Night temperature effects on morphology, phenology, yield and yield components of indeterminate field-grown soybean. *Agron. J.* **76**: 824–828.
- Sionit, N., Strain, B. R. and Beckford, H. A. 1981.** Environmental control on the growth and yield of okra. I. Effects of temperature and of CO₂ enrichment at cool temperature. *Crop Sci.* **21**: 885–888.
- Sionit, N., Rogers, H. H., Bingham, G. E. and Strain, B. R. 1984.** Photosynthesis and stomatal conductance with CO₂ enrichment of container- and field-grown soybeans. *Agron. J.* **76**: 447–451.
- Sionit, N., Strain, B. R. and Flint, E. P. 1987.** Interaction of temperature and CO₂ enrichment on soybean: Growth and dry matter partitioning. *Can. J. Plant Sci.* **67**: 59–67.
- Strain, B. R. 1985.** Physiological and ecological controls on carbon sequestering in ecosystems. *Biogeochemistry* **1**: 219–232.
- Strain, B. R. and Armentano, T. V. 1982.** Responses of “Unmanaged” Ecosystems, 013, Vol. II, Part 12. Environmental and Societal Consequences of a Possible CO₂-Induced Climate Change. Prepared for U.S. Dept. of Energy, DOE/EV/10019-12. 104 pp.
- Thomas, J. F. and Raper, C. D., Jr. 1976.** Photoperiod control of seed filling for soybeans. *Crop Sci.* **16**: 667–672.
- Thomas, J. F. and Raper, C. D., Jr. 1978.** Effect of day and night temperatures during floral induction on morphology of soybeans. *Agron. J.* **70**: 893–898.
- Thomas, J. F. and Raper, C. D., Jr. 1981.** Day and night temperature influence on carpel initiation and growth in soybeans. *Bot. Gaz.* **142**: 183–187.
- Valle, R., Mishoe, J. W., Campbell, W. J., Jones, J. W. and Allen, L. H., Jr. 1985.**

Photosynthetic responses of 'Bragg' soybean leaves adapted to different CO₂ environments. *Crop Sci.* **25**: 333-339.

Wulff, R. D. and Strain, B. R. 1982. Effect of

CO₂ enrichment on growth and photosynthesis of *Desmodium paniculatum*. *Can. J. Bot.* **60**: 1084-1091.