

## 16. SILICOFLAGELLATES, EBRIDIANS, AND ACTINISCIDIANS FROM PLIOCENE AND QUATERNARY SEDIMENTS OFF SOUTHERN CHILE, ODP LEG 141<sup>1</sup>

Sigurd Locker<sup>2</sup>

### ABSTRACT

Three groups of siliceous flagellates are discussed from Sites 860 and 861 in the southeastern South Pacific Ocean off southern Chile. Three silicoflagellate zones and two ebridian/actiniscidian zones are recognized in upper Pliocene to Quaternary sediments. Repeated occurrences of *Dictyocha messanensis messanensis* demonstrate that the cool subpolar water masses of the Peru Current were mixed with or replaced by warm transitional water masses of the central South Pacific several times during late Pliocene to middle Pleistocene times. Continuous occurrences of *Dictyocha messanensis aculeata* in Zone NN21 sediments reflect an enhanced influence of transitional surface-water also during late Pleistocene and Holocene times. These Pliocene and Quaternary warming episodes may have been caused by strong El Niño activities, which resulted in a southward shift of the transitional water belt and an expansion of warmer surface-water over the study area.

### INTRODUCTION

During Leg 141 of the Ocean Drilling Program (ODP), 13 holes were drilled at five sites between about 45°40'S and 47°00'S, off southern Chile (Fig. 1). This is the region where the active oceanic spreading center of the Chile Ridge is being subducted under the continental plate of South America. The main objective of drilling was to investigate the geologic and tectonic processes that are associated with the subduction of oceanic crust and sediment material under the continent.

To determine structural and depositional characters of the collision zone, three sites (Sites 859, 860, and 861) were arranged along a transect across the accretionary wedge and the forearc basin just north of the present-day Chile Triple Junction. The sediment sequences retrieved consist of upper Pliocene to Quaternary predominantly clayey to silty sediments of varying consolidation. These sediments contain diversified assemblages of calcareous and siliceous microfossils, of both planktic and benthic origin.

This study deals with three groups of planktic microorganisms possessing siliceous skeletons (i.e., silicoflagellates, ebridians, and actiniscidians). Neglecting specific taxonomic affinities, these groups may be summarized as siliceous flagellates. In general, representatives of siliceous flagellates were minor constituents of the plankton communities during Pliocene and Quaternary times. But, nevertheless, they provide important stratigraphic and paleoceanographic information. Because sediments at Sites 860 and 861 yielded plankton assemblages that were more diversified and better preserved, investigations were restricted to those sites. Stratigraphic results are summarized in Figure 2 and Tables 1, 2, and 3, and paleoceanographic interpretations are displayed in Figures 3 and 4 and Table 2.

### METHODS

To concentrate skeletons of siliceous flagellates, sediment samples were processed successively with hydrogen peroxide, hydrochloric acid, and sodium pyrophosphate. After the use of acid and lye, respectively, samples were washed three times. Decalcified residues were strewn on cover glasses 22 × 22 mm in size and then embedded in Canada balsam.

Generally, eight traverses across a cover glass (21%) were counted with a magnification of ×390 and tabulated for species abundances. But, when searching for stratigraphically indicative species, more traverses were scanned. The following abundance classes were used:

- B = barren
- T = traces (specimens out of count)
- R = rare (1–5 specimens)
- F = few (6–15 specimens)
- C = common (16–50 specimens)
- A = abundant (>50 specimens)
- + = reworked specimens

Total abundances of silicoflagellates and ebridians/actiniscidians were classified as described above. Because siliceous flagellates were generally minor constituents of the protophyte assemblages that consisted mainly of diatoms, diatom abundances were estimated, too, and noted as an indicator for expected flagellate abundances. The following abundance classes were adopted:

- B = barren
- T = traces, single valves and/or remnants of corrosion
- F = few, minor content of valves and fragments
- C = common, major content of valves and fragments

In most studies, the preservational state of siliceous flagellates remains undescribed, as it is difficult to determine the original extent of fragmentation and corrosion. The fragile skeletons of silicoflagellates may be fragmented during preparation, and corrosion stages of ebridians and actiniscidians may be confused with juvenile skeletons. Because the slides of this study were prepared from suspended material, the state of preservation can be indicated in a general manner as follows. In samples containing complete skeletons, preservation is moderate; in samples containing only fragments, preservation is poor. The star-shaped skeletons of actiniscidians are most resistant and disappear last.

### BIOSTRATIGRAPHIC ZONATIONS

Plankton biostratigraphy of upper Neogene to Quaternary sediments is strongly influenced by paleoceanography, especially by ecological adaptations of the former plankton communities to certain water masses and current systems. The biozonations used for Leg 141 obviously represent high-latitude subdivisions, which may be recog-

<sup>1</sup> Lewis, S.D., Behrmann, J.H., Musgrave, R.J., and Cande, S.C. (Eds.), 1995. *Proc. ODP, Sci. Results*, 141: College Station, TX (Ocean Drilling Program).

<sup>2</sup> GEOMAR Research Center for Marine Geosciences, Wischhofstrasse 1-3, D-24148 Kiel, Federal Republic of Germany.




Figure 1. Location of ODP Leg 141 sites in the southeastern South Pacific and position of the subduction zone off southern Chile. Bathymetric contours in meters.

Age	NN zones	Silicoflagellate zones	Ebridian/actiniscidian zones	Datum indicators
Quaternary	21	<i>D. messanensis aculeata</i>		FCO <i>D. messanensis aculeata</i>
	20	<i>Distephanus speculum</i>	<i>Actiniscus pentasterias</i>	
	19			
Pliocene	16-18	<i>Distephanus aculeatus</i>	<i>Ammodochium serotinum</i>	LO <i>D. aculeatus</i> LO <i>A. serotinum</i>
		Unzoned	Unzoned	
	15	Barren	Barren	

Figure 2. Silicoflagellate and ebridian/actiniscidian zonations used for Sites 860 and 861, FOs and LOs of diagnostic species and correlation to standard nannofossil zones (Spiegler and Müller, this volume).




Figure 3. Distribution of surface-water masses in the southeastern South Pacific, based on present-day annual mean potential temperatures (according to Levitus, 1982). Location of Site 861 and position of Sites 594 and 688 where *Dictyocha messanensis aculeata* was found in Zone NN19–NN21 sediments.

nized in both hemispheres, but which differ distinctly from subdivisions described for middle and low latitudes.

The stratigraphic sequences cored at Sites 860 and 861 can be divided into three silicoflagellate zones and two ebridian/actiniscidian zones (Fig. 2). Definitions are given below. The correlations to nannoplankton zones are based on the data of Spiegler and Müller (this volume). The Pliocene/Pleistocene boundary was placed at the first occurrence of *Gephyrocapsa oceanica* (Spiegler and Müller, this volume), which corresponds to the base of Zone CN14 (Okada and Bukry, 1980) or an uplifted Zone NN18/NN19 boundary (Martini and Müller, 1986). This level has been dated at 1.6 Ma by Berggren et al. (1985). Paleomagnetic data could not be used for comparison, because of the strongly overprinted paleomagnetic signals.

### Silicoflagellates

#### *Distephanus aculeatus* Zone

**Definition:** Interval from the last occurrence (LO) of *Mesocena diodon* to the LO of *Distephanus aculeatus* (Bukry and Foster, 1973; base redefined by Locker and Martini, 1989). Lower to uppermost Pliocene.

**Occurrence:** Found in Holes 860B and 861C. The LO of *M. diodon* was not discovered in both holes. The LO of *D. aculeatus* is well developed in Hole 861C, but poorly preserved in Hole 860B. In Hole 861C, the LO of *D. aculeatus* probably lies in the uppermost part of Zone NN18.

**Remarks:** The last appearance datum (LAD) of *M. diodon* is at or just above the Miocene/Pliocene boundary, which may correspond to the top of Zone NN12. The LAD of *D. aculeatus* (synonym *Dictyocha boliviensis*) approaches the Pliocene/Pleistocene boundary or the top of Zone NN18. This has been found in the eastern Pacific Ocean at Site 157 (Bukry and Foster, 1973), and in the Southern Ocean at Sites 266, 269, and 274 (Ciesielski, 1975). At other high-latitude sites, the LAD of *D. aculeatus* may be earlier, and then, some equivalent *Distephanus speculum* zones have been used by authors.

#### *Distephanus speculum* Zone

**Definition:** Interval from the LO of *Distephanus aculeatus* to the first consistent occurrence (FCO) of *Dictyocha messanensis aculeata* (Ciesielski, 1975; top modified in this study). Uppermost Pliocene to Pleistocene.

**Occurrence:** Found in Holes 860B and 861C.

**Remarks:** Depending on the LO of *D. aculeatus* and the first occurrence (FO) and FCO of *D. messanensis aculeata*, respectively, the content of this zone may vary regionally.

#### *Dictyocha messanensis aculeata* Zone

**Definition:** Interval above the FCO of *Dictyocha messanensis aculeata* (Dumitrica, 1973; Bukry and Foster, 1973; base modified in this study). Pleistocene to Holocene.




Figure 4. Generalized stratigraphic distribution of *Distephanus speculum* and *Dictyocha messanensis* at Site 861 and correlation to standard nannofossil zones (Spiegler and Müller, this volume). Abbreviations: Ds = only *D. speculum* present, Ds > Da = minor occurrence of *D. messanensis*, Ds < Da = major occurrence of *D. messanensis*.

**Occurrence:** Found in Holes 860B and 861C. The FCO of *D. messanensis aculeata* commences in Zone NN21 of both holes.

**Remarks:** Because specimens of *Mesocena quadrangula* were not found in the material, the original zonal definition had to be modified for Leg 141 use. Here, the base has been placed at the FCO of *D. messanensis aculeata*, which corresponds to a paleoceanographic level above its first appearance datum (FAD).

The FAD of *D. messanensis aculeata* is in the middle Pleistocene or within Zone NN19. This has been found, for instance, in the northwestern and northern Pacific Ocean at Sites 299 and 310 (Ling, 1975; Bukry, 1978), in the eastern Pacific at Sites 157, 425, and 688 (Bukry and Foster, 1973; Bukry, 1980; Martini, 1990), in the southwestern Pacific at Sites 206, 278, and 594 (Dumitrica, 1973; Perch-Nielsen, 1975; Locker and Martini, 1986b), in the northeastern and northern Atlantic at Sites 397 and 412 (Bukry, 1979a, 1979b).

Table 1. Stratigraphic distribution of siliceous flagellates in upper Pliocene to Quaternary sediments of Hole 860B and correlation to standard nanno-fossil zones (Spiegler and Müller, this volume).

Nanno-plankton zone	Hole 860B, core, section, interval (cm)	Depth (mbsf)	Slide area (%)	Diatom abundance							
					<i>Da. messanensis</i> , total	<i>Da. mess. aculeata</i>	<i>Da. mess. messanensis</i>	<i>Da. mess. stapedia</i>	<i>Da. perlaevis</i>	<i>Da. spp.</i>	<i>Ds. aculeatus</i>
NN21	1H-1, 90-91	00.90	63	C	R	T	R	.	.	1+	.
	2H-1, 40-42	01.80	21	C	F	R	R	.	.	.	.
	2H-2, 19-21	03.09	21	T	Barren						
?	2H-4, 29-31	06.19	42	C	F	.	F	.	.	.	.
	2H-5, 18-20	07.58	42	F	.	.	.	.	.	.	.
	2H-7, 19-21	10.59	63	C	R	R	.	.	.	.	.
NN20	3H-3, 100-101	14.90	42	C	.	.	.	.	.	.	.
NN19	4H-3, 100-101	24.40	21	T	.	.	.	.	.	.	.
	5H-3, 37-38	33.27	21	T	Fragments						
	6H-3, 94-95	43.34	21	T	Barren						
	7H-4, 60-61	54.00	21	T	Fragments						
	12X-3, 100-101	91.70	42	T	.	.	.	.	.	.	.
	14X-2, 48-49	101.08	21	T	Barren						
	15X-3, 100-101	111.20	42	F	R	.	R	.	.	1+	.
NN16-18	16X-2, 68-69	118.98	21	T	Fragments						
	17X-CC	136.10	21	T	Barren						
	19X-3, 100-101	141.10	63	C	R	.	R	R	T	1,2+	.
	20X-1, 127-128	147.07	63	F	.	.	.	.	R	1+	.
	21X-1, 72-73	156.22	42	F	.	.	.	.	.	.	T
	22X-2, 103-104	167.63	42	C	.	.	.	.	.	.	.
	24X-2, 25-26 to	186.15	10	T	Barren						
	30X-2, 75-76	243.82	05	T	Barren						
	31X-3, 6-7	254.96	21	F	Fragments						
	32X-1, 27-28	261.87	63	F	.	.	.	.	.	1+	.
	34X-3, 66-67	284.46	05	B	Barren						
	35X-1, 79-80	291.19	05	T	Barren						
	36X-3, 91-92	304.01	42	F	Barren						
	37X-3, 98-100	313.78	42	T	.	.	.	.	.	.	.
	39X-2, 76-77 to	330.96	05	B	Barren						
	60X-3, 44-46	514.84	05	B	Barren						
	61X-4, 60-62	526.80	21	T	Fragments						
NN15?	64X-3, 19-20 to	552.39	05	B	Barren						
	67X-3, 83-84	583.03	05	B	Barren						

Note: Barren samples are summarized in Table 3. Numbers under *Dictyocha* spp. refer to trace or rare occurrences of *Dictyocha fibula* (1) and *Dictyocha varia* (2).

## Ebridians and Actiniscidians

### *Ammodochium serotinum* Zone

**Definition:** Interval from the LO of *Ebriopsis cornuta* to the LO of *Ammodochium serotinum* (Ling, 1973, cited as *Ammodochium rectangulare*). Upper Pliocene.

**Occurrence:** Found in Holes 860B and 861C. The LO of *E. cornuta* was not discovered in both holes. The LO of *A. serotinum* is well developed in Hole 861C, but indistinctly represented in Hole 860B. In Hole 861C, the LO of *A. serotinum* probably lies in the uppermost part of Zone NN18.

**Remarks:** Both the LADs of *E. cornuta* and *A. serotinum* are loosely defined at some northern Pacific sites, due to the absence of calcareous nannofossils. Generally, the LAD of *E. cornuta* is known for the early to late Pliocene transition, and the LAD of *A. serotinum* for the latest Pliocene. This is documented, for instance, in the northeastern Pacific Ocean at Site 173 (Ling, 1977), and in the northern Pacific at Site 192 (Ling, 1973). Both levels are most exactly determined at Site 644 in the Norwegian Sea (Locker and Martini, 1989) and at Site 798 in the Japan Sea (Ling, 1992). At Site 644, the LAD of *E. cornuta* is at the top of the Gauss Chron (top of Zone NN16) and the LAD of *A. serotinum* is below the Olduvai Subchron (near the top of Zone

NN18). At Site 798, the LAD of *E. cornuta* is above the Gauss Chron (probably Zone NN17) and the LAD of *A. serotinum* lies within the Olduvai Subchron.

### *Actiniscus pentasterias* Zone

**Definition:** Interval above the LO of *Ammodochium serotinum* (Locker and Martini, 1989). Uppermost Pliocene to Holocene.

**Occurrence:** Found in Holes 860B and 860C.

**Remarks:** This zone fills the interval above the LO of *A. serotinum*, as the last distinct ebridian marker.

## SITE SUMMARIES

### Site 860 (Tables 1 and 3)

From this site, one hole was investigated for siliceous flagellates. Hole 860B (45°51.972'S, 75°45.101'W; water depth 2145.9 m) was drilled on the middle slope of the Chile Trench, as the central member of the transect across the forearc basin. The sediment sequence recovered comprises predominantly clayey to silty sediments

Table 1 (continued).

<i>Ds. bioctonarius</i>	<i>Ds. speculum</i> , total	<i>Ds. speculum</i> f. <i>speculum</i>	Silicoflagellate abundance	Silicoflagellate zone	<i>A. serotinum</i>	<i>A. sp.</i> , small triodes	<i>E. tripartita</i>	<i>E. tripartita</i> , large triodes	<i>A. pentasterias</i>	Ebrid./actin. abundance	Ebridian/actiniscidian zone
.	R	R	R	<i>Dictyochoa</i>	.	.	.	.	R	R	<i>Actiniscus pentasterias</i>
.	R	R	F	<i>mess. aculeata</i>	.	.	.	.	R	R	
.	Barren		B	<i>Distephanus speculum</i>	.	.	Barren	.	R	B	
.	R	R	F		.	.	T	.	R	R	
.	R	R	R		.	.	.	.	R	R	
.	R	R	R		.	.	.	.	T	T	
.	R	R	R		.	.	.	.	R	R	
.	R	R	R		.	.	.	.	R	R	
.	Fragments		T		.	.	.	.	R	R	
.	Barren		B		.	.	Barren	.	R	B	
.	Fragments		T		.	.	.	.	R	R	
.	R	R	R		.	.	Barren	.		B	
.	Barren		B		.	.	Barren	.		B	
.	R	R	R		.	.	.	.	R	R	
.	Fragments		T		.	.	Barren	.		B	
.	Barren		B		.	.	.	.	R	R	
.	R	R	F		T	R	R	R	F	<i>Ammodochium serotinum</i>	
.	F	F	F		.	.	.	R	R		
.	R	R	R	<i>Distephanus</i>	.	.	.	.	T	T	Unzoned
.	F	F	F	<i>aculeatus</i>	R	T	.	R	R	R	
.	Barren		B	Unzoned	.	.	Barren	.		B	
.	Fragments		T		.	.	.	.	R	R	
R	R	R	R		.	.	.	.	R	R	
.	Barren		B		.	.	Barren	.		B	
.	Barren		B		.	.	Barren	.		B	
.	Barren		B		.	.	.	.	R	R	
.	R	R	R		.	.	.	.	R	R	
.	Barren		B		.	.	Barren	.		B	
.	Barren		B		.	.	.	.		B	
.	Fragments		T		.	.	Barren	.		B	
.	Barren		B		.	.	Barren	.		B	
.	Barren		B		.	.	.	.		B	

with a higher degree of consolidation toward the base (Shipboard Scientific Party, 1992a).

The sediments studied contain rather poorly preserved but, nevertheless, diversified assemblages of siliceous microfossils. These include protophytes (diatoms, silicoflagellates, ebridians, actiniscidians), protozoans (radiolarians), and remnants of metazoans (sponge spicules). Although abundances may be very low in certain layers, 19 out of 46 samples yielded siliceous flagellates. With a few exceptions, most of the flagellate assemblages are low in abundance and moderate to poor in preservation.

Samples 141-860B-1H-1, 90–91 cm, and -2H-1, 40–42 cm, contain *D. messanensis aculeata* and *A. pentasterias* and have been assigned to the *Dictyochoa messanensis aculeata* and the *Actiniscus pentasterias* Zones. The interval between Samples 141-860B-2H-2, 19–21 cm, and -22X-2, 103–104 cm, shows fluctuating abundances of siliceous flagellates that range from few to barren. But occurrences of *D. speculum* and *D. aculeatus* allow one to place Samples 141-860B-2H-2, 19–21 cm, to -20X-1, 127–128 cm, into the *Distephanus speculum* Zone, and Samples 141-860B-21X-1, 72–73 cm, and -22X-2, 103–104 cm, into the *Distephanus aculeatus* Zone. The presence of *A. pentasterias* and *A. serotinum* (including *Ammodochium* sp.)

indicates the *Actiniscus pentasterias* Zone down to Sample 141-860B-17X-CC, and the *Ammodochium serotinum* Zone from Sample 141-860B-19X-3, 100–101 cm, to -22X-2, 103–104 cm. Several occurrences of *D. messanensis* in Cores 141-860B-2H, -15X, and -19X obviously mark time levels of warmer surface-water conditions. With the exception of four samples displaying nonindicative silicoflagellate and actiniscidian species, the interval from Sample 141-860B-24X-2, 25–26 cm, down to the bottom of the hole remains barren of siliceous flagellates.

According to nannofossil data (Spiegler and Müller, this volume), the Pleistocene/Pliocene boundary lies above the LOs of *D. aculeatus* and *A. serotinum*. Although reworked Miocene to lower Pliocene silicoflagellates, such as *Dictyochoa fibula* and *Dictyochoa varia*, were found, the bulk of skeletons seems to be in place. No slumping or major reworking could be recognized.

#### Site 861 (Tables 2 and 3)

From this site, two holes were investigated for siliceous flagellates. Holes 861C (45°51.025'S, 75°41.531'W; water depth of 1652.2 m) and 861D (45°51.008'S, 75°41.499'W; water depth of 1652.2 m)

Table 2. Stratigraphic distribution of siliceous flagellates in upper Pliocene to Quaternary sediments of Holes 861C/D and correlation to standard nanofossil zones (Spiegler and Müller, this volume).

Nanno-plankton zone	Holes 861C/D, core, section, interval (cm)	Depth (mbsf)	Slide area (%)	Diatom abundance										
					<i>Da. messanensis</i> , total	<i>Da. mess. aculeata</i>	<i>Da. mess. messanensis</i>	<i>Da. mess. stapedia</i>	<i>Da. perlaevis</i>	<i>Da. spp.</i>	<i>Ds. aculeatus</i>	<i>Ds. bioctonarius</i>	<i>Ds. speculum</i> , total	<i>Ds. spec. f. pentagonus</i>
NN21	1H-1, 10-11	00.10	42	C	F	R	F	.	.	.	.	.	R	.
	1H-2, 40-41	01.90	21	C	F	R	F	.	R	.	.	.	F	.
	2H-5, 80-81	09.80	21	C	F	R	F	.	.	.	.	.	F	R
	3H-3, 99-100	16.49	42	C	R	.	R	.	.	.	.	.	F	R
	4H-1, 110-11	23.10	42	F	.	.	Fragments	.	.	.	.	.	Fragments	.
	4H-3, 33-34	25.33	42	C	R	.	R	.	.	.	.	.	.	.
	4H-4, 86-87	27.36	42	F	.	.	.	.	.	.	.	.	.	R
4H-6, 130-131	30.80	42	F	.	.	.	.	.	.	.	.	.	R	
NN20	5H-2, 60-62	32.40	21	F	R	.	R	.	.	.	.	.	R	.
	5H-4, 60-62	35.40	21	T	.	.	.	.	.	.	.	.	R	R
	5H-6, 60-62	38.40	42	F	.	.	Fragments	.	.	.	.	Fragments	.	
	5H-8, 58-60	41.38	42	F	.	.	.	.	.	.	.	.	R	.
	6H-2, 29-30	42.79	42	F	.	.	.	.	.	.	.	.	R	.
	6H-4, 79-80	46.29	21	T	.	.	.	.	.	.	.	.	R	.
	6H-6, 90-91	49.40	21	F	.	.	.	.	.	.	.	.	R	.
	6H-7, 60-61	50.60	42	F	.	.	.	.	.	2+	.	.	R	.
	7H-2, 120-121	53.20	21	T	.	.	Barren	.	.	.	.	Barren	F	.
	7H-3, 89-90	54.39	21	F	.	.	.	.	.	.	.	.	.	.
	7H-6, 25-26	58.25	42	F	F	.	F	.	.	.	.	.	.	.
	8H-3, 71-72	63.71	42	C	R	.	R	.	.	.	.	.	R	.
	8H-5, 10-11	66.10	42	C	R	R	R	T	.	.	.	.	.	.
	10H-3, 100-101	75.00	84	F	T	T	.	.	.	.	.	T	R	T
	12X-2, 50-51	92.10	63	C	.	.	.	.	.	.	.	.	F	.
	13X-1, 35-36	100.15	42	F	.	.	.	.	.	.	.	.	R	R
	14X-2, 70-71	111.60	63	F	.	.	.	.	.	.	.	.	R	.
15X-2, 68-69	121.18	42	C	R	R	.	.	.	.	.	.	F	T	
15X-4, 75-76	124.25	42	C	.	.	.	.	.	.	.	.	R	.	
NN19	16X-1, 72-73	129.62	42	C	T	T	.	.	.	.	.	.	R	.
	16X-5, 60-61	135.50	21	C	T	T	.	.	.	.	.	.	R	.
	17X-1, 109-110	139.79	42	C	.	.	.	.	.	.	.	.	R	.
	18X-3, 119-120	152.69	42	F	.	.	Fragments	.	.	.	.	Fragments	.	
	19X-CC	168.20	21	T	.	.	Barren	.	.	.	.	Barren	.	
	20X-CC	170.20	42	F	R	.	R	.	.	3+	.	.	R	.
?	21X-1, 100-101	171.20	84	F	.	.	.	.	.	.	R	.	R	.
	22X-4, 87-88	185.27	84	C	F	.	F	.	.	2+	.	.	C	R
	25X-3, 99-100	212.89	42	C	R	.	R	.	.	.	R	.	R	.
NN16-18	27X-2, 30-31	221.90	42	C	.	.	.	.	.	.	R	.	C	R
	27X-4, 30-31	223.81	21	C	.	.	.	.	.	2+	F	.	C	.
	28X-3, 40-41	231.50	21	C	T	.	T	.	.	.	F	.	F	.
	28X-5, 29-30	234.03	21	C	R	.	R	.	.	.	R	.	R	.
	29X-1, 75-76	238.45	42	T	.	.	.	.	.	.	R	.	R	.
	32X-2, 44-46	258.94	42	F	.	.	.	.	.	.	.	.	R	.
	33X-2, 70-71	268.80	21	T	.	.	Barren	.	.	.	.	Barren	.	
NN16	34X-3, 73-74	278.18	10	B	.	.	Barren	.	.	.	.	Barren	.	
	35X-3, 100-101	289.03	63	F	.	.	.	.	.	1+	.	.	.	
	36X-3, 100-101	299.60	42	F	.	.	.	.	.	.	T	.	R	
	37X-1, 90-91	305.80	42	F	.	.	.	.	.	.	.	.	R	
	39X-3, 100-101	327.32	21	T	.	.	Barren	.	.	.	.	Barren	.	
	40X-3, 87-88	337.77	21	T	.	.	Barren	.	.	.	.	Barren	.	
	1R-3, 100-101	346.30	42	F	R	.	R	.	.	2+	.	.	R	
	41X-3, 99-100	347.39	42	F	R	.	R	T	.	.	.	.	R	
	2R-1, 130-131 to 16R-2, 21-22	353.20 487.71	05 05	B B	.	.	Barren	.	.	.	.	.	Barren	.

Note: Barren samples are summarized in Table 3. Numbers under *Dictyocha* spp. refer to trace or rare occurrences of *Dictyocha fibula* (1), *Dictyocha varia* (2), and *Dictyocha* sp. (3).



Table 2 (continued).

<i>Ds. spec. f. speculum</i>	<i>Ds. spec. f. septenarius</i>	<i>Ds. spec. f. octonarius</i>	Silicoflagellate abundance	Silicoflagellate zone	<i>A. serotinum</i>	<i>A. sp., small triodes</i>	<i>E. tripartita</i>	<i>E. sp., large triodes</i>	<i>A. pentasterias</i>	Ebrid./actin. abundance	Ebridian/actiniscidian zone
R	.	T	F	<i>Dietyocha mess. aculeata</i>	.	.	.	T	R	R	<i>Actiniscus pentasterias</i>
F	.	.	C		.	.	T	T	R	R	
R	.	.	F	.	.	.	.	.	R	R	
R	.	.	F	.	.	R	R	R	R	R	
Fragments			T	.	.	.	.	.	R	R	
.	.	.	R	.	.	.	R	R	R	F	
R	.	.	R	.	.	.	.	.	R	R	
R	T	.	R	.	.	.	.	.	R	R	
R	.	.	R	.	.	.	.	.	T	T	
.	.	.	R	.	.	.	.	.	T	T	
Fragments			T	.	.	.	.	.	R	R	
R	.	.	R	.	.	.	.	.	T	T	
R	.	.	R	.	.	.	.	.	R	R	
R	.	.	R	.	.	.	.	.	T	T	
R	.	.	R	.	.	.	.	.	R	R	
R	.	.	R	.	.	.	.	.	R	R	
R	.	.	R	.	.	.	.	.	R	R	
F	Barren	.	B	.	.	Barren	.	.	R	B	
F	.	.	F	.	.	Barren	.	.	F	B	
.	.	.	F	.	.	R	R	F	R	F	
R	.	.	R	.	.	R	R	R	R	R	
.	.	.	R	.	.	.	.	.	R	R	
R	.	R	R	.	R	T	.	F	F	F	
F	.	.	F	.	.	R	R	R	R	R	
R	.	.	R	.	.	R	R	T	R	R	
R	.	.	R	.	.	.	.	.	R	R	
F	T	.	F	.	.	.	.	T	T	T	
R	.	.	R	.	.	T	.	R	R	R	
R	.	.	R	.	.	.	R	R	R	R	
R	.	.	R	.	.	Barren	.	.	R	B	
R	.	.	R	.	.	R	.	.	R	R	
Fragments Barren			B	.	Barren	Barren	.	.	B	Barren	
R	.	.	R	.	Barren	Barren	.	.	B	Barren	
R	T	.	R	.	R	.	.	.	R	R	<i>Ammodochium serotinum</i>
C	T	R	C	.	R	.	R	F	F	C	
R	.	R	R	.	.	.	.	R	R	R	
C	.	R	C	.	.	.	.	F	F	F	
C	.	R	C	.	.	.	.	F	F	F	
R	.	F	C	.	R	R	.	C	C	C	
R	.	.	R	.	R	.	.	F	F	F	
.	.	R	R	.	T	.	R	R	R	R	
R	.	.	R	.	.	.	.	R	R	R	
Barren			B	.	Barren	Barren	.	.	B	B	
Barren			B	.	Barren	Barren	.	.	B	B	
.	.	.	R	.	R	T	.	R	R	R	
R	.	R	R	.	R	.	.	T	R	R	
R	.	.	R	.	T	T	.	R	R	R	
Barren			B	.	Barren	Barren	.	.	B	Barren	
Barren			B	.	Barren	Barren	.	.	B	Barren	
R	.	.	F	.	R	.	.	.	R	R	
.	.	R	R	.	R	.	.	R	R	R	
Barren			B	.	Barren	Barren	.	.	B	Barren	
Barren			B	.	Barren	Barren	.	.	B	Barren	

**Table 3. Samples from Holes 860B, 861C, and 861D that are barren of siliceous flagellates.**

Core, section, interval (cm)	Core, section, interval (cm)
141-860B-	64X-3, 19-20
2H-2, 19-21	65X-CC
6H-3, 95-95	141-861C-
14X-2, 48-49	7H-2, 120-121
24X-2, 25-26	19X-CC
25X-2, 32-33	33X-2, 70-71
29X-1, 126-127	34X-3, 73-74
30X-2, 75-76	39X-3, 100-101
34X-3, 66-67	40X-3, 87-88
35X-1, 79-80	
39X-2, 76-77	141-861D-
41X-3, 100-101	2R-1, 130-131
43X-1, 34-35	6R-3, 99-100
46X-1, 48-49	7R-3, 99-100
47X-CC	8R-1, 56-57
48X-3, 100-101	9R-2, 39-40
53X-2, 75-76	11R-2, 70-71
55X-CC	12R-2, 36-37
56X-CC	13R-3, 102-103
57X-CC	15R-2, 25-26
60X-3, 44-46	16R-2, 21-22

were drilled on the upper slope of the Chile Trench, as the eastern endmembers of the transect across the forearc basin. The sediment sequences recovered consist of clayey, silty, and sandy sediments with a higher degree of consolidation toward the base (Shipboard Scientific Party, 1992b).

The sediments studied contain assemblages of siliceous microfossils that are more diversified and better preserved than those in Hole 860B. The assemblages include protophytes (diatoms, silicoflagellates, ebridians, actiniscidians, and scattered chrysomonads), protozoans (radiolarians), remnants of metaphytes (scattered phytoliths) and metazoans (sponge spicules). Forty-six out of 62 samples yielded siliceous flagellates. Abundances of flagellate assemblages may be higher than those in isochronous intervals of Hole 860B, but preservation is likewise moderate to poor.

The interval between Samples 141-861C-1H-1, 10-11 cm, and -2H-5, 80-81 cm, is characterized by *D. messanensis aculeata* and *A. pentasterias* and has been placed into the *Dictyochoa messanensis aculeata* and the *Actiniscus pentasterias* Zones. Rather continuous occurrences of *D. speculum* and *D. aculeatus* indicate the *Distephanus speculum* Zone from Sample 141-861C-3H-3, 99-100 cm, to -20-CC, and the *Distephanus aculeatus* Zone from Sample 141-861C-21X-1, 100-101 cm, to -36X-3, 100-101 cm. Skeletons of *A. pentasterias* reflect the *Actiniscus pentasterias* Zone down to Sample 141-861C-18X-3, 119-120 cm. As evidenced by well-preserved specimens of *A. serotinum* (including *Ammodoichium* sp.), the *Ammodoichium serotinum* Zone follows from Samples 141-861C-21X-1, 100-101 cm, to -41X-3, 99-100 cm. Distinct occurrences of *D. messanensis* and of *Ebria tripartita* in the Quaternary to Pliocene sediment sequence obviously correspond to time levels of warmer surface-water conditions. The interval from Sample 141-861D-2R-1, 130-131 cm, down to the bottom of the hole remains barren of siliceous flagellates.

According to nanofossil data (Spiegler and Müller, this volume), the Pliocene/Pleistocene boundary lies immediately above the LOs of *D. aculeatus* and *A. serotinum*. Reworked Miocene to lower Pliocene silicoflagellates, such as *D. fibula* and *D. varia*, were found, but the bulk of skeletons seems to be in place. The somewhat misleading occurrences of *A. serotinum* and *Ammodoichium* sp. in Sample 141-861C-10H-3, 100-101 cm, may be explained by minor slumping or by reworking.

### PALEOCEANOGRAPHIC IMPLICATIONS

In recent time, the oceanography of the southeastern South Pacific is determined by subpolar and transitional surface-water masses (Fig. 3). South of the study area are polar and subpolar waters that are

driven by the Antarctic Circumpolar Current in an easterly direction around the Antarctic continent. West and northwest of the study area are transitional and subtropical waters that are part of the South Pacific Central Water mass and are driven within the South Pacific Gyre. The most important feature of the study area is the Peru Current, which branches from the Antarctic Circumpolar Current and drives subpolar water along the coast of South America to the north (Bearman, 1993; Pickard and Emery, 1993).

Surface-water masses generally are inhabited by plankton organisms. According to their ecological preferences, these organisms may reflect particular surface-water conditions, especially temperature, salinity, and nutrient content. This holds true also for the plankton groups studied (i.e., silicoflagellates, ebridians, and actiniscidians). Recent data are scarce for the study area off southern Chile. Nevertheless, certain aspects of Pliocene and Quaternary paleoenvironmental changes can be revealed, if planktologic and oceanographic analogies are taken into consideration.

In recent time, the distribution of the silicoflagellate species *Dictyochoa messanensis* and *Distephanus speculum* follows the temperature gradient from equatorial to polar regions, with *D. messanensis* solely present at the equator and *D. speculum* solely near the pole. The region inbetween is usually inhabited by mixed populations at varying proportions. This has been revealed by the investigation of water samples from the South Atlantic (Gemeinhardt, 1934) and of surface sediment samples from the North Pacific (Poelchau, 1976). The very detailed investigation of Poelchau exhibited, furthermore, that *D. messanensis messanensis* is sedimented in higher abundance beneath tropical and transitional waters and *D. messanensis aculeata* (cited as *Dictyochoa epiodon*) beneath transitional water.

During late Pliocene time, plankton communities were obviously determined by cool surface-water conditions at Site 861 (Fig. 4, Table 2), which may be assumed to have been under the influence of an early Peru Current. This is indicated by higher percentages of the cool-adapted species *D. speculum* and the closely related *D. aculeatus*. But, during four time intervals prior to 1.6 Ma (top of Zone NN18, as used here), probably transitional surface-water masses intruded into the investigation area, which diminished the influence of the Peru Current. This may be deduced from the occurrence of the warm-adapted taxon *D. messanensis messanensis*. In general, communities of siliceous flagellates contained more taxa during late Pliocene time than during Quaternary time, which may hint at more favorable paleoenvironmental conditions.

During early and middle Pleistocene times, plankton communities were living mainly under cool surface-water conditions at Site 861, as continuous occurrences of *D. speculum* and *Actiniscus pentasterias* demonstrate. But, surprisingly, also during those periods, several expansions of transitional water masses proceeded off the coast of southern South America. This is indicated especially by occurrences of the warm- to temperate-adapted taxa *D. messanensis messanensis* and *D. messanensis aculeata*. The most prominent expansion of warm water masses was probably between 0.5 and 0.3 Ma (within Zone NN20), when *D. speculum* disappeared temporarily, but *Distephanus bioctonarius* and *Ebria tripartita* occurred in the area. In recent time, *D. bioctonarius* (often cited as *Octactis pulchra*) attains maximum densities in tropical and subtropical waters off Middle America and Peru.

After 0.3 Ma (base of Zone NN21), during late Pleistocene and Holocene times, warm surface-water masses intruded once more into the study area. According to the continuous occurrence of *D. messanensis aculeata*, transitional water expanded, probably iteratively rather than permanently, from the central South Pacific Ocean toward the coast. This may have caused mixing with the cool water masses of the Peru Current, thus diminishing its intensity. But, narrowing or dislodging of the Peru Current may have occurred, too. Although definitive explanations cannot be given at present, the Pliocene to Quaternary warming episodes may have been affected by enhanced El Niño activities, which resulted both in a southward shift of the



transitional surface-water belt and an eastward expansion of respective water masses.

## TAXONOMY

Silicoflagellates, ebridians, and actiniscidians represent specific groups of siliceous flagellates that may be placed into two classes of protophytes (Tappan, 1980). Silicoflagellates are related to the Chrysophyceae, and ebridians and actiniscidians to the Dinophyceae. As the taxonomy and synonymies of the species found were discussed earlier (Locker, 1974; Locker and Martini, 1986a, 1986b, 1990), only a list of taxa is given. Remarks about autochthonous species have been added.

### Silicoflagellates

- Dictyocha fibula* Ehrenberg, 1840 (not figured)  
*Dictyocha messanensis* Haeckel, 1861  
*Dictyocha messanensis* subsp. *aculeata* (Lemmermann, 1901) Locker and Martini, 1986: Pl. 1, Fig. 2  
*Dictyocha messanensis* Haeckel, 1861 subsp. *messanensis*: Pl. 1, Fig. 4  
*Dictyocha messanensis* subsp. *stapedia* (Haeckel, 1887) Locker and Martini, 1986: Pl. 1, Fig. 3  
*Dictyocha perlaevis* Frenguelli, 1951: Pl. 1, Fig. 1  
*Dictyocha varia* Locker, 1975; synonym *Dictyocha pulchella* Bukry, 1975 (not figured)  
*Distephanus aculeatus* (Ehrenberg, 1841) Haeckel, 1887; synonym *Distephanus boliviensis* Frenguelli, 1940  
*Distephanus aculeatus* (Ehrenberg, 1841) f. *aculeatus*: Pl. 1, Fig. 14  
*Distephanus aculeatus* f. *binoculus* (Ehrenberg, 1845) Locker and Martini, 1986: Pl. 1, Fig. 15  
*Distephanus bioctonarius* (Ehrenberg, 1846) Martini, 1990; synonym *Octactis pulchra* Schiller, 1925: Pl. 1, Fig. 13  
*Distephanus speculum* (Ehrenberg, 1840) Haeckel, 1887  
*Distephanus speculum* f. *octonarius* (Ehrenberg, 1845) Locker and Martini, 1986: Pl. 1, Fig. 12  
*Distephanus speculum* f. *pentagonus* (Lemmermann, 1901) Locker and Martini, 1986: Pl. 1, Fig. 9  
*Distephanus speculum* f. *septenarius* (Ehrenberg, 1845) Locker and Martini, 1986: Pl. 1, Fig. 11  
*Distephanus speculum* (Ehrenberg, 1840) f. *speculum*: Pl. 1, Fig. 10  
*Dictyocha messanensis* typically has a rhombic basal ring and a longitudinal apical bar with a short pike (subsp. *messanensis*). One subspecies displays a basal ring with more curved bars (subsp. *stapedia*), and another one, a basal ring with additional spines (subsp. *aculeata*). Variants without an apical pike are not taxonomically discriminated here. As in other species, populations may include skeletons with thicker and with thinner skeletal bars.  
*Dictyocha perlaevis* possesses a large basal ring with distinctly curved bars and a short longitudinal apical bar. Basal pikes are usually missing.  
*Distephanus aculeatus* usually has a hexagonal basal ring and an apical ring without diagnostic pikes (f. *aculeatus*). This skeletal structure corresponds to *D. speculum*, but in *D. aculeatus* the basal ring generally appears larger, and apical pikes, if present, do not exhibit a preferential orientation. Variants with more than six basal spines (not found in Leg 141 material) or with a fenestrated apical window (f. *binoculus*) are rare in populations.  
*Distephanus bioctonarius* normally has an octagonal basal ring and a delicate apical ring which often breaks off. Variants with more, or with less, than eight basal spines are usually rare in populations and have not been found in the material.  
*Distephanus speculum* usually possesses a hexagonal basal ring and an apical ring with two pikes (f. *speculum*). But populations may contain a lot of other variants: forms with five basal spines (f. *pentagona*), with seven spines (f. *septenarius*), with eight spines (f. *octonarius*), or with a differently constructed apical apparatus (not found in the material). Here, a broad species concept is applied that embraces skeletons with basal pikes and a narrow apical ring, and skeletons without basal pikes and a wider apical ring.

### Ebridians and Actiniscidians

- Ammodoichium serotinum* Locker and Martini, 1986: Pl. 1, Fig. 6  
*Ammodoichium* sp.; small triodes: Pl. 1, Fig. 5  
*Ebria tripartita* Schumann, 1867: Pl. 1, Fig. 8  
*Ebria* sp.; large triodes: Pl. 1, Fig. 7  
*Actiniscus pentasterias* Ehrenberg, 1841: Pl. 1, Fig. 16

*Ammodoichium serotinum* has a triode that bears triplets of proclades and opisthoclares plus highly arched synclades. Juvenile (and corroded?) skeletons may be delivered by the triode alone, or by the triode and rudimentary proclades and opisthoclares. Small triodes that probably belong to this species have been indicated as *Ammodoichium* sp. in tables.

*Ebria tripartita* has a large triode that is connected with the ring of synclades by short proclades. Juvenile (and corroded?) skeletons may solely consist of the large triode. Large triodes, which probably belong to this species, are indicated as *Ebria* sp. in tables.

*Actiniscus pentasterias* is usually represented by five-rayed stellate bodies that possess crests and depressions. Sometimes populations contain also four-, six- or seven-rayed specimens. Here, a broad species concept was used that includes skeletons having larger and smaller peripheral depressions, and longer and shorter arms.

## CONCLUSIONS

1. The sediment sequences investigated from Holes 860B, 861C, and 861D were assigned to the *Distephanus aculeatus*, *Distephanus speculum*, and *Dictyocha messanensis aculeata* Zones of silicoflagellates and to the *Ammodoichium serotinum* and *Actiniscus pentasterias* Zones of ebridians/actiniscidians. This indicates a late Pliocene to Quaternary age of related intervals. Sediments having a higher age could not be proven due to the absence of siliceous flagellates in the lower part of sequences.

2. According to the LOs of *Distephanus aculeatus* and *Ammodoichium serotinum*, the Pliocene/Pleistocene boundary can be placed above 141.10 mbsf in Hole 860B, and above 171.20 mbsf in Hole 861C. This confirms, in general, the nannofossil results of Spiegler and Müller (this volume), who draw this boundary between 128.07 and 126.50 mbsf in Hole 860B and between 170.64 and 170.20 mbsf in Hole 861C.

3. Fluctuating occurrences of *Dictyocha messanensis messanensis* throughout the upper Pliocene to Pleistocene section of Holes 860B and 861C record varying paleoceanographic conditions. Repeated occurrences of this subspecies indicate that the cool surface-water masses of the Peru Current were mixed with or replaced by warm transitional water masses of the central South Pacific several times.

4. The consistent occurrence of *Dictyocha messanensis aculeata* in sediments of late NN21 age reflects a longer period of prevailing warm surface-water conditions, caused by repeated expansions of transitional water masses toward the coast of southern South America.

5. The Pliocene and Quaternary warming episodes may have been caused by strong El Niño activities, which resulted in a southward shift of the transitional water belt and an expansion of warmer surface water over the study area.

## ACKNOWLEDGMENTS

This study profited from discussions with J.O. Backhaus (Hamburg), J.H. Behrmann (Giessen), M. Knoll (Kiel), and D. Spiegler (Kiel). The ODP staff at College Station, Texas, supplied the initial sample set and, later, some additional samples. The manuscript was carefully reviewed by K. McCartney, and K. von Salis Perch-Nielsen. The Deutsche Forschungsgemeinschaft provided financial support. All this help is gratefully acknowledged.

## REFERENCES\*

- Bearman, G. (Ed.), 1993. *Ocean Circulation* (3rd ed.): Walton Hall (The Open University).  
 Behrmann, J.H., Lewis, S.D., Musgrave, R.J., et al., 1992. *Proc. ODP, Init. Repts.*, 141: College Station, TX (Ocean Drilling Program).  
 Berggren, W.A., Kent, D.V., Flynn, J.J., and Van Couvering, J.A., 1985. Cenozoic geochronology. *Geol. Soc. Am. Bull.*, 96:1407-1418.

\* Abbreviations for names of organizations and publications in ODP reference lists follow the style given in *Chemical Abstracts Service Source Index* (published by American Chemical Society).

- Bukry, D., 1978. Cenozoic silicoflagellate and coccolith stratigraphy, south-eastern Atlantic Ocean, Deep Sea Drilling Project Leg 40. In Bolli, H.M., Ryan, W.B.F., et al., *Init. Repts. DSDP*, 40: Washington (U.S. Govt. Printing Office), 635–649.
- , 1979a. Coccolith and silicoflagellate stratigraphy, northern Mid Atlantic Ridge and Reykjanes Ridge, Deep Sea Drilling Project Leg 49. In Luyendyk, B.P., Cann, J.R., et al., *Init. Repts. DSDP*, 49: Washington (U.S. Govt. Printing Office), 551–581.
- , 1979b. Comments on opal phytoliths and stratigraphy of Neogene silicoflagellates and coccoliths at Deep Sea Drilling Project Site 397 off northwest Africa. In Luyendyk, B.P., Cann, J.R., et al., *Init. Repts. DSDP*, 49: Washington (U.S. Govt. Printing Office), 977–1009.
- , 1980. Silicoflagellate biostratigraphy and paleoecology in the eastern equatorial Pacific, Deep Sea Drilling Project Leg 54. In Rosendahl, B.R., Hekinian, R., et al., *Init. Repts. DSDP*, 54: Washington (U.S. Govt. Printing Office), 545–573.
- Bukry, D., and Foster, J.H., 1973. Silicoflagellate and diatom stratigraphy, Leg 16, Deep Sea Drilling Project. In van Andel, T.H., Heath, G.R., et al., *Init. Repts. DSDP*, 16: Washington (U.S. Govt. Printing Office), 815–871.
- Ciesielski, P.F., 1975. Biostratigraphy and paleoecology of Neogene and Oligocene silicoflagellates from cores recovered during Antarctic Leg 28, Deep Sea Drilling Project. In Hayes, D.E., Frakes, L.A., et al., *Init. Repts. DSDP*, 28: Washington (U.S. Govt. Printing Office), 625–691.
- Dumitrica, P., 1973. Paleocene, late Oligocene and post-Oligocene silicoflagellates in southwestern Pacific sediments cored on DSDP Leg 21. In Burns, R.E., Andrews, J.E., et al., *Init. Repts. DSDP*, 21: Washington (U.S. Govt. Printing Office), 837–883.
- Gemeinhardt, K., 1934. Die Silicoflagellaten des südatlantischen Ozeans. *Wiss. Ergebnisse Deutsch. Exped. "Meteor" 1925–27*, 12:274–312.
- Levitus, S., 1982. *Climatological Atlas of the World Ocean*. NOAA Prof. Pap., 13.
- Ling, H.Y., 1973. Silicoflagellates and ebridians from Leg 19. In Creager, J.S., Scholl, D.W., et al., *Init. Repts. DSDP*, 19: Washington (U.S. Govt. Printing Office), 751–775.
- , 1975. Silicoflagellates and ebridians from Leg 31. In Karig, D.E., Ingle, J.C., Jr., et al., *Init. Repts. DSDP*, 31: Washington (U.S. Govt. Printing Office), 763–777.
- , 1977. Late Cenozoic silicoflagellates and ebridians from the eastern North Pacific region. In Saito, T., and Ujiie, H. (Eds.), *Proc. First Int. Congr. Pacific Neogene Stratigraphy*, 1:205–233.
- , 1992. Late Neogene silicoflagellates and ebridians from Leg 128, Sea of Japan. In Pisciotto, K.A., Ingle, J.C., Jr., von Breymann, M.T., Barron, J., et al., *Proc. ODP, Sci. Results*, 127/128 (Pt. 1): College Station, TX (Ocean Drilling Program), 237–248.
- Locker, S., 1974. Revision der Silicoflagellaten aus der Mikrogeologischen Sammlung von C.G. Ehrenberg. *Eclogae Geol. Helv.*, 67:631–646.
- Locker, S., and Martini, E., 1986a. Ebridians and actiniscidians from the southwest Pacific. In Kennett, J.P., von der Borch, C.C., et al., *Init. Repts. DSDP*, 90: Washington (U.S. Govt. Printing Office), 939–951.
- , 1986b. Silicoflagellates and some sponge spicules from the southwest Pacific, DSDP Leg 90. In Kennett, J.P., von der Borch, C.C., et al., *Init. Repts. DSDP*, 90: Washington (U.S. Govt. Printing Office), 887–924.
- , 1989. Cenozoic silicoflagellates, ebridians, and actiniscidians from the Vøring Plateau (ODP Leg 104). In Eldholm, O., Thiede, J., Taylor, E., et al., *Proc. ODP, Sci. Results*, 104: College Station, TX (Ocean Drilling Program), 543–585.
- Martini, E., 1990. Tertiary and Quaternary silicoflagellates, actiniscidians, and ebridians from the eastern Pacific off Peru (Leg 112). In Suess, E., von Huene, R., et al., *Proc. ODP, Sci. Results*, 112: College Station, TX (Ocean Drilling Program), 157–173.
- Martini, E., and Müller, C., 1986. Current Tertiary and Quaternary calcareous nannoplankton stratigraphy and correlations. *Newsl. Stratigr.*, 16:99–112.
- Okada, H., and Bukry, D., 1980. Supplementary modification and introduction of code numbers to the low-latitude coccolith biostratigraphic zonation (Bukry, 1973; 1975). *Mar. Micropaleontol.*, 5:321–325.
- Perch-Nielsen, K., 1975. Late Cretaceous to Pleistocene silicoflagellates from the southern Southwest Pacific, DSDP, Leg 29. In Kennett, J.P., Houtz, R.E., et al., *Init. Repts. DSDP*, 29: Washington (U.S. Govt. Printing Office), 677–721.
- Pickard, G.L., and Emery, W.J., 1993. *Descriptive Physical Oceanography*: New York (Pergamon).
- Poelchau, H.S., 1976. Distribution of Holocene silicoflagellates in North Pacific sediments. *Micropaleontology*, 22:164–193.
- Shipboard Scientific Party, 1992a. Site 860. In Behrmann, J.H., Lewis, S.D., Musgrave, R.J., et al., *Proc. ODP, Init. Repts.*, 141: College Station, TX (Ocean Drilling Program), 159–238.
- , 1992b. Site 861. In Behrmann, J.H., Lewis, S.D., Musgrave, R.J., et al., *Proc. ODP, Init. Repts.*, 141: College Station, TX (Ocean Drilling Program), 239–299.
- Tappan, H., 1980. *The Paleobiology of Plant Protists*: San Francisco (W.H. Freeman).

**Date of initial receipt: 5 October 1993**

**Date of acceptance: 2 November 1994**

**Ms 141SR-042**




Plate 1. Upper Pliocene and Quaternary siliceous flagellates. All specimens magnified  $\times 800$ , bar = 10  $\mu\text{m}$ . 1. *Dictyocha perlaevis* Frenguelli, Sample 141-860B-19X-3, 100–101 cm, Pliocene. 2. *Dictyocha messanensis* subsp. *aculeata* (Lemmermann), Sample 141-861C-1H-1, 10–11 cm, Quaternary. 3. *Dictyocha messanensis* subsp. *stapedia* (Haeckel), Sample 141-861C-41X-3, 99–100 cm, Pliocene. 4. *Dictyocha messanensis* Haeckel subsp. *messanensis*, Sample 141-861C-20X-CC, Quaternary. 5. *Ammodochium* sp., small triode, Sample 141-861C-28X-3, 40–41 cm, Pliocene. 6. *Ammodochium serotinum* Locker and Martini, Sample 141-861C-22X-4, 87–88 cm, Pliocene. 7. *Ebria* sp., large triode, Sample 141-861C-1H-1, 10–11 cm, Quaternary. 8. *Ebria tripartita* Schumann, Sample 141-860B-2H-4, 29–31 cm, Quaternary. 9. *Distephanus speculum* f. *pentagonus* (Lemmermann), Sample 141-861C-22X-4, 87–88 cm, Pliocene. 10. *Distephanus speculum* (Ehrenberg) f. *speculum*, Sample 141-860B-12X-3, 100–101 cm, Quaternary. 11. *Distephanus speculum* f. *septenarius* (Ehrenberg), Sample 141-861C-15X-2, 68–69 cm, Quaternary. 12. *Distephanus speculum* f. *octonarius* (Ehrenberg), Sample 141-861C-22X-4, 87–88 cm, Pliocene. 13. *Distephanus bioctonarius* (Ehrenberg), Sample 141-861C-22X-4, 87–88 cm, Pliocene. 14. *Distephanus aculeatus* (Ehrenberg) f. *aculeatus*, Sample 141-861C-27X-2, 30–31 cm, Pliocene. 15. *Distephanus aculeatus* f. *binoculus* (Ehrenberg), Sample 141-861C-28X-3, 40–41 cm, Pliocene. 16. *Actiniscus pentasterias* Ehrenberg, Sample 141-861C-7H-6, 25–26 cm, Quaternary.