



## ***Corrigendum to*** **“Delivery of halogenated very short-lived substances from the west Indian Ocean to the stratosphere during the Asian summer monsoon” published in Atmos. Chem. Phys., 17, 6723–6741, 2017**

**Alina Fiehn<sup>1,2</sup>, Birgit Quack<sup>1</sup>, Helmke Hepach<sup>1,a</sup>, Steffen Fuhlbrügge<sup>1</sup>, Susann Tegtmeier<sup>1</sup>, Matthew Toohey<sup>1</sup>, Elliot Atlas<sup>3</sup>, and Kirstin Krüger<sup>2</sup>**

<sup>1</sup>GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

<sup>2</sup>Meteorology and Oceanography Section, Department of Geosciences, University of Oslo, Oslo, Norway

<sup>3</sup>Rosenstiel School of Marine and Atmospheric Science, University of Miami, Miami, USA

<sup>a</sup>now at: Environment Department, University of York, York, UK

**Correspondence:** Kirstin Krüger (kkrueger@geo.uio.no)

Published: 30 April 2021

In the abovementioned paper, two errors appear in Eq. (2) and the following sentence. They are corrected to

$$k_w = k_{600} \cdot \left( \frac{Sc}{600} \right)^{-\frac{1}{2}} \quad (2)$$

Nightingale et al. (2000) determined the transfer coefficient ( $k_{600}$ ) as a function of the wind speed at 10 m height ( $u_{10}$ ):  
 $k_{600} = 0.222(u_{10})^2 + 0.333u_{10}$ .

*Data availability.* The underlying data are now available at the open-access library Pangaea: <https://doi.org/10.1594/PANGAEA.876115> (Fiehn et al., 2017).

### **References**

Fiehn, A., Quack, B., Hepach, H., Fuhlbrügge, S., Tegtmeier, S., Toohey, M., Atlas, E. L., and Krüger, K.: Delivery of halogenated very short-lived substances from the west Indian Ocean to the stratosphere during Asian summer monsoon, PANGAEA, <https://doi.org/10.1594/PANGAEA.876115>, 2017.