



Interaction between mafic magma and lithospheric mantle: Evidence from the geochemistry of olivines and olivine-hosted melt inclusions in lavas from Kibblewhite Volcano, Kermadec arc

Yasuhiro Hirai (1,2), Yoshihiko Tamura (2), Kaj Hoernle (3), Christian Timm (4), Folkmar Hauff (3), Reinhard Werner (3), Takeshi Hanyu (2), Bogdan Vaglarov (2), Qing Chang (2), Takashi Miyazaki (2), and Jun-Ichi Kimura (2)

(1) Division of Natural System, Kanazawa University, Kanazawa, Ishikawa, Japan, (2) Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokosuka, Kanagawa, Japan, (3) GEOMAR Helmholtz Centre for Ocean Research, Kiel, Germany, (4) GNS Science, Lower Hutt, New Zealand

During German R/V SONNE cruise (SO-255) mafic basaltic to magnesian andesitic lavas were recovered from the Kermadec arc front volcano Kibblewhite (Hirai et al., EGU2018). Notably, these primitive lavas contain forsteritic olivines and olivine-hosted melt inclusions, which were re-homogenized using a heating stage. Here, we present the chemical variations of the melt inclusions and their host olivines and discuss the interaction between mafic melts and subarc lithospheric mantle.

The olivines are divided into two groups based on their occurrence in different host rock and NiO contents. Olivines ($\text{Fo}_{90.3-92.4}$) found as phenocrysts in ankaramites (olivine-clinopyroxene rich basalts; 50.4–50.6 wt.% SiO_2 ; 13.1–13.5 wt.% MgO ; $\text{Mg\#} = 71.8-72.1$) show unusually low Ni contents (912–1462 ppm). Olivines ($\text{Fo}_{91.0-92.2}$) occurring as xenocrysts in magnesian andesites (57.5–57.6 wt.% SiO_2 ; 5.5–5.6 wt.% MgO ; $\text{Mg\#} = 56.8-57.3$) show higher Ni contents (1108–3057 ppm) than olivines in ankaramites.

All re-homogenized melt inclusions have primitive basalt compositions (49.6–52.0 wt.% SiO_2 ; 12.7–15.9 wt.% MgO , 0.36–0.58 wt.% TiO_2 , 1.2–2.0 wt.% $\text{Na}_2\text{O}+\text{K}_2\text{O}$, 0.06–0.37 wt.% SO_3 , and 0.05–0.20 wt.% Cl), which could represent parental melts for the ankaramites. However, melt inclusions hosted by olivines in ankaramites have lower Al_2O_3 and higher CaO contents than those hosted by olivine xenocrysts in magnesian andesites; $\text{CaO}/\text{Al}_2\text{O}_3$ values are 1.2–1.6 and 0.9–1.3 respectively, indicating that there is a negative correlation between Ni contents in host olivines and $\text{CaO}/\text{Al}_2\text{O}_3$ values in their melt inclusions.

The variable Ni and constant Fo contents in the host olivines could be explained by an interaction of primary mafic melts with lithospheric mantle, which causes assimilation of pyroxenes in the lithosphere and simultaneous fractional crystallization of olivines (AFC; Kelemen, 1990; Tamura et al., 2018). Moreover, the high $\text{CaO}/\text{Al}_2\text{O}_3$ values in the melt inclusions hosted by the ankaramite olivines cannot be produced by partial melting of lherzolite (e.g. as compilation by Médard et al., 2004). These melts have most likely extensively interacted with the subarc lithospheric mantle. In addition, a negative correlation between Ni content in the host olivine and $\text{CaO}/\text{Al}_2\text{O}_3$ ratio in their melt inclusions supports the melt-mantle interaction. Conversely, xenocrystic olivines in the andesites with high Ni contents must have been crystallized from more primitive basaltic melts within lithospheric mantle prior to significant melt-mantle interaction. These olivines were then trapped by primary magnesian andesite magmas as xenocrysts.