



Comparing CLE-AdCSV applications using SA and TAC to determine the Fe-binding characteristics of model ligands in seawater

Loes J. A. Gerringa¹, Martha Gledhill², Indah Ardiningsih¹, Niels Muntjewerf¹, and Luis M. Laglera³

¹Royal Netherlands Institute for Sea Research (NIOZ), Department of Ocean Systems, Texel, the Netherlands

²GEOMAR Helmholtz Centre for Ocean Research, 24148 Kiel, Germany

³FI-TRACE, Departamento de Química and Laboratori Interdisciplinari sobre Canvi Climàtic, Universidad de las Islas Baleares, Palma, Balearic Islands 07122, Spain

Correspondence: Loes J. A. Gerringa (loes.gerringa@nioz.nl)

Received: 27 May 2021 – Discussion started: 23 June 2021

Revised: 10 August 2021 – Accepted: 11 August 2021 – Published: 30 September 2021

Abstract. Competitive ligand exchange–adsorptive cathodic stripping voltammetry (CLE-AdCSV) is used to determine the conditional concentration ($[L]$) and the conditional binding strength ($\log K^{\text{cond}}$) of dissolved organic Fe-binding ligands, which together influence the solubility of Fe in seawater. Electrochemical applications of Fe speciation measurements vary predominantly in the choice of the added competing ligand. Although different applications show the same trends, $[L]$ and $\log K^{\text{cond}}$ differ between the applications. In this study, binding of two added ligands in three different common applications to three known types of natural binding ligands is compared. The applications are (1) salicylaldoxime (SA) at 25 μM (SA25) and short waiting time, (2) SA at 5 μM (SA5), and (3) 2-(2-thiazolylazo)- ρ -cresol (TAC) at 10 μM , the latter two with overnight equilibration. The three applications were calibrated under the same conditions, although having different pH values, resulting in the detection window centers (D) $\text{DTAC} > \text{DSA25} \geq \text{SA5}$ (as $\log D$ values with respect to Fe^{3+} : $12.3 > 11.2 \geq 11$).

For the model ligands, there is no common trend in the results of $\log K^{\text{cond}}$. The values have a considerable spread, which indicates that the error in $\log K^{\text{cond}}$ is large. The ligand concentrations of the nonhumic model ligands are overestimated by SA25, which we attribute to the lack of equilibrium between Fe-SA species in the SA25 application. The application TAC more often underestimated the ligand concentrations and the application SA5 over- and underestimated the ligand concentration. The extent of overestimation and un-

derestimation differed per model ligand, and the three applications showed the same trend between the nonhumic model ligands, especially for SA5 and SA25. The estimated ligand concentrations for the humic and fulvic acids differed approximately 2-fold between TAC and SA5 and another factor of 2 between SA5 and SA25.

The use of SA above 5 μM suffers from the formation of the species $\text{Fe}(\text{SA})_x$ ($x > 1$) that is not electro-active as already suggested by Abualhaija and van den Berg (2014). Moreover, we found that the reaction between the electro-active and non-electro-active species is probably irreversible. This undermines the assumption of the CLE principle, causes overestimation of $[L]$ and could result in a false distinction into more than one ligand group.

For future electrochemical work it is recommended to take the above limitations of the applications into account. Overall, the uncertainties arising from the CLE-AdCSV approach mean we need to search for new ways to determine the organic complexation of Fe in seawater.

1 Introduction

The trace element Fe is an important micro-nutrient for phytoplankton (De Baar and La Roche, 2003; Achterberg et al., 2018; Lauderdale et al., 2020). Together with light it limits the growth of phytoplankton in 30 % to 40 % of the oceans (De Baar, 1990; Martin et al., 1990; Rijkenberg et al., 2018;

Boyd et al., 2000). One of the reasons for the limiting role of Fe is its low solubility in seawater, which can be enlarged at least 10-fold by complexation with dissolved organic ligands (Liu and Millero, 2002). The organic complexes of dissolved iron (DFe) in the oceans are important since these decrease the inorganic Fe concentration and therefore reduce precipitation as Fe(oxy)hydroxides and adsorption onto particles (scavenging). Organic ligands can also be oxidized under the influence of light and reduce Fe(III) into the labile but more bio-available Fe(II) via ligand–metal charge transfer reactions (Barbeau et al., 2001; Barbeau, 2006; Rijkenberg et al. 2006). Furthermore, organic complexation of Fe can be expected to modify Fe bioavailability, although the relationship between DFe speciation and bioavailability appears to be complex (van den Berg, 1995; Hutchins et al., 1999; Shaked et al., 2005, 2020; Salmon et al., 2006; Morrissey and Bowler, 2012; Gledhill and Buck, 2012). The significance of Fe speciation to its biogeochemistry has led to incorporation of chemical Fe speciation into global biogeochemical models, with varying levels of complexity (Tagliabue and Völker, 2011, 2015; Ye and Völker, 2017). Recent modeling work has also highlighted the potential importance of the physicochemical environment on Fe speciation, in particular highlighting the role that pH plays in modifying Fe speciation (Ye et al., 2020). The role of both pH and temperature is potentially of great significance considering climate change and ocean acidification.

Out of the natural ligand pool, the following Fe-binding organic ligands groups have been identified:

1. siderophores, relatively strong Fe-binding ligands excreted by micro-organisms to bind Fe and make it bio-available (Gledhill et al., 2004; Mawji et al., 2008, 2011; Boiteau and Repeta, 2015; Boiteau et al., 2018);
2. humic substances (HSs), a diverse group of large molecules that include ligands with affinity for iron in a broad range possibly spanning from weak to as strong as some siderophores (Laglera et al., 2007, 2011, 2019; Su et al., 2018; Slagter et al., 2019);
3. polysaccharides, a group of ligands binding Fe relatively weakly (Hassler et al., 2011, 2015), although stronger polysaccharides have also been reported (Norman et al. 2015).

Organic ligands increase the solubility and residence time of Fe. Although specific methods exist that focus on analyzing siderophores, humic materials and polysaccharides, the connection between the actual abundance of these groups and the overall Fe-binding capacity is often not well resolved. A few exceptions include Laglera et al. (2019), who determined specifically humic Fe-binding ligands; Boiteau et al. (2018), who focused on DFe bound to siderophores; and Bundy et al. (2014, 2015), who combined methods to determine the abundance of specific groups.

For approximately 3 decades, competitive ligand exchange–adsorptive cathodic stripping voltammetry (CLE-AdCSV) has been used to estimate the overall Fe-binding capacity of organic matter in seawater. The application of this method enlarged our knowledge on the marine chemistry of Fe, and the results formed the base of the explanation why DFe depth profiles deviated from those of other trace metals (van den Berg, 1995; Rue Bruland, 1995; Hutchins et al., 1999; Croot et al., 2004; Laglera and van den Berg, 2009; Gledhill and Buck, 2012; Bundy et al., 2014; Buck et al., 2015; Hassler et al., 2019; Lauderdale et al., 2020). The technique estimates the conditional concentration of ligands in the sample ($[L]$) and the conditional stability constant ($K_{\text{FeL}}^{\text{cond}}$) of their complexes without specifying the different contributions of specific ligands (Gledhill and van den Berg, 1994; Rue Bruland, 1995; Wu and Luther, 1995; Croot Johansson, 2000; Boye et al., 2001; Buck et al., 2007; Cabanes et al., 2020; Ardiningsih et al., 2020). The term “conditional” is extremely important and means that the obtained results are specific to the composition of the sample matrix analyzed (DFe, temperature, pH, ionic strength). Since $[L]$ also depends on the conditions like pH, salinity and dissolved organic matter, we will use the term conditional for both parameters. The results cannot therefore be considered as an absolute quantification of the properties of all the available Fe-binding sites and extrapolated to other conditions or matrices (Gledhill and Gerringa, 2017; Town and van Leeuwen, 2014). The technique uses an added ligand (AL) with known concentration and conditional stability constant with Fe that form an electro-active Fe–ligand complex that competes with the natural ligands present in a sample for Fe. The sample is equilibrated for a defined time period under controlled conditions of pH, light and temperature. The Fe bound to the artificial ligand is analyzed through its electro-active properties. By adding increasing amounts of Fe to subsamples, the competing natural organic ligands are titrated until the natural binding sites are no longer strong or abundant enough to compete successfully with the AL. The competition is reflected by the increased proportion of Fe bound to the artificial ligand, and from this the conditional concentration and binding strength can be calculated (van den Berg, 1982). Although the method does not provide information on the molecular composition of the binding sites, CLE-AdCSV does give information on the Fe binding capacity of seawater at the measurement pH, temperature and DFe concentration of the sample. Thus, an indication of the potential capacity for further Fe binding in a particular sample can be assessed (van den Berg, 1995; Tagliabue and Völker, 2011; Pham and Ito, 2019). Application of CLE-AdCSV allowed in some samples the division of the overall ligand in two broad ligand groups as a function of their conditional stability constants indicated with 1, ($K_{1,\text{FeL}}^{\text{cond}}$), for the relatively strong ligand group and with 2, ($K_{2,\text{FeL}}^{\text{cond}}$), for the relatively weak ligand group (Rue

and Bruland, 1995, 1997; Buck et al., 2015, 2018; Bundy et al., 2014, 2015).

Four different ALs have been reported as forming effective electro-active complexes for the purposes of CLE-AdCSV: 1-nitroso-2-naphthol (NN) (Gledhill and van den Berg, 1994), salicylaldehyde (SA) (Rue and Bruland, 1995), TAC 2-(2-thiazolylazo)-*p*-cresol (Croot and Johansson, 2000) and 2,3-dihydroxynaphthalen (DHN) (van den Berg, 2006). The two ALs, SA and TAC, are the usual selection in field studies (Rue and Bruland, 1995, 1997; Croot and Johansson, 2000; Croot et al., 2004; Boye et al., 2005; Thuróczy et al., 2011a, b, 2012; Kondo et al., 2012; Bundy et al., 2014, 2015; Buck et al., 2015, 2018; Gerringa et al., 2015, 2017; Abualhaija et al., 2015; Kleint et al., 2016; Slagter et al., 2017, 2019), and basin-scale data sets now exist for $K_{\text{FeL}}^{\text{cond}}$ and $[L]$ obtained using these two ALs (Caprara et al., 2016; Cabanes et al., 2020; Schlitzer et al., 2018), which provide an important resource for our understanding of iron biogeochemistry in the ocean (Boyd and Ellwood, 2010; Boyd and Tagliabue, 2015; Völker and Tagliabue, 2015; Tagliabue et al., 2016; Lauderdale et al., 2020). However, results of inter-comparisons of field data suggest that although trends may be similar for different ALs, the different methods may not be directly inter-comparable as conditional $[L]$ differed significantly, with SA giving higher $[L]$ and often identifying more than one ligand group compared with TAC (Buck et al., 2012, 2015). With SA, often two ligand groups can be distinguished, while TAC distinguishes only one, complicating comparison of trends in $K_{\text{FeL}}^{\text{cond}}$. The $K_{\text{FeL}}^{\text{cond}}$ obtained by TAC is in between the two $K_{\text{FeL}}^{\text{cond}}$ values of the two groups obtained with SA. The question is therefore – what is the underlying cause of these differences? It was found to be urgent within the SCOR work group 139 to test or calibrate methods with model ligands. Although there are studies that determined $[L]$ and $K_{\text{FeL}}^{\text{cond}}$ for model ligands such as siderophores, with some success with respect to $[L]$ at least (Rue and Bruland, 1995; Buck et al., 2000; Witter et al., 2000; Croot and Johansson, 2000), a thorough examination of multiple ligands and approaches that also sought to compare determined $[L]$ and $K_{\text{FeL}}^{\text{cond}}$ with values calculated from thermodynamic constants has not been previously undertaken to our knowledge. In this work we chose to examine potential bias between ALs via a series of carefully controlled studies of selected Fe-binding ligands that are likely representative of those found in the marine environment. We chose to work with SA and TAC and further compared two reported SA methods. Our three approaches comprised (a) 10 μM TAC with overnight equilibration at pH = 8.05 (Croot and Johansson, 2000), (b) 25 μM SA (SA25) at pH = 8.2 with a short waiting time of 15 min for the competing reaction to occur (Rue and Bruland, 1995; Buck et al., 2007), and (c) 5 μM SA (SA5) at pH = 8.2 with overnight equilibration as described by Abualhaija et al. (2015). Since all parameters derived in CLE-AdCSV are fundamentally dependent on the side reaction coefficient of the Fe-AL under the conditions of analysis,

we calibrated each ligand in the same laboratory under comparable conditions for consistency and to avoid any issues of bias relating to the choice of calibrating ligand, the calculation methods employed in the original papers and the choice of side reaction coefficient for Fe. We made some (arbitrary) choices on conditional binding constants between DTPA and Fe; however, we worked with one set of thermodynamic constants to make comparison between the methods consistent. We therefore press the point that the focus of the paper is on comparing the empirical outcome of the three applications and not on the accuracy of $K_{\text{FeL}}^{\text{cond}}$.

We used diethylenetriaminepentaacetic acid (DTPA) as a simple well-defined model molecule; the naturally occurring phytic acid; the hydroxamate siderophores desferrioxamine *B*, ferrioxamine *E* and ferrichrome; the catecholates siderophore vibriobactin; and fulvic and humic acids. Moreover, we carried out a specific study on the kinetics of complex formation and ligand exchange of $\text{Fe}(\text{SA})_x$ complexes for the first time. All the experiments were performed in a single laboratory in order to minimize inter-laboratory variations in protocol, material and reagent variations. We begin with a short review of previous criticisms of the CLE-AdCSV approach, since this provides important context for our study. Our overall aim was to shed light on the processes that lead to method discrepancies in the determination of natural iron ligand concentrations.

2 Potential origins of bias in the determination of binding parameters by CLE-AdCSV

CLE-AdCSV is based on many limitations and assumptions which have been discussed at some length in the literature (e.g., Apte et al., 1988; Turoczy and Sherwood, 1997; Town and Filella, 2000; Hudson et al., 2003; Croot and Heller, 2012; Laglera et al., 2013; Town and van Leeuwen, 2014; Gerringa et al., 2014; Laglera and Filella, 2015; Pižeta et al., 2015; Turner et al., 2016; Gledhill and Gerringa, 2017). If the assumptions are sufficiently satisfied, the calculation of ligand complexation parameters like the conditional ligand concentration $[L]$ and the conditional stability constant $\log K_{\text{FeL}}^{\text{cond}}$ can be undertaken, usually using the Langmuir isotherm (e.g., Gerringa et al., 2014, and references herein).

Here we give a brief overview of the limitations and assumptions.

1. Thermodynamic equilibrium between Fe, added ligand and natural ligands must be established. Failure to achieve equilibrium can lead to incorrect estimates of $[L]$ and the conditional constants (Hudson, 1998; Gerringa et al., 2014; Town and van Leeuwen, 2014; Laglera and Filella, 2015). Non-equilibrium conditions arise if the electro-active complex is a reaction intermediate, if insufficient time is allowed for equilibration of the reactants or if there is a fraction of DFe that is ki-

netically inert in the timescale of the equilibrium period (e.g., aged inorganic colloids or Fe(AL) complexes).

- There must be a detectable level of competition between the added and natural ligands. The competitive interaction is summarized by the side reaction coefficients for the natural and added ligands (α_{FeL} , α_{FeAL} , respectively). The side reaction coefficient, which is often expressed as a logarithm, is defined as

$$\alpha_{\text{FeL}} = K_{\text{FeL}}^{\text{cond}} \times [L'] = \frac{\text{FeL}}{\text{Fe}'} \quad (1)$$

or

$$\alpha_{\text{FeAL}} = K_{\text{FeAL}}^{\text{cond}} \times [\text{AL}'] = D, \quad (2)$$

where $[L']$ and $[\text{AL}']$ are the conditional concentration of the organic ligand not bound by Fe and the concentration of AL not bound by Fe, respectively, and Fe' is the Fe concentration not bound to L . The side reaction coefficient of AL defines the center of the detection window or analytical window, which we defined here as D to prevent confusion between side reaction coefficients of added and natural ligands. The window is assumed to be approximately 3 to 3 orders of magnitude wide and 1 to 2 orders of magnitude above and below D (Apte et al., 1988; van den Berg and Donat, 1992; Milller and Bruland, 1997; Laglera et al., 2013; Laglera and Fillela, 2015). In practice, the upper limit of D is defined by the analytical sensitivity of the AdCSV method, as it is bound by the limit of detection of FeAL. The lower limit of D is bound by ligands that are outcompeted by the AL within the range of Fe added during the titration. Since AdCSV is internally calibrated via standard additions, in practice the lower limit of the detection window is bound by the value of α_{FeL} achieved when the analytical response is deemed to be linear (Apte et al., 1988; Laglera and Fillela, 2015).

- The concentration of the FeAL complex can be accurately determined at each titration point. AdCSV is internally calibrated via standard additions, and D and values obtained for $K_{\text{FeL}}^{\text{cond}}$ and $[L]$ are strongly influenced by our ability to accurately calculate the sensitivity (Turoczy and Sherwood, 1997; Hudson et al., 2003; Pizeta et al., 2015).
- Complexes of Fe with natural ligands cannot be electro-labile under the experimental conditions since this could result in interferences with the actual detection of the Fe–AL complex (Yang and van den Berg, 2009; Laglera et al., 2011).
- The equilibrated Fe–AL complex must be electro-active since it is the reaction on which the detection is based.
- The AL should not react with the natural ligands altering or canceling their binding ability.

In the last decade, many studies have questioned the compliance of the CLE-AdCSV methodology to these assumptions and their influence on method discrepancies. Laglera et al. (2011) showed the inability of TAC to measure fulvic and humic acids as Fe-binding dissolved organic ligands, which might be due to either assumption 2 or 6. Humic substances are ubiquitous, they form large diverse molecules and they are broadly recognized as Fe-binding ligands (Krachler et al., 2015; Su et al., 2018; Laglera et al., 2019; Whitby et al., 2020; Yamashita et al., 2020). According to other work, TAC is able to detect at least some humics as Fe-binding organic ligands (Batchelli et al., 2010; Slagter et al., 2017; Dulaquais et al., 2018).

The SA25 application has been criticized for not meeting assumption 1; SA25 has a waiting time of 15 to 20 min in contrast with the overnight equilibration used for the TAC and SA5 applications (Abualhaija and van den Berg, 2014; Abualhaija et al., 2015; Slagter et al., 2019).

Abualhaija and van den Berg (2014) found that two Fe–SA complexes are formed, FeSA and Fe(SA)₂, and only FeSA is electro-active. At higher [SA], the proportion of Fe(SA)₂ increases and the analytical signal decreases, resulting in a negative relationship between sensitivity and competitive force. Finally, we would like to point out that the pH of the analysis may have a larger influence on the organic complexation of DFe than previously thought (Gledhill et al., 2015; Avendaño et al., 2016; Ye et al., 2020), but the same competition of OH ions in binding Fe, irrespective of the buffered pH values of the SA and TAC applications, is sometimes used (8.2 and 8.05, respectively, which is a factor of 1.4 different in terms of H⁺ concentration). This complicates a direct comparison of data even more.

3 Methods

The natural seawater used in the experiments consisted of mixed leftover filtered samples of the northwestern Atlantic GEOTRACES cruise GA02 (stored frozen at -20°C) (Rijkenberg et al., 2014). A sample volume, assumed to be necessary for the following few days, was thawed, mixed, UV-irradiated to destroy the natural organic Fe-binding ligands and stored in the refrigerator. Consequently, one batch differs from others with respect to the DFe content, and also potentially in other constituents, such as other trace metals. Since surface samples were not used, we do not expect large differences in salinity. The average salinity was 35.09 ± 0.61 ($N = 434$), obtained as an average of all samples > 100 m depth taken for the ligand analysis in Gerringa et al. (2015). Samples for DFe analysis were taken from every batch. UV-irradiated seawater was stored for 3 d at most.

3.1 Equipment and measuring conditions

3.1.1 Equipment and electrochemical parameters

We carefully followed procedures as described in the literature to ensure methodology was as close as possible to that originally described (Tables S1 and S2). Three different voltammetric setups were used (Table S1). A standard Metrohm set up was used for TAC. For use with SA, a separate Metrohm system was modified to allow for air purging whilst the mercury drop formation was still executed under nitrogen pressure. Nitrogen did not leak into the headspace of the sample during the measurements in our Metrohm stand. However, when drops are formed pulses of nitrogen are released and end up in the headspace of the sample, and purging with air would remove (at least part of) the nitrogen. To check a potential effect of this nitrogen, a kinetic experiment with SA25 was executed. To five identical subsamples SA was added at the same time. These subsamples were each measured repetitively during 1 h to several hours, one after the other. No effect was seen after subsample replacements (Fig. S1). We concluded that nitrogen from the stand did not influence the kinetic process, since the measured FeAL concentrations had a gradual change over time, independent of the subsample. For the other kinetic experiments with SA, BASi equipment was used (Table S1). The electrochemical settings used by Croot and Johansson (2000), Buck et al. (2007), and Abualhaija and van den Berg (2014) were used without alteration and are summarized in Table S2.

3.1.2 Conditioning and equilibration

Electro-active complexes with Fe typically have low solubility and thus tend to adsorb on the walls of containers. Conditioning and equilibration of all contact surfaces is thus an important pre-treatment step in order to minimize losses of Fe and ligand species during the course of the experiment. Different materials do not have the same adsorption properties (Fischer et al., 2006). All cells and titration vials were made of Teflon. Other bottles, sample bottles and those used for kinetic experiments were low-density polyethylene (LDPE) bottles (Nalgene™, Fisher Scientific). For all three applications, the same materials were used, canceling any deviation among methods from the interaction of solution component and containers. Equilibration between the samples and AL was attained at room temperature.

Before use, all materials such as vials, bottles and cells were conditioned overnight in UV-irradiated seawater with the prepared combinations of seawater and ligand. The conditioning procedure was performed at least three times for the analysis with TAC and at least five times for analysis with SA. The cell with electrodes, stirrer and purge tube were kept overnight in low-metal seawater. Before a titration was started, first two measurements were executed with seawater containing all chemicals but no Fe addition. These mea-

surements also served as check for possible contamination of the cell. Hereafter, two zero additions were measured (see Sect. 3.4), of which the second was used as the start of the titration.

Before starting kinetic measurements, a 30 mL vial with the same content and treatment as the sample was used for three analyses (thus three times 10 mL) in order to condition the cell wall, electrodes and stirrer.

The 200 mL bottles used for kinetic studies were conditioned with 6 nM Fe, in the absence of the added ligand. For tests with UV-irradiated seawater without a model ligand, 200 mL bottles were conditioned by rinsing the bottle three times for 2 min with 30 mL of the test seawater. Since UV-irradiated seawater did not contain Fe-binding organic ligands, most of the added 6 nM DFe would adsorb on the bottle walls or precipitate.

Samples were equilibrated according to the specific method descriptions, which was overnight equilibration for the TAC and 5 μ M SA and 15 min for 25 μ M SA (Croot and Johansson, 2000; Buck et al., 2007; Abualhaija and van den Berg, 2014). The 15 min equilibrations were applied precisely using a stopwatch, whereas overnight equilibration resulted in a period of at least 14 h.

3.1.3 UV irradiation

Samples, without any additions, were poured into 30 mL Nalgene FEP bottles and placed in a custom-made UV box between 4 TUV 15W/G15 T8 fluorescent tubes (Phillips) for 4 h (Rapp et al., 2017; Wuttig et al., 2019). Precipitates were not observed. After UV irradiation, samples were transferred into a clean 1 L trace-metal LDPE bottle and kept in the refrigerator.

3.1.4 Model ligands

The following discrete synthetic ligands of known concentration (model A ligands) were used at a concentration of 2 nM, unless otherwise stated. No tests were undertaken to check the purity of the siderophores. The solutions were used within 2 weeks after preparation and kept in the refrigerator in the dark at 4 °C, which should at least for DFOB be short enough to prevent degradation (Hayes et al., 1994). The aim of our research was to compare the three applications. Humics (model B ligands, 0.1 or 0.2 mg/L) were added in a concentration to give an iron-binding capacity of approximately 3 nM (Laglera and van den Berg, 2009; Yang et al., 2017; Sukekava et al., 2018). The stoichiometry of the formed Fe-model ligand complexes differs for each model ligand. In order to simplify the comparison of binding strengths, stability constants are given for a 1 : 1 stoichiometry.

Model A ligands

- Diethylenetriaminepentaacetic acid (DTPA C₁₄H₂₃N₃O₁₀, Sigma-Aldrich D6518-5G) was

used to calibrate the added ligands via reverse titration according to methods described previously (Croot and Johansson, 2000). We calculated a conditional binding constant $\log K_{\text{FeDTPA,Fe}^{3+}}^{\text{cond}}$ of 19.0 using the ion-pairing speciation software Visual MINTEQ (Gustafsson, 2012), disregarding the formation of FeOHDTPA. The $\log K_{\text{FeDTPA,Fe}^{3+}}^{\text{cond}}$ value was independent of the pH difference 8.05–8.2. This is 0.34 higher than the value (18.65) used by Croot and Johansson (2000) at $I = 0.7$, pH = 8.05, with the difference most likely arising as a result of the lower ionic strength predicted by the ion-pairing model. Addition of 2 and 4 nM DTPA did not increase the DFe content of the UV-irradiated seawater (detection limit = 25 pM)

- Phytic acid ($\text{C}_6\text{H}_{18}\text{O}_{24}\text{P}_6 \times x\text{Na}^+ \times y\text{H}_2\text{O}$, Sigma 68388). According to Witter et al. (2000), $\log K_{\text{FePA,Fe}^{3+}}^{\text{cond}} = 22.3\text{--}22.4$. Rijkenberg et al. (2006) warned that at high phytic acid concentrations aggregates are formed, but at our concentrations (2 nM) this should not be a problem.
- The hydroxamate siderophore desferrioxamine *B* ($\text{C}_{25}\text{H}_{48}\text{N}_6\text{O}_8$, CH_4SO_3 Novartis RVG03984 U.R., 477881 NL.) has a thermodynamic stability constant of 30.5 ($I = 0.1$; Hider and Kong, 2010). According to Witter et al. (2000) the conditional stability constant, $\log K_{\text{FeDFOB,Fe}^{3+}}^{\text{cond}}$, is between 21.6 and 22.1 ($I = 0.7$). Van den Berg (2006) found $\log K_{\text{FeDFOB,Fe}^{3+}}^{\text{cond}} = 21.5$, whereas Croot and Johansson (2000) concluded that this conditional stability constant was too high and outside *D* of their TAC method ($\log K_{\text{FeDFOB,Fe}^{3+}}^{\text{cond}} > 23.4$). However, new side reaction coefficients of major cations have been determined since, which give rise to a $\log K_{\text{FeDFOB,Fe}^{3+}}^{\text{cond}}$ of 24.3 at seawater salinity (Schijf and Burns, 2016).
- The hydroxamate siderophore ferrichrome ($\text{C}_{27}\text{H}_{45}\text{N}_9\text{O}_{12}$ ferrichrome iron-free from *Ustilago sphaerogena*, Sigma Aldrich (F8014-1MG)). Hider and Kong (2010) gave a $\log K_{\text{FeL,Fe}^{3+}} = 29.1$ ($I = 0.1$). According to Witter et al. (2000), $\log K_{\text{FeL,Fe}^{3+}}^{\text{cond}}$ in seawater varies between 21.6 and 22.9 depending on the applied method. Kinetic measurements determining formation constant resulted in 22.9; the equilibrium approach with Fe titration resulted in 21.6.
- The hydroxamate siderophore ferrioxamine *E*, ($\text{C}_{27}\text{H}_{45}\text{FeN}_6\text{O}_9$ ferrioxamine *E* from *Streptomyces antibioticus*, Sigma Aldrich (38266-3MG-F)). According to Hider and Kong (2010) ferrioxamine *E* has a higher affinity for Fe than ferrioxamine *B* ($\log K_{\text{FeL,Fe}^{3+}} = 32.5$, at $I = 0.1$). But Bundy et al. (2018) estimated $\log K_{\text{FeL,Fe}^{3+}}$ values of ferrioxamine *B* and *E*, using SA5 in seawater, to be close with 14.4

and 14, respectively. However, this model ligand as purchased was already saturated with Fe.

- The triscatecholate siderophore vibriobactin ($\text{C}_{35}\text{H}_{33}\text{FeN}_5\text{O}_{11}$ vibriobactin (iron-free) from *Vibrio cholerae* V69, EMC micro collections). No information is available on the Fe-binding characteristics of this model ligand, but in general catecholates have higher binding strengths with Fe than hydroxamates because of their ortho phenolate binding groups (Hider and Kong, 2010).

Model B ligands

Humic substances are the heterogeneous mix of hydrophobic compounds originating from chemical and microbial transformation of living matter as it decays in the environment.

- Fulvic acid (FA) is the smaller and more soluble fraction of humic substances (Buffle, 1990). Therefore, this is not just a ligand but also a series of compounds of which a fraction function as iron-binding ligands. Laglera and van den Berg (2009) determined that 1 mg of this specific FA binds 16.7 ± 2.0 nM Fe with $K_{\text{FeL,Fe}^{3+}}^{\text{cond}} = 10.6$. (IHSS Suwannee River Fulvic Acid Standard II, 1R101F). Yang et al. (2017) and Sukekava et al. (2018) found that 1 mg could bind 14.6 ± 0.7 nM Fe. (IHSS Suwannee River Fulvic Acid Standard II 2S101F). It must be noted that the batches are different between the above results, and these values can differ per produced batch. Still we assumed 2.92 nM equivalents (nMEq) of ligand sites to be added with 0.2 mg SRFA per liter.
- Humic acids (HAs) are the larger fraction of humic substances that precipitate at low pH (pH 2) (Buffle, 1990). Laglera and van den Berg (2009) determined that 1 mg of this specific HA binds 32 ± 2.2 nmol Fe with $\log K_{\text{FeL}}^{\text{cond}} = 11.1$. (IHSS Suwannee River Humic Acid Standard II, 2S101H). We assumed that 0.1 mg of added HA per liter would add 3.2 and 0.2 mg 6.4 nM Eq of ligand sites.

3.2 AL calibration

Seven or eight conditioned Teflon 30 mL vials were filled with 10 mL of UV-irradiated seawater spiked with buffer, 6 nM Fe (Table S1) and increasing amounts of the calibrating ligand DTPA. For TAC the pH was 8.05, and for SA the pH was 8.2 according to the original method specifications (according to the NSB scale). The calibrations were repeated four times. The buffer used for all applications was ammonium borate (Abualhaija and van den Berg, 2014; Buck et al., 2007). Details can be found in the Supplement.

DTPA additions were 0, 10, 100, 200, 400, 1000 and 2500 nM DTPA for TAC; 0, 1, 10, 40, 80, 100 and 200 nM DTPA for SA5; and 0, 1, 40, 100, 200, 400 and 1000 nM DTPA for SA25. Mixtures of UV seawater with buffer, DFe

and DTPA were equilibrated for at least 8 h, after which SA or TAC was added. New mixtures were equilibrated either overnight or for 15 min in the case of SA25, after which peak heights for FeAL were determined following the procedures described in Table S1. We calibrated SA25 using a short waiting time instead of 5 h of equilibration (Rue and Bruland, 1995; Buck et al., 2007) to ensure consistency with the approach applied to samples. For the calibration of TAC the normal Metrohm instrument was used, and for SA the BASi instrument was used (Tables S1, S2). The measurements were done in sequence of increasing DTPA concentrations, without rinsing cells in between.

3.3 Signal stability tests

As we found the CSV signal after SA addition lacked stability and decreased with time, we performed a series of experiments to find the cause. We tested the following aspects for both instruments: the influence of a purge step with air (Fig. S2), the influence of the size of the mercury drop (Fig. S3, Table S3), the influence of the mercury puddle on the bottom of the cell, and the influence of the SA concentration. For the kinetic measurements (Sect. 2.4) of the SA applications, both BASi and Metrohm stands were used. Details on procedures are given in the Supplement.

3.4 Titrations

3.4.1 TAC

Fifteen vials were prepared with increasing Fe content, in a mixture of UV-irradiated seawater and model ligand (Table S1, Croot and Johannsson, 2000; Ardiningsih et al., 2020). Blanks were obtained by analysis in the absence of model ligands.

3.4.2 SA5

The application followed Abualhaija and van den Berg (2014) but used the above-described BASi instrument. SA (added to a final concentration of 5 μM), buffer, Fe additions (Table S1) and samples were left to equilibrate overnight.

3.4.3 SA25

For SA25, the buffer and DFe were added 1 h before analysis. SA was added to a final concentration of 25 μM separately to each vial, 15 min before the measurement.

3.5 Kinetic measurements

The samples contained either UV-irradiated seawater or UV-irradiated seawater with a model ligand to which buffer and 6 nM Fe were added in a pre-conditioned bottle. If a model ligand was present, this was first allowed to equilibrate overnight with the buffer and 6 nM Fe. In samples with

only UV-irradiated seawater, two approaches were followed: one in which Fe was added together with TAC or SA at $t = 0$ and one in which Fe was equilibrated overnight prior to addition of TAC or SA. In the latter case, there is the possibility that Fe-oxide precipitates were formed prior to the addition of TAC or SA and were probably dissolved after the addition of TAC or SA. At $t = 0$, TAC or SA was added.

At $t = 0$, the first measurements were done as rapidly as possible until approximately $t = 1$ h, followed by subsequent measurements every 20 min, every 40 min and 1 h until either $t = 4$ or $t = 7$ h. The number of analyses depended on the application and experiment duration (4 or 7 h) but contained a minimum of 14 duplicate measurements. In this way the FeAL_(x) formation in time can be followed. However, the model ligand dissociation rates cannot be calculated from the rate of peak increment because the addition of Fe (6 nM) was in excess of the model ligand concentration (2 nM Eq, if not indicated differently). Therefore, the excess Fe formed hydroxides and adsorbed on the cell and electrode surfaces. The increment of signal reflects the competition of TAC with all these iron species and not just with the model ligand.

Two protocols were followed (Fig. S4).

1. *In-cell experiments.* With repeated scans of the same sample contamination was prevented, and more measurements could be undertaken, especially at the start of the experiments. The total time of the experiments lasted 4 or 7 h. Samples of 30 mL were prepared, of which the first 20 mL was used to condition the cell twice, after which the last 10 mL was transferred to the cell and the experiment undertaken. The AL was added to the cell at $t = 0$. For TAC, the addition took place after the purge step to reduce the time lapse between addition and first measurement. An extra set of in-cell experiments were carried out with UV-irradiated seawater, natural seawater and UV-irradiated seawater spiked with DTPA, 40 nM for SA = AL and 200 nM for TAC = AL. Other experiments for all model ligands were repeated with 2 nM of added model ligand. In this protocol, mercury accumulated in the cell during the experiment.
2. *Bottle experiments.* Scans were carried out on separate aliquots of one sample. In this experiment, a fresh aliquot of 10 mL was pipetted into the voltammetric cell for the determination of peak height at each time point. The total sample volume was 200 mL, and the AL was added at $t = 0$. In this experiment, accumulation of mercury at the bottom of the cell was limited. The experiment lasted for 4 or 7 h, consistent with the in-cell approach. The first 10 mL was transferred as quickly as possible into the preconditioned cell and the measurement started. In order to determine the amount of adsorbed Fe on the 250 mL bottle walls, the bottles were rinsed carefully with 5 mL of elution acid (1.5 M Teflon distilled HNO₃ that contained rhodium; see be-

low Sect. 3.6 ICPMS analysis) and Fe concentration in the acid rinse determined by inductively coupled plasma mass spectrometry (ICPMS) (Sect. 3.7).

3.6 Calculations

The sensitivity, S ; the ligand concentration, $[L]$; and the conditional stability constant (K^{cond}) were calculated by direct non-linear fitting of the Langmuir isotherm (Gerringa et al., 2014) with inherent co-dependence of $[L]$ and K^{cond} (Apte et al., 1988; Hudson et al., 2003; Gerringa et al., 2014).

The inorganic side reactions of DFe with dissolved hydroxides, $\alpha_{\text{Fe}^{\cdot}}$, were calculated using the constants from Liu and Millero (2002), resulting in an inorganic alpha for Fe (α_{inorg}) of $\log\alpha_{\text{inorg}} = 9.9$ at pH = 8.05 and $\log\alpha_{\text{inorg}} = 10.4$ at pH = 8.2. These are slightly different from literature values of Croot and Johansson (2000) and Abualhaija and van den Berg (2014). The conditional binding strength of DTPA was obtained using Visual MINTEQ. We used an average seawater major ion composition, and an average deep sea DFe concentration of 0.5 nM was chosen for these calculations. DTPA was added to the composition at the concentrations used, and the pH was fixed at values of 8.05 and 8.2. According to the Visual MINTEQ calculations, $K_{\text{FeDTPA, Fe}^{3+}} = 27.3$, the logarithm of side reaction coefficients for DTPA with major cations was 8.26, resulting in $K_{\text{FeDTPA, Fe}^{3+}}^{\text{cond}} = 19.0$.

3.7 ICPMS analysis of dissolved Fe

Dissolved Fe was analyzed with a Thermo Finnigan HR-ICPMS element 2 (for details see Middag et al., 2015; Gerringa et al., 2020). Briefly, seawater aliquots, with and without the addition of model ligands, were concentrated using a seaFAST system after UV destruction.

For the analyses of DFe in UV-irradiated seawater with and without added model ligands, the limit of detection for DFe was $22 \text{ pM} \pm 8 \text{ pM}$. The DFe of model ligands DTPA, phytic acid, desferrioxamine *B* and ferrichrome was (2 and 4 nM) below the detection limit in the dilutions used. This means the addition of the model ligand did not increase the DFe in the UV-irradiated seawater. Analyses of 2 nM vibriobactin resulted in a value under the detection limit once and in 0.1 nM DFe once. However, ferrioxamine *E*, FA and HA contained measurable amounts of DFe: 2 nM ferrioxamine *E* = $1.76 \pm 0.04 \text{ nM}$ ($N = 2$), 0.2 mg FA = $1.1 \pm 0.02 \text{ nM}$ ($N = 3$) and 0.2 mg HA = $3.39 \pm 0.05 \text{ nM}$ ($N = 2$).

Background Fe concentrations in TAC, SA and both buffers were determined by pipetting 100 μL in 20 mL of elution acid (1.5 M Teflon distilled HNO_3 containing rhodium). These samples were measured by ICPMS without further sample handling as were the acid rinse samples to measure adsorption on bottle walls. The results of the ICPMS on samples with added ligands are given in Table 2 as the DFe of the samples. Upon addition of the buffers, 0.04 nM

DFe was added inadvertently to the samples. The addition of 10 μM TAC added 0.2 nM Fe, 25 μM SA 0.2 nM and 5 μM SA 0.04 nM Fe. These inadvertent additions have been included in the Fe concentrations. The acid rinse of the 250 mL LDPE bottles contained 106 nM Fe, when conditioned with 6 nM Fe and TAC and 59 nM Fe when conditioned by 6 nM Fe, 2 nM phytic acid and TAC. This means that the potential release in a 200 mL sample could be at maximum 1.3 and 0.7 nM Fe. The difference in Fe adsorption on the bottle wall shows the effect of conditioning very well.

4 Results and discussion

4.1 Calibration of Fe–AL α coefficients

Details of the calibration are given in the Supplement; here we explain our choice to use an overall α coefficient for SA as AL instead of the sum of separate α coefficients of FeSA and FeSA₂. We further present and discuss the resulting binding characteristics of the ALs.

The competition by DTPA causes a reduction in peak height compared to the situation without DTPA (Fig. S5). At equilibrium, dissolved Fe is distributed over the following species:

$$\text{DFe} = [\text{FeDTPA}] + [\text{FeAL}_2] + [\text{FeAL}] + [\text{Fe}^{\cdot}]. \quad (3)$$

For the application of TAC, the contribution of FeTAC is thought to be negligible with respect to the formation of Fe(TAC)₂ (Croot and Johansson, 2000). For SA in the micromolar range, both FeSA₂ and FeSA are formed, although only FeSA is the electro-active species (Abualhaija and van den Berg, 2014). Using Eqs. (1) and (3) gives

$$\begin{aligned} \text{DFe} &= \alpha_{\text{FeDTPA}} \times [\text{Fe}^{3+}] + \alpha_{\text{FeAL}_2} \times [\text{Fe}^{3+}] + \alpha_{\text{FeAL}} \\ &\times [\text{Fe}^{3+}] + \alpha_{\text{inorg}}[\text{Fe}^{3+}] = [\text{Fe}^{3+}] \times (\alpha_{\text{FeDTPA}} \\ &+ \alpha_{\text{FeAL}_2} + \alpha_{\text{FeAL}} + \alpha_{\text{inorg}}). \end{aligned} \quad (4)$$

The α coefficients determine the distribution of Fe over the complexes with DTPA and AL. When $[\text{FeDTPA}] = \Sigma [\text{FeAL}]$, the α coefficients of DTPA and AL are equal, illustrating that a calibration is actually comparing α values of the added ligand (AL) and the calibrating ligand. From the α values at a determined AL concentration, $K_{\text{FeAL}}^{\text{cond}}$ and/or $\beta_{\text{FeAL}_2}^{\text{cond}}$ are calculated. The calculation of $\beta_{\text{Fe(SA)}_2}^{\text{cond}}$ cannot be done with precision using only our two SA concentrations. Since we actually need the α values for calculating the ligand characteristics from the titration data, we do not need to calculate $\beta_{\text{Fe(SA)}_2}^{\text{cond}}$. The α values include the contributions of $K_{\text{FeAL}}^{\text{cond}}$ and $\beta_{\text{FeAL}_2}^{\text{cond}}$ (Table 1) (for more details see the Supplement).

The obtained α values (Table 1) differ from the original literature values, which is likely due to the toolbox we used, Visual MINTEQ. If we consider the $\log\alpha_{\text{FeAL, Fe}^{3+}}$ values, the calibration results for TAC and SA5 we obtained

Table 1. Average beta and alpha values of the added ligands (AL) with the standard deviation around the mean of N experiments. In bold are the parameters used in this study to calculate the model ligand characteristics, 1: assuming one FeAL is formed, either FeSA or Fe(SA)₂, 2: assuming both FeSA and Fe(SA)₂ are formed. (a–c) indicate literature values: ^a Croot and Johanson (2000) using $\log \alpha_{\text{inorg}} = 10$; ^b Abualhaija et al. (2015) using $\log \alpha_{\text{inorg}} = 9.98$; ^c Buck et al. (2007) using $\log \alpha_{\text{inorg}} = 10$. Alpha values of the AL are the direct outcomes of the calibration exercises; therefore, these have a standard deviation added, which is the standard deviation around the mean of four calibrations. Since K and/or β are directly derived by dividing through the AL concentration or squared concentration, the standard deviations of $\log K_{\text{FeAL,Fe}'}^{\text{cond}}$ and $\log \beta_{\text{FeAL}_2\text{,Fe}'}^{\text{cond}}$ have the same values.

AL	$\alpha_{\text{FeAL,Fe}'}$	$\log \alpha_{\text{FeAL,Fe}^{3+}}$	N	$\log \beta_{\text{FeAL}_2\text{,Fe}'}^{\text{cond}}$	$\log \beta_{\text{FeAL}_2\text{,Fe}^{3+}}^{\text{cond}}$	or	$\log K_{\text{FeAL,Fe}'}^{\text{cond}}$	$\log K_{\text{FeAL,Fe}^{3+}}^{\text{cond}}$
TAC	275 250 ^a	12.3 ± 0.2 12.4 ^a	4	12.4 12.4 ^a	22.3 22.4 ^a			
SA5	4.16 17.87 ^b	11.1 ± 0.1 11.23 ^b	4	11.3 ¹ 10.7 ^b	21.7 ¹ 20.7 ^b		5.9² 6.5 ^b	16.3² 16.5 ^b
SA25	5.49 78.7 ^c	11.3 ± 0.23 11.9 ^c	5	10.1 ¹ 11.1 ^c	20.5 ¹ 21.1 ^c		5.34²	15.8²

(Table 1) compare very well with values from the literature (Croot and Johansson, 2000; Abualhaija and van den Berg, 2014). Our $\log \alpha_{\Sigma\text{FeSA,Fe}^{3+}}$ of SA25 (considering both FeSA and Fe(SA)₂ formation) shows a larger discrepancy with Buck et al. (2007) than the above comparisons (our $\log \alpha_{\Sigma\text{FeSA,Fe}^{3+}} = 11.2$ versus $\log \alpha_{\text{FeSA}_2\text{,Fe}^{3+}} = 11.9$ of Buck et al. (2007)).

The difference becomes larger when calculated with respect to inorganic Fe (Fe') when using the pH-adjusted values of $\log \alpha_{\text{inorg}} = 9.9$ for 8.05 and $\log \alpha_{\text{inorg}} = 10.4$ for pH = 8.2 (Liu and Millero, 2002). For SA5 and SA25, the comparison between our data and literature values is thus offset with respect to Fe', due to the application of $\log \alpha_{\text{inorg}} = 10.4$. It is possible that the larger deviation in SA25 from previously reported values is partly due to the shorter waiting time used in our study, 15 min instead of 5 h (Rue and Bruland, 1995). However, the calibration should be executed according to the published protocol of the analyses.

4.2 Titrations

We present the concentrations of FeAL determined during the titrations of the selected ligands with the three different methods to allow direct comparison between the approaches (Fig. 1, Tables 2 and 3). Differences due to variations in sample materials are assumed to be small. However, a variance in the content of metals that could compete with Fe for ligand sites can have influenced the results and might have caused an underestimation of the model ligand concentration and indirectly also have influenced the value of K^{cond} . This could not have influenced the comparison between the applications since the same mixed sample was always used per experiment for the three applications. We again emphasize that CLE-AdCSV titrations in natural waters result in the derivation of conditional parameters, and this applies to the ligand concentration as well as the stability constant.

Overall, K^{cond} and α_{FeL} values of the model ligands were highest with TAC. In other words, they were highest with the application with the highest D . For model A ligands like siderophores, this points to bias in the determination of K^{cond} , perhaps as a result of true values too high to be measured with accuracy. For example, an estimate of 24.25 for K^{cond} for FeDFO in seawater can be calculated using the side reaction coefficient of 6.25 for DFO binding to Ca and Mg at pH 8.0 (Schijf and Burns, 2016; Wuttig et al., 2013) and the stability constant for FeDFO given by Hider and Kong (2010). For the complex ligands, model B, like humic and fulvic acids, which contain multiple binding sites with a range of affinities, an average K^{cond} will be determined based on D of the method being applied (Tables 1 and 2). Both factors highlight important concepts that relate to the CLE-AdCSV approach in general that need to be taken into consideration when interpreting K^{cond} derived from CLE-AdCSV titrations.

Ligand concentrations were highest with SA25 and lowest with TAC (Tables 2 and 3, Fig. S6) and thus showed the opposite trend to K^{cond} . Comparison with the actual added concentrations of the model A ligands shows that $[L]$ was, with only the exception of the Fe-saturated ferrioxamine E , relatively underestimated by TAC (5%–58%) and systematically overestimated by SA25 (26%–125%, Fig. S6). The overestimation by SA25 might be due to a lack of equilibrium. In theory when Fe-binding ligands are not yet in equilibrium with the AL, the dissociation of FeL complexes required to reach equilibrium is incomplete, and the so-called straight part is curved and not straight. In principle this will underestimate the ligand concentration. The overestimates observed for SA25 might therefore be caused by disequilibrium in the Fe–SA species. The extent of overestimation and underestimation differed per model ligand (see below), and the three applications showed the same trend between the model A ligands (Fig. S6), especially for SA5 and SA25. As-




Figure 1. Iron titrations of UV-irradiated seawater containing model ligands in competition with TAC, SA5 and SA25 as added ligand. (FeAL) is Fe-added ligand complex using sensitivity (S) = 1 and (Fe) is total iron concentration. See Table 2 for the DFe at zero addition. (a) DTPA, (b) phytic acid, (c) desferrioxamine B, (d) ferrichrome, (e) ferrioxamine E (saturated with Fe), (f) vibriobactin, (g) fulvic acid FA and (h) humic acid HA. Data for TAC and SA5 are from duplicate experiments, and data for SA25 are from single experiments, except for FA and HA where for SA25 duplicate experiments were also done. Note the different HA concentrations, 0.1 and 0.2 mg.

suming the concentration of ligand sites per weight unit determined by Laglera and van den Berg (2009) and Sukekava et al. (2018) to be correct, the overestimation by SA25 was larger for model B ligands (Tables 2 and 3). The difference from the average value for duplicate measurements of $[L]$ was 0.3 and 0.2 nMEq of Fe for the TAC and SA5 application, respectively (excluding ferrioxamine E because it was saturated with Fe; see below). The standard deviation with SA25 ($N = 5$) was 0.3 nMEq of Fe. In the following we will assume ± 0.3 nMEq of Fe as precision for $[L]$. The differences between the applications are smaller when the α_{FeL} values are compared (Table 3), which is understandable, since it is α_{FeL} that is titrated and also because α_{FeL} is calculated from the product of $[L]$ not bound by Fe ($[L']$) and K^{cond} and thus compensates for any codependence between K^{cond} and $[L]$.

We “back-calculated” the titration curves using our present results, K^{cond} and $[L]$, and we presented this in log-log plots of $[\text{FeAL}]$ versus total dissolved Fe together with the actual data points (Fig. 2). For DTPA and desferrioxamine B we added the theoretical titration curves that should be obtained given the K^{cond} calculated from the thermodynamic constants and the 2 nM of added model A ligands. We presented the back calculations in log-log plots in order to magnify the initial part of the titration (Fig. 2).

4.2.1 DTPA

All applications have been calibrated by reverse titration with DTPA. We would expect to recover comparable binding parameters for DTPA during the Fe titration. However, in all cases the $\log K^{\text{cond}}$ for DTPA calculated from the Fe titration was overestimated. The overestimation of $K_{\text{FeDTPA}}^{\text{cond}}$ for all three added ligands is likely a result of $\alpha_{\text{FeDTPA}} < D$ and thus theoretically below the detection window for all applications. For determinations in marine samples, Caprara et al. (2016) showed in a compilation of data from the open ocean that, with the exception of NN, the ligands were above D of the used AL; thus deviation caused by ligands with $\alpha < D$ is likely a minor problem in seawater samples. For TAC and SA5, $[L]$ was underestimated. Although such a discrepancy in $[L]$ could be a result of incorrect estimation of the Fe present in the titration, analysis with ICPMS showed that the Fe concentration increased only by 0.04 nM upon 2 nM DTPA addition, and thus we ruled out contamination as a cause for the underestimation of $[L]$ for TAC and SA5. The K^{cond} values are comparable between the applications (ratios vary between 1 and 1.03, Table 3), although the range 21.3–21.8 is substantially higher than 19.0, the K^{cond} used for the calibration. Due to the codependence, the DTPA ligand concentration (2 nM) should have been underestimated (Apte et al., 1988), which is the case for the results from

Table 2. Results of the titrations following the three applications, SA5, SA25 and TAC for model A ligands, with a well-described composition and a specific added concentration of 2 or 4 nM, and for model B ligand, the humic substances FA and HA that do not have a fixed composition and were added in weight units (0.1–0.4 mg/L). DFe was measured by ICPMS. $\log K$ is used for $\log K^{\text{cond}}$ with respect to Fe^{3+} . $\log K^{\text{cond}}$ and $[L]$ are calculated using the non-linear Langmuir isotherm. Alpha ($K^{\text{cond}} \times [L']$) is calculated using $[L']$ and not by simple $[L]$ minus DFe. DFe is in nM, $[L]$ is in nM Eq Fe and K^{cond} in M^{-1} . For the model ligands 2 nM were used unless otherwise stated. Most model ligands have been analyzed in duplicate with TAC and SA5, and once with SA25. The addition of the humics was determined using Laglera and van den Berg (2009) for HA and Sukekava et al. (2018) for FA. Since K' is log transformed, the standard error (SE) is asymmetric to lower and upper values; therefore two SE values are obtained, one to lower (down) and to upper (up) values.

Name	TAC							SA 5 μM							SA 25 μM						
	DFe	$\log K$	SE	SE	$[L]$	SE	Log	DFe	$\log K$	SE	SE	$[L]$	SE	Log	DFe	$\log K$	SE	SE	$[L]$	SE	Log
	nM		Down	Up	nMEq		$\alpha_{\text{Fe}'}$			Down	Up	nMEq		$\alpha_{\text{Fe}'}$			Down	Up	nMEq		$\alpha_{\text{Fe}'}$
Model A																					
DTPA	0.31	21.8	0.4	0.2	1.35	0.20	2.96	0.12	21.7	NA	0.4	1.15	0.09	2.29	0.25	21.3	0.3	0.2	2.62	0.15	2.23
2 nM	0.31	21.7	0.2	0.2	1.73	0.24	2.90	0.12	21.6	NA	0.6	1.14	0.21	2.16							
PhA	0.29	21.9	0.2	0.1	1.85	0.13	3.18	0.44	20.7	0.3	0.2	2.85	0.30	1.69	0.57	20.1	0.3	0.2	3.12	0.95	1.14
2 nM	0.29	22.8	NA	0.3	1.32	0.09	2.92	0.44	20.8	0.3	0.2	2.34	0.20	1.67							
DesferB	0.29	22.4	0.6	0.2	1.43	0.12	3.53	0.3	21.4	2.4	0.3	1.80	0.13	2.22	0.43	20.2	0.3	0.12	2.98	0.84	1.16
2 nM	0.29	22.3	0.4	0.2	1.41	0.11	3.47	0.3	21.4	NA	0.34	2.04	0.20	2.28							
Ferchr	0.61	22.1	0.2	0.1	2.1	0.11	3.40	0.42	21.5	0.6	0.2	2.66	0.16	2.41	0.55	20.3	0.2	0.2	4.6	0.99	1.48
2 nM	0.61	22.6	0.2	0.1	1.66	0.05	3.68	0.42	21.4	0.4	0.2	3.00	0.15	2.37							
Ferriox	2.39	22.7	0.3	0.2	2.79	0.08	3.35	2.2	22.5	NA	0.8	2.84	0.16	2.92	2.33	21.7	0.2	0.2	3.94	0.12	2.50
2 nM	2.39	22.8	0.53	0.23	2.77	0.08	3.50	2.2	22.9	NA	1.1	2.83	0.16	3.29							
Ferriox	4.34	23.1	0.2	0.2	5.03	0.06	4.04	4.15	22.3	NA	0.6	5.01	0.14	2.86	4.28	21.6	0.2	0.1	6.81	0.14	2.63
4 nM								4.15	22.2	0.8	0.3	5.26	0.08	2.82							
Vibrio	0.58	23	NA	0.4	1.25	0.07	3.94	0.39	21.6	NA	0.5	1.40	0.19	2.23	0.52	21.2	0.1	0.1	3.18	0.18	2.24
2 nM	0.58	23.5	NA	0.5	1.03	0.05	4.21	0.39	21.5	NA	0.5	1.19	0.16	1.99							
Model B																					
FA, 0.2	1.39	22.3	0.2	0.1	2.04	0.08	3.21	1.2	20.6	0.2	0.1	3.68	0.32	1.62	1.33	20.4	0.1	0.1	8.11	1.28	1.84
mg/L	1.39	22.6	0.8	0.3	1.69	0.11	3.13	1.2	21.2	0.2	0.2	3.95	0.15	2.26	1.33	20.6	0.1	0.1	6.75	0.65	1.92
HA, 0.1								2.2	20.7	0.3	0.2	5.43	0.47	1.80	2.33	20.6	0.1	0.1	10.45	1.01	2.08
mg/L								2.2	20.8	0.2	0.1	5.82	0.35	1.95							
HA, 0.2	3.89	21.7	0.2	0.2	3.69	0.28	1.35							2.33	21.2	0.01	0.1	*	0.68	2.80	
mg/L	3.89	21.9	NA	0.4	3.49	0.75	0.98														

* Unreliable result: Fe is added up to 12.5 nM; therefore $[L] = 13.23$ cannot be calculated in a correct way, even though the SD of the fitted value is relatively low. NA: SE down could not be determined for data that fitted the Langmuir isotherm less well.

TAC and SA5 (by a factor of 0.56–0.87, Table 3). But the DTPA ligand concentration was overestimated by SA25 (by a factor of 1.31, Table 3; see also titration in Fig. 1). Indeed, in the log-log plots (Fig. 2a) our results are well off from the theoretical titration line. Additionally, at the very low concentrations the data points of all applications deviate from the modeled curves.

4.2.2 Phytic acid

The differences between the applications are also not large for phytic acid; the two SA applications are even quite similar. The large difference for the K^{cond} values of phytic acid estimated by TAC was not expected when comparing the two very similar titration curves. We found that small changes in determined FeTAC2 concentrations at low Fe additions could be responsible for this difference (Fig. 2b). When the calculation was repeated for the combined titrations, we obtained $\log K^{\text{cond}} = 22.2 (\pm 0.2 \text{ and } 0.2)$ and $[L] = 1.6 \pm 0.1 \text{ nMEq}$.

4.2.3 Siderophores

The siderophores have high K^{cond} , so high that the AL should not be able to compete. However, although the here-estimated K^{cond} values are the highest compared to other model A ligands, curved titrations were still obtained (Fig. 1c, d, e), although there is considerable variance ($K^{\text{cond}} = 22.1\text{--}23.5, 21.4\text{--}22.9$ and $20.2\text{--}21.7$ for TAC, SA5 and SA25, respectively). The K^{cond} obtained for ferrichrome is close, almost identical, to those for desferrioxamine B for the three applications. However, $[L]$ values obtained for ferrichrome are higher than found for desferrioxamine B, with a factor of 1.4–1.6. The K^{cond} values for desferrioxamine B are lower than the value calculated from thermodynamic stability constants (Hider and Kong, 2010; Schijf and Burns, 2016) (Tables 2 and 3). Although we want to focus on comparing the applications and not on the exact values, here we need to compare with literature values. The $K^{\text{cond}} = 20.2$ for desferrioxamine B obtained for SA25 is much lower than measured by Rue and Bruland (1995), $K^{\text{cond}} \geq 23$, although they recovered 100 % of the added 2.5 nM. However, we note that they used another protocol and applied 4 min of nitrogen purging before every measurement, which would have interfered with the signal stability according to Abualhajja and

Table 3. The differences between the results in Table 3 of the three applications, ratios or difference in concentration are given. Added model ligand concentrations are given in column 1. The far right column contains the percentual deviation from the added ligand concentration as $E(\%) = (([L_{AL}] - [x])/[x]) \times 100$, with $[L_{AL}]$ as the result of the applied ligand method and $[x]$ as the added concentration of the model ligand being 2, 4, 2.9, 3.2 or 6.4 nM (as indicated in the first column at the left side under the name of the model ligand). $\log K$ is used for $\log K^{\text{cond}}$ with respect to Fe^{3+} . Data containing the ligand concentration are from the model ligands added at a concentration of 2 nM and thus exclude HA and FA. For these we used the ligand site concentrations of 2.92 and 6.4 nM Eq Fe for 0.2 mg of added fulvic and humic acids from Sukekava et al. (2018) and Laglera and van den Berg (2009). Since humic acids are not discrete ligands, the estimate % E is in italic.

Name	Logarithm values									$E_{AL}(\%) = ([L_{AL}] - [x])/[x] \times 100\%$		
	Log values			Log values with respect to Fe^{3+}			$L_{SA5}/$	$L_{TAC}/$	$L_{TAC}/$			
	$K_{SA5}/$	$K_{TAC}/$	$K_{TAC}/$	$\alpha_{SA5}/$	$\alpha_{TAC}/$	$\alpha_{TAC}/$	$L_{SA5}/$	$L_{TAC}/$	$L_{TAC}/$	E_{SA5}	E_{SA25}	E_{TAC}
Model A												
DTPA	1.02	1.01	1.03	1.01	1.01	1.02	0.44	1.18	0.52	-43	31	-33
2 nM		1.00			1.02			1.52		-43		-14
Phytic acid	1.03	1.06	1.09	1.05	1.08	1.13	0.91	0.65	0.59	42	56	-8
2 nM		1.10			1.06			0.56		17		-35
Desferrioxam. B	1.06	1.04	1.11	1.09	1.06	1.16	0.61	0.79	0.48	-10	49	-29
2 nM		1.04			1.05			0.69		2		-30
Ferrichrome	1.06	1.03	1.09	1.08	1.04	1.12	0.58	0.79	0.46	33	130	5
2 nM		1.06			1.06			0.55		50		-17
Ferrioxamine E	1.04	1.01	1.04	1.03	1.00	1.03	0.72	0.98	0.71	42	97	39
2 nM		1.00			0.98			0.98		42		38
Ferrioxamine E	1.03	1.03	1.07	1.02	1.05	1.07	0.74	1.00	0.74	25	70	25
4 nM										31		
Vibriobactin	1.02	1.06	1.08	1.00	1.10	1.10	0.44	0.89	0.39	-30	59	-38
2 nM		1.09			1.14			0.87		-40		-49
Model B												
FA	1.01	1.08	1.09	0.98	1.09	1.07	0.45	0.55	0.25	26	178	-30
0.2 mg/2.9 nM	1.03	1.06	1.10	1.03	1.03	1.06	0.59	0.43	0.25	35	131	-42
HA	1.01	1.05	1.06	0.98			0.52	0.34	0.18	70	226	-41
0.1 mg/3.2 nM	0.98	1.05	1.03	0.94			0.44	0.30	0.13	82	317	
HA												-43
0.2 mg/6.4 nM												-46

van den Berg (2014). Buck et al. (2010) also successfully recovered 100 % of a different siderophore (aerobactin) using SA25. Witter et al. (2000) measured siderophores with CLE-AdCSV but using NN, and their results compare better with the results obtained here. They found a range of K^{cond} values for a range of siderophores and measured 21.6 for both desferrioxamine B and ferrichrome. These values are very close to those obtained here by SA5 (21.4–21.5). However, their K^{cond} for phytic acid was K^{cond} higher (22.3 with respect to Fe^{3+}) than what we found with all the SA applications. However, other research reported lower K^{cond} for Fe–phytic acid complexes. Schlosser and Croot (2008) combined cross-flow ultrafiltration with the Fe radioisotope (^{55}Fe) and obtained a substantially lower value (18.6 with respect to Fe^{3+}). Moreover, phytic acid can form colloids with FeIII (Anderson, 1963). Colloid formation will interfere in several ways with the analysis by the loss of Fe, since formation of colloids results in a potentially inert fraction of Fe, the loss of phytic acid and interference of the colloids on the mercury electrode surface. However, the formation of these colloids is dependent on the phytic acid concentration (Rijkenberg et

al., 2006; Purawatt et al. 2007), and at 2 nM phytic acid we do not expect colloids to be formed. The ratios of added $[L]$ and obtained values by CLE-AdCSV by Witter et al. (2000) varied between 0.8 and 1.7, resembling our results, although the ratio was 1 for both desferrioxamine B and ferrichrome. Thus, even for model ligands there is no consistence in the literature between ligands or methods, suggesting problems in the standardization of the methodology. It is possible that the siderophores used are not of 100 % purity, which would result in a systematic underestimation of $[L]$. However, whilst it is interesting to note absolute values, our research focuses on differences between the three applications, which should not be impacted by any impurities.

The theoretical titration curve for desferrioxamine B has a relatively large offset at low concentrations compared to the modeled results (Fig. 2c). This indicates that the theoretical K^{cond} was not even approached by the three applications. That we obtained (and not for the first time) clearly curved titrations, where we should not within the applied D values, is hard to explain. One possible explanation could be a reaction taking place at the electrode surface in CSV pro-




Figure 2. Iron titrations of UV-irradiated seawater containing model ligands in competition with TAC, SA5 and SA25 as added ligand (FeAL) versus total dissolved Fe. The same data as in Fig. 1 are presented but with a log-log transformation. The lines represent back calculated titration curves with the data from Table 2, and the markers are the actual data points. The dashed black lines in (a) and (c) are back-calculated titration curves using theoretical K^{cond} calculated from the thermodynamic constants and 2 nM as model ligand concentration. (a) DTPA, (b) phytic acid, (c) desferrioxamine B, (d) ferrichrome, (e) ferrioxamine E (saturated with Fe), (f) vibriobactin, (g) fulvic acid FA and (h) humic acid HA. Data for TAC and SA5 are from duplicate experiments, and data for SA25 are from single experiments, except for FA and HA where for SA25 duplicate experiments were also done. Note the different HA concentrations, 0.1 and 0.2 mg.

moting ligand exchange of Fe(III) siderophore complexes, which produces a current. Another alternative explanation might be aluminum competition (which is not accounted for by the thermodynamic constants) since Al complexes with siderophores are detected in MS analysis of samples (Gledhill et al., 2019). The Al content, however, is unknown.

Ferrioxamine E was the only model ligand that was saturated with Fe prior to the start of the experiment. Moreover, none of the ALs should sequester Fe from ferrioxamine E, which is required in order to estimate K^{cond} because its K^{cond} is too high, outside D (Apte et al., 1988; Hudson et al., 2003; Gerringa et al., 2014). This can be explained by considering the Langmuir isotherm, used to derive K^{cond} and $[L]$:

$$[\text{FeL}] = \frac{K^{\text{cond}} [L] [\text{Fe}^{3+}]}{K^{\text{cond}} [\text{Fe}^{3+}] + 1}, \quad (5)$$

or

$$\frac{[\text{FeL}]}{[L]} = \frac{K^{\text{cond}} [\text{Fe}^{3+}]}{K^{\text{cond}} [\text{Fe}^{3+}] + 1}, \quad (6)$$

which shows that when $K^{\text{cond}} \times [\text{Fe}^{3+}] = 1$, $[\text{FeL}] = 0.5 [L]$, the equivalence point of the titration, where an almost linear relationship between $[\text{FeL}]$ and $[L]$ changes into an asymptotic relationship. In an Fe titration when $K^{\text{cond}} [\text{Fe}^{3+}] \gg 1$,

$[\text{FeL}]$ will approach $[L]$. When a titration starts at initial $[\text{FeL}] > 0.5 [L]$, the ability to estimate K^{cond} diminishes substantially. In the asymptote, at much larger values of $[\text{FeL}] / [L]$, the dependence of K^{cond} is lost and K^{cond} becomes impossible to derive. Therefore, the titration of ferrioxamine E was more or less a standard addition and in theory $[L]$ should be equivalent to the determined DFe concentration. However, all three applications overestimated the ligand concentration, but the estimated $[L]$ of ferrioxamine E compares very well for TAC and SA5 and even SA25. The relatively large difference in resulting K^{cond} between the two SA applications for the hydroxamate siderophores was not expected since the D values are close. The two most probable explanations are D and lack of equilibrium. Possibly the siderophores have α_{FeL} at the borders of and greater than D , and therefore a small decrease in D still had a consequence for the outcome of the calculations. The short waiting time may be the other reason for the deviation of SA25, which we will discuss in the next section.

The $[L]$ of the catechol vibriobactin was underestimated by SA5 and TAC and overestimated by SA25. We believe that this divergence was a combination of lack of equilibrium due to the high stability of Fe–vibriobactin complexes during the short equilibrium period of SA25 and consequent overestimation of $[L]$ and possibly by the tendency of cat-

echolates to oxidize or hydrolyze in water (Brickman and McIntosh, 1992), which could have resulted in partial loss of vibriobactin during overnight equilibration.

4.2.4 Humic substances

Our titrations of FA and HA show remarkable differences between the applications, $[L]$ by TAC was 13 %–25 % of $[L]$ by SA25 (Table 3). TAC detected 55 %–70 % of FA and HA in contrast to an early report that TAC could not detect any portion of IHSS humic reference material (Laglera et al., 2011). Our FA and HA results are in line with the partial detection of HS by TAC that was observed already by several field studies, where an increase in $[L]_{\text{TAC}}$ correlated with an increase in natural humics (Gerringa et al., 2017; Dulaquais et al., 2018; Slagter et al., 2017, 2019; Laglera et al., 2019). HS showed the largest deviations from the expected (literature) results of all tested ligands in $[L]$. Humic substances are ubiquitous in seawater (Laglera et al., 2009; Whitby et al., 2020; Yamashita et al., 2020) and potentially more representative of the dominant fraction of dissolved organic matter actually present in seawater than the model A ligands tested here, since although siderophores are detected in seawater, they are typically only present at pM concentrations (Mawji et al., 2008, 2011; Velasquez et al., 2016; Boiteau et al., 2018). The different results between the three applications could explain a major part of the offset between the TAC and SA methods in natural waters (Buck et al., 2012, 2015; Slagter et al., 2019; Ardiningsih et al., 2021). Moreover, the deviation in K^{cond} obtained by TAC from the other two applications ($K_{\text{TAC}}^{\text{cond}}/K_{\text{SA5}}^{\text{cond}}$ and $K_{\text{TAC}}^{\text{cond}}/K_{\text{SA25}}^{\text{cond}}$ up to 1.1) is greater than for most model A ligands (Table 3). This is also likely to be linked to the heterogeneity of humic substances, which means the detection window of each method will have a greater influence on the groups of binding sites titrated during the experiments. We cannot provide a definitive explanation for the $[L]$ spread. TAC showed almost straight-line patterns for FA and HA (Fig. 1g, h), as in HS-rich estuarine waters (Gerringa et al., 2007; Croot and Johansson, 2000). This could be compatible with a fraction of HS being too strong and a fraction too weak to compete with TAC (both fractions would be at or beyond the upper and lower limits of D). There is an abundant presence of strong binding sites in HS that may not be outcompeted by TAC, since desferrioxamine *B* could also not outcompete all HS binding sites in Arctic Ocean samples (Laglera et al., 2019). Another possible explanation for the similar recoveries for FA and HA, despite their reported different affinity for iron, is that TAC could form interactions with some of the binding groups of HS, canceling their interaction with iron. In other words, the use of TAC would not obey Langmuir assumption 5. For the SA applications, $[L]$ with SA25 seems to be substantially over the literature values in contrast to SA5. Titration data of HA with SA5 showed detectable levels of Fe–AL at low total Fe concentrations, while for SA25

they could not be seen. Thus, the formation of the electroactive Fe–SA complex does not happen until after 6–7 nM Fe has been added (0.1 mg HA, or over 10 nM for 0.2 mg HA). This is most probably an effect of ongoing association and dissociation processes between Fe, SA and HA, i.e., a lack of equilibration. Another explanation could be a substantial decrease in the sensitivity caused by adsorption of free and complexed humics onto the surface of the electrode, shielding the electrode from interaction with Fe(Al) complexes (Laglera et al., 2011, 2017). Adsorption of humics at the mercury electrode has been extensively discussed by Buffle and co-authors (Buffle and Cominoli, 1981; Cominoli et al., 1980), and the drop of sensitivity for CLE-AdCSV was discussed in Laglera et al. (2011 and 2017). The SA25 application does not obey Langmuir assumption 1. This might also explain the large data spread in Fig. 2h for SA25.

4.2.5 Overall

The log-log plots for DTPA and desferrioxamine *B*, between known and observed conditional stability constants, show that the data points obtained by TAC are closest to the theoretical curve of DTPA, and those obtained by SA5 are closest to desferrioxamine *B* (Fig. 2a, c). However, the TAC application has the highest D and should therefore be better equipped to detect desferrioxamine *B* and least equipped to detect DTPA. The different results between applications are mostly due to data in the first curved part of the titration as shown in Fig. 1 and illustrated when compared with the theoretical titration curves. At this part of the titration curve, peaks should in many cases be below the detection limit. Thus the precision of these measurements is very low. The log-log plots (Fig. 2) emphasize the differences between expected and observed values. The observations seem to overestimate the FeAL at low metal additions. Possible reasons are

1. electrochemical, for example a catalytic effect becoming more important at low concentrations and enhancing the signal or tiny peaks caused by impurities of the reagent or the methanol solvent as shown in previous work with NN (Boye et al., 2001),
2. concentrations are more likely to be overestimated near the detection limit,
3. desorption from conditioned cells and electrode surfaces is more significant at low concentrations.

Further analysis is required in order to resolve these possibilities and verify that the response of FeAL is linearly related to $[\text{Fe}']$ or $[\text{Fe}^{3+}]$, even at very low Fe concentrations (< 0.5 – 1 nM). Moreover, in order to obtain reliable estimates of $[L]$ and K^{cond} , we suggest that samples should have $[L']$ greater than $2 \cdot D_{\text{Fe}}$ to ensure the titration starts at low enough $[\text{Fe}']$ or $[\text{Fe}^{3+}]$ since it is this part of the curve that is used to calculate K^{cond} (i.e., where $[\text{FeL}] \leq 0.5 [L']$; Eqs. 5 and 6).

It is also possible that reactions occurred during the cathodic scan, which could also explain the deviating results of $[L]$ for vibriobactin, which was underestimated by 52 %–62 % using TAC and 60 %–70 % using SA5. Free catechols can be electro-active (Fakhari et al., 2008), and even a small contribution to the CSV peak from the ligand side of the complex would lead to a significant underestimation of the complexed fraction. A last explanation of underestimating ligand concentrations can be contamination after sampling for Fe determination by ICPMS took place.

Considering average $[L]$ and the spread in $[L]$ in the titrations with 2 nM of model A ligands (without considering the saturated ferrioxamine), average $[L]$ was 1.51 ± 0.32 , 3.30 ± 0.76 and 1.96 ± 0.73 nM Eq Fe for TAC, SA25 and SA5, respectively. There appeared to be model-ligand- and AL-dependent variations in the estimation of $[L]$ as also illustrated in Fig. S6. We can conclude that TAC underestimated most added model ligand concentrations, with a model-ligand-dependent degree of underestimation between 0.55–0.7 for HS and an average of 0.8 (0.52–1.05) for the model A ligands. The application with SA5 both over- and underestimated model ligand concentrations but less than SA25 and TAC, respectively, although HS was overestimated by a factor of 1.27–1.82 (assuming the number of binding sites from literature). The application with SA25 overestimated the concentrations by a factor of 1.3–2.3 for model A ligands and 2.33–4.17 for HS. However, it must be remembered that $[L]$ and K^{cond} are not determined independently, and unfortunately, comparison of thermodynamic constants with our results suggests that K^{cond} cannot be estimated precisely. Here SA applications result in worse estimates of K^{cond} due to a lack of data at $\text{FeL} < L$ and the possibility that the ligands are outside D , the detection window. As far as we know, four publications describing an intercomparison exist. Three of these compared SA25 and TAC (Buck et al., 2012, 2016; Slagter et al., 2019) and one compared SA5 and TAC (Ardiningsih et al., 2021). In all publications, $[L_{\text{SA}}]$ is larger than $[L_{\text{TAC}}]$ when the data are fitted with a one-ligand model. In Buck et al. (2016) it was concluded that, when using the same calculation method, comparison between the results of both applications seemed good, with one exception that SA could measure a second ligand whereas TAC could not, and therefore the total ligand concentration obtained with SA25 was always considerably larger (their Fig. 2e). This difference was attributed to an underestimation by TAC because TAC does not detect binding sites of humic substances and cannot discriminate a second ligand as well as SA25 (Buck et al., 2012, 2016; Slagter et al., 2019). Slagter et al. (2019) sampled in the Arctic Ocean where the Transpolar Drift transports high concentrations of humic substances in the upper 50 to 80 m. Since the humic content was an important feature, TAC was compared with SA25 (Slagter et al., 2019) and the voltammetric determination of humic acids (Sukekava et al., 2018). Slagter et al. (2019) found $[L_{\text{TAC}}]/[L_{\text{SA25}}] = 0.6$. However, this ratio hardly varied with the concentration of

humic substances, a strong indication that the underestimation of humics with the TAC method was not the only explanation for the difference in ligand concentration between the two methods. Ardiningsih et al. (2021) compared TAC and SA5 in the Arctic Fram Strait and also concluded that the offset between TAC and SA5 could not only be directly ascribed to underestimation of binding sites in humic substances. Moreover, Ardiningsih et al. (2021) found a relatively constant $[L_{\text{TAC}}]/[L_{\text{SA5}}]$ on the Greenland shelf but a variation in $[L_{\text{TAC}}]/[L_{\text{SA5}}]$ between 0.6 and 1 in Fram Strait. It was largely the inconsistencies in these studies, where humic substances were believed to potentially play a key role in Fe speciation, that led to this study. In future use of CLE-AdCSV, careful consideration is needed for the interpretation of the obtained $[L]$ in relation to the application and environmental variations in ligand groups, especially the humic substances.

No intercomparison between SA5 and SA25 has been undertaken since the SA5 application was published in 2014. The question remains as to why SA5 and SA25 in this work give different results. One explanation might be disequilibrium of the SA25 application. To further study the equilibration process, we executed some kinetic experiments.

4.3 Kinetic measurements

4.3.1 In-cell kinetics

In-cell kinetics were first performed with three types of samples: normal seawater, UV-irradiated seawater and UV-irradiated seawater containing DTPA at large concentrations (200 for TAC and 40 for both SA concentrations) (Fig. 3). Further, in-cell kinetic experiments were done on a subset of the model ligands at lower concentrations (2 nM or 0.2 mg). For the model A ligands, DTPA was chosen since it was used as the calibrating ligand. Desferrioxamine *B* was chosen to represent the hydroxamates, and vibriobactin was chosen to represent the catecholates. Phytic acid was also included because the titration results of all applications were in agreement. FA was used to represent model B ligands, as heterogeneous natural organic matter.

The in-cell kinetic measurements with high DTPA gave completely different results between TAC and SA (Figs. 3 and 4, Metrohm stand used). For SA25, the peak decrease was high at $t = 3$ min (initial value) and dropped sharply to values close to the limit of detection within an hour in UV-irradiated seawater and UV+DTPA. For TAC we observed a slow increase in the $\text{Fe}(\text{TAC})_2$ concentration followed by an asymptotic change to a constant value, as expected for a product of a ligand exchange reaction tending towards equilibrium.

Kinetic experiments with the low concentrations of model ligands in a volume of 10 mL were not pursued with the SA applications, only with TAC. In these experiments, equilibrium between TAC and model ligands was reached after




Figure 3. In-cell kinetic experiments, with the three AL applications, in UV-irradiated seawater, UV-irradiated seawater + DTPA (200 nM for TAC and 40 nM for the SA25 application) and natural seawater. For SA25 UV and UV + 40 nM DTPA was done. At $t = 0$ AL is added. (a) TAC. (b) SA25.




Figure 4. In-cell kinetic experiments at different $[SA]$ in UV-irradiated seawater, peak heights versus time. Measurements versus time are done in the same 10 mL, using the Metrohm electrode; drop size = 1, with regular purging with air. At $t = 0$ SA was added.

approximately 6 h, as observed previously (Croot and Johansson, 2000), 4 h for most ligands and 6 h for desferrioxamine *B* (Fig. S7). Although the rate of $Fe(TAC)_2$ formation changed with the type of model ligand, a steep increase where the relative weaker ligands were added, DTPA, FA and phytic acid, compared with a slow and steady increase where desferrioxamine *B* and vibriobactin were added, all model ligands followed the theoretical ligand exchange concentration evolution (Fig. S7).

Possible explanations of the rapid decrease in peak height are as follows.

1. The decrease can be due to formation of the non-electro-labile $Fe(SA)_2$ complex. $Fe(SA)_2$ is the non-electro-active species (Abualhaija and van den Berg, 2014) and becomes the dominant $Fe(SA)_x$ complex at higher SA concentrations. The two forms of $Fe-SA$ would have

different formation kinetics, with a slower formation of $Fe(SA)_2$, with Fe coming not from the dissociation of the model complex but from the dissociation of $Fe(SA)$. This process increases the time to reach equilibrium, and D changes accordingly. We monitored the electro-labile $Fe(SA)$ concentrations after SA additions in the range 2.5 and 50 μM using the adapted Metrohm instrument with a small mercury drop (size 1) and regular air purging (Fig. 4). Possible contributions due to decreasing oxygen were excluded and due to adsorption on mercury on the cell bottom were minimized. At 25 μM SA the concentration of the electro-active species practically disappeared after 2 h. At $SA < 25 \mu M$, the concentration of the electro-labile species $Fe(SA)$ decreased exponentially with time for a period of at least 13 h. At concentrations $\leq 5 \mu M$ there is a decrease to a constant value above zero. These results support the formation of a non-electro-active species $Fe(SA)_2$ irrespective of adsorption on the mercury drop, confirming the results of Abualhaija and van den Berg (2014).

2. Formation of $Fe(SA)_2$ from $FeSA$ is slow and probably also irreversible. We investigated this possibility by trying to force dissociation of $Fe(SA)_2$ by adding the competing model ligand DTPA during the decrease in the CSV signal in a kinetic experiment. Addition of DTPA did show a sudden decrease in signal with SA5, but not with higher SA concentrations (the experiment was done at 5 and 15 μM). We suggest that at the low SA (5 μM) DTPA competed with $FeSA$, causing a decrease in $FeSA$ and thus in peak height. Adding DTPA at the higher SA concentration of 15 μM, where $Fe(SA)_2$ is dominant, only a slight decrease in peak height was possible because only $FeSA$ could dissociate and not $Fe(SA)_2$ within the 2 h of the experiment (Fig. 5). This

result indicates irreversible formation of $\text{Fe}(\text{SA})_2$ and has important implications for overnight equilibration.

- Adsorption on the mercury at the bottom of the cell as indicated by Buck et al. (2007) and contradicted by Abualhaija and van den Berg (2014). However, both used different analytical equipment, with the latter testing with a Metrohm stand, characterized by smaller mercury drops and automatic air purge. We checked the effect of the drop size for all three applications, including TAC. The TAC application did not show any decrease in signal with time and with increasing mercury at the cell bottom. On the contrary, the two SA applications did show a decrease that was steeper and larger with increasing drop size. The decrease in the SA5 application was larger than in the SA25 application (Fig. S3). A positive linear relationship ($r^2 = 0.98$) exists for SA25 between the decrease in peak height within 43 min and the volume of dispensed mercury at the bottom of the cell. However, no relation exists for SA5, although the reduction is strongest in that application (Fig. S3). We tested whether SA was reversibly adsorbed to the mercury puddle at the bottom of the cell by transferring mercury accumulated under SA5 and SA25 protocols into a cell filled with 10 mL UV-irradiated seawater with 6 nM of added Fe. No SA was released from the mercury into the seawater as no peak could be detected when analyzed with the normal AdCSV procedure. An explanation could be that only $\text{Fe}(\text{SA})_2$ adsorbs on the mercury, causing a direct relationship with peak reduction and mercury volume at high SA. Since $\text{Fe}(\text{SA})_2$ might be formed irreversibly, no release of SA into the solution that could lead to the formation of the electro-active FeSA species would be possible. It remains hard to explain the strong reduction of the peak height without a relationship with the mercury volume at SA5.
- The lack of purging influences the conditions. The BASi electrode only allows either stirring or purging in an automatic measurement, and stirring is the normal practice. Purging with air should maintain a constant concentration of oxygen and should increase the sensitivity and prevent decreasing peak heights with time (Abualhaija and van den Berg, 2014). We checked the effect of an air purge step on the SA measurements of both electrodes, BASi and Metrohm. The decrease in peak height with time was not influenced by an air purge step in both electrodes (Fig. S2).

We can conclude that the formation kinetics of $\text{Fe}(\text{TAC})_2$ using in-cell experiments with model ligands reached equilibrium within 8 h. In-cell kinetic experiments with SA did not reach equilibrium and showed a continuous decay of the peak height. This can be explained by a combination of processes like adsorption of $\text{Fe}(\text{SA})$ complexes on dispensed mercury




Figure 5. Reversibility of Fe–SA formation upon addition of 150 nM DTPA at $t = 80$ min. The experiment was undertaken with a Metrohm electrode in-cell in UV-irradiated seawater with 6 nM Fe and two SA concentrations, 5 μM SA (right-hand side y axis) and 15 μM SA (left-hand side y axis).

at the cell bottom and formation of the irreversible, according to Abualhaija and van den Berg (2014), non-electro-active $\text{Fe}(\text{SA})_2$ ($\text{FeSA} + \text{SA}$ forming FeSA_2). The formation of irreversible species is not compatible with techniques such as CLE-AdCSV that require a dynamic equilibrium between competing ligands before analysis.

4.3.2 Bottle kinetics

The kinetic experiments were repeated extracting 10 mL aliquots from a 200 mL bottle. Experiments carried out were UV-irradiated seawater, UV-irradiated seawater with desferrioxamine *B*, UV-irradiated seawater with phytic acid and UV-irradiated seawater with FA (Fig. 6). Some points have to be considered for interpretation. The conditioning procedure of the bottle in which the reaction takes place raises the question of whether to condition with or without the AL. Addition of AL should occur at $t = 0$, and conditioning of the bottle with AL is thus not possible. Conditioning without AL with UV-irradiated seawater with 6 nM of added DFe could cause Fe precipitation and/or adsorption on the bottle walls. DFe at the end of the experiment is probably higher than 6 nM due to Fe desorption from the bottle wall.

TAC

The results with TAC showed the same pattern as for the in-cell experiments: an increase that levels off to equilibrium for UV-irradiated seawater, phytic acid and FA (Fig. 6) characteristic of a ligand exchange reaction reaching a steady state. Higher equilibrium signals prove an extra Fe input from the bottle wall. The experiment with desferrioxamine *B* did not level off after 8 h, and the slope was less steep. This concurs with our in-cell observation that desferrioxamine *B* dissociates more slowly. That equilibrium is not reached even after 8 h explains, at least for this model ligand, the underestimation of $[L]$ by TAC (Table 2; Croot and Johansson, 2000). It




Figure 6. Kinetic measurements for the formation of Fe–SA complexes the change in peak height versus time. At $t = 0$ the AL is added. In-cell means the whole experiment is done in the same 10 mL which was placed in the cell; dispensed mercury accumulates at the bottom of the cell. Bottle experiment means for every measurement a fresh 10 mL was taken from a large volume of sample. Here the reaction takes place in the large volume, and AL was added at $t = 0$. (a, b) UV-irradiated seawater (UV), (c, d) UV + desferrioxamine B, (e, f) UV + phytic acid bottle (a, c, f) and in-cell experiments (b, d, f) are shown. (g) For FA only bottle experiments were done with SA5 and SA25 as AL.

also indicates that with this protocol kinetics were dependent not just on ligand exchange but also on Fe desorption and/or redissolution of precipitated Fe.

SA

The results of the bottle experiments for SA5 changed dramatically with respect to those of the in-cell protocol. They show that equilibrium appears to be achieved for phytic acid and fulvic acid, and the increase in signal over time in the first 1–2 h is similar to that for TAC (Fig. 6). An equilibrium

is reached after approximately 4 h with phytic acid and FA. More time, 8 h, is needed to reach an equilibrium with desferrioxamine *B*, in line with the slower formation of FeTAC_2 in the presence of desferrioxamine *B* compared to the other model ligands (Figs. 6, S6).

Equilibrium or a steady state is difficult to establish for SA25. A plateau is reached after approximately 1 h. This seemingly depends on the added ligand, but after the plateau the signal decreases at a steady rate. We can conclude thus far that the steep decrease shown in the BASi electrode in the in-cell kinetics does not happen with the bottle experiments. This result points to adsorption on the dispensed mercury puddle as the main cause of the disappearing signal. Equilibrium is not reached at short waiting times after addition of 25 μM SA. Therefore, the use of the Langmuir isotherm to calculate the ligand characteristics is not possible. We hypothesize that the distinction in more than one ligand group, which was often possible with this application, could have been caused by the absence of equilibrium.

5 Conclusions

All applications have drawbacks; however the SA25 application clearly does not obey the main assumption of the Langmuir isotherm: no equilibrium is reached and therefore the results cannot be reliable. In addition, the most upsetting conclusion is that the estimation of the conditional stability constant K^{cond} is a very rough estimate only and systematically biased by the AL. Comparing $[L]$ obtained by the three applications with the added concentrations of the model ligands, $[L]$ is underestimated by TAC with a factor 0.5 to 1.05 for model A ligands and 0.52–0.7 for HS model B ligands. The SA5 application both under- and overestimated ligand concentrations (0.57–1.5 for model A ligands and 1.27–1.8 for model B ligands), but our kinetic studies suggest that true equilibrium may still be an issue for strong ligands. The SA25 application overestimated $[L]$ by a factor of 1.31–2.3 for model A ligands and 2.33–4.17 for model B ligands. Moreover, for all approaches, ligand-specific interferences occurred, as for humic acids.

We confirm the conclusion of Abualhajja and van den Berg (2014) that the SA concentration needs to be low, $\leq 5 \mu\text{M}$, to prevent formation of not electro-active $\text{Fe}(\text{SA})_2$. Our experiments suggest that the formation of $\text{Fe}(\text{SA})_2$ is irreversible and thus does not obey the Langmuir equation. Possibly $\text{Fe}(\text{SA})_2$ also adsorbs on dispensed mercury on the bottom of the cell. When using a BASi electrode, small mercury drops are to be preferred.

All methods have drawbacks, and the characterization of Fe-binding organic ligands at nM concentrations in seawater with a high ionic strength and at 40–80 $\mu\text{Mol/kg}$ DOC (Hansell et al., 2009) is and remains challenging. In case voltammetric methods are used, we recommend the SA5 application, but it should not be overinterpreted, and issues

relating to the technique should be fully acknowledged. In particular, it is hard to justify determination of equilibrium constants using an adsorption isotherm that assumes equilibrium if a signal is not stable and the experimental system is therefore not at equilibrium (for whatever reason), especially bearing in mind the method specifically uses the term “equilibrium” in its title. Furthermore, it appears that apart from the constraint forced by *D*, these estimates have a much larger error than apparent from those obtained from the fits to the Langmuir isotherm, which may explain why direct links between speciation predicted from conditional parameters do not relate strongly to bioavailability (Shaked et al., 2020, 2021), although this could also be caused by the limited role assigned to iron reduction prior to bio-assimilation. Given that the research questions that are typically addressed when determining conditional parameters from the Langmuir isotherm actually relate to how variations in the parameters impact Fe speciation (e.g., abundance of Fe not bound to organic matter and thus assumed to be bioavailable), it would seem appropriate to broaden the methodology applied to these questions to other methods capable of, e.g., estimating concentrations and distributions of the dominant groups of binding sites and/or the lability and solubility of Fe. For example, Whitby et al. (2020) and Laglera et al. (2019) have used alternative methods based on voltammetry to suggest that humic substances are more important than thought before. Other recent studies have focused on the influence of the pH on metal organic complexation and vary the pH of titrations (Ye et al., 2020; Gledhill et al., 2015; Zhu et al., 2021). Here we note that the acid–base chemistry of organic matter – which underpins metal binding – is severely understudied but can nevertheless quantify the distribution and concentration of the total cation binding sites present, at least in the portion of dissolved organic matter that can be isolated from seawater (Lodeiro et al., 2020). With further knowledge of acid–base chemistry of dissolved organic matter and how it changes with inputs of fresh material from biological activity, the total binding site concentration could be independently constrained and only binding affinities derived. Alternatively models can be applied that allow for the estimation of metal speciation for ambient conditions (e.g., Hiemstra and van Riemsdijk, 2006; Stockdale et al., 2015), with predictions of how binding to organic matter could influence e.g., iron solubility (Zhu et al., 2021). The wider application of methods employing cation chelating resins, such as diffusive thin film gradients (Zhang and Davison, 2015; Town et al., 2009; Bayens et al., 2018), might also offer alternative insights into the lability of metals in seawater. Another way to apply voltammetric methods is to characterize metal-binding ligands by pseudovoltammetry (Luther et al. (2021), although unfortunately this method is not currently suitable for DFe. Whilst continued application of the CLE-AdCSV approach will no doubt further develop our knowledge on how operationally defined ligand concentrations and stability constants vary in the ocean under a restrictive set of conditions appar-

ently specific to one added ligand, new approaches to both the determination and the interpretation of metal binding to organic matter will surely stimulate discussions in the field of the organic metal complexation and furthermore be likely lead to new insights. It is clear that further work needs to be done to effectively contextualize the database, and robust quality control procedures are urgently required. We recommend that these procedures include the determination of a standard ligand with independently determined thermodynamic constants that is within the detection window of the applied methodology. Our work suggests that none of the ligands examined here are ideal for this, since they might have been outside the detection window or do not have available thermodynamic constants for comparison.

Data availability. Data used in figures are available at <https://doi.org/10.25850/nioz/7b.b.qb> (last access: 10 August 2021, Gerringa et al., 2021).

Supplement. The supplement related to this article is available online at: <https://doi.org/10.5194/bg-18-5265-2021-supplement>.

Author contributions. The organization of the experiments was done by LG, LL and MG. The experiments were executed by or under supervision of NM, IA and LG. Evaluation of the data was done by LG, MG and LL. LG wrote the manuscript with substantial contributions of MG and LL.

Competing interests. The contact author has declared that neither they nor their co-authors have any competing interests.

Disclaimer. Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Acknowledgements. We are grateful for the help of Kristin Buck during the time we struggled to get the SA method working. The bachelor students David Amptmeijer and Robert Sluijter did preliminary experiments whereas Ismael Salazar and Martijn Korpelaar executed experiments used in this paper. The critical reading of Rebecca Zitoun helped to improve the manuscript. The critical comments of Dario Omanović and the anonymous reviewer are gratefully acknowledged. The PhD work of Hans Slagter and co-author Indah Ardiningsih made this research possible.

Financial support. This research has been supported by the Ministerio de Ciencia e Innovación (grant no. CTM2017-84763-C3-3-R) and the Lembaga Pengelola Dana Pendidikan (grant in the name of Indah Ardiningsih).

Review statement. This paper was edited by Carolin Löscher and reviewed by Dario Omanović and one anonymous referee.

References

- Abualhaja, M. M. and van den Berg, C. M. G.: Chemical speciation of iron in seawater using catalytic cathodic stripping voltammetry with ligand competition against salicylaldehyde, *Mar. Chem.*, 164, 60–74, 2014.
- Abualhaja, M. M., Whitby, H., van den Berg, C. M. G.: Competition between copper and iron for humic ligands in estuarine waters, *Mar. Chem.*, 172, 46–56, 2015.
- Achterberg, E. P., Steigenberger, S., Marsay, C. M., LeMoigne, F. A. C., Painter, S. C., Baker, A. R., Connelly, D. P., Moore, C. M., Tagliabue, A., and Tanhua, T.: Iron Biogeochemistry in the High Latitude North Atlantic Ocean, *Sci. Rep.*, 8, 1283, <https://doi.org/10.1038/s41598-018-19472-1>, 2018.
- Anderson, G.: Effect of iron/phosphorus ratio and acid concentration on precipitation of ferric inositol hexaphosphate, *J. Sci. Food Agric.*, 14, 352–359, 1963.
- Apte, S. C., Gardner, M. J., and Ravenscroft, J. E.: An evaluation of voltammetric titration procedures for the determination of trace metal complexation in natural waters by use of computer simulation, *Anal. Chim. Acta*, 212, 1–21, 1988.
- Ardiningsih, I., Gledhill, M., Lodeiro, P., Krisch, S., Reichart, G.-J., Achterberg, E. P., Middag, R., and Gerringa, L. J. A.: Natural Fe-binding organic ligands in Fram Strait and over the Northeast Greenland shelf, *Mar. Chem.*, 224, 103815, <https://doi.org/10.1016/j.marchem.2020.103815>, 2020.
- Ardiningsih, I., Zu, K., Lodeiro, P., Gledhill, M., Reichart, G.-J., Achterberg, E. P., Middag, R., and Gerringa, L. J. A.: Iron speciation in Fram Strait and over the northeast Greenland shelf: An inter-comparison study of voltammetric methods, *Front. Mar. Sci.*, 7, 609379, <https://doi.org/10.3389/fmars.2020.609379>, 2021.
- Avendaño, L., Gledhill, M., Achterberg, E. P., Rérolle, V. M. C., and Schlosser, C.: Influence of ocean acidification on the organic complexation of iron and copper in Northwest European shelf seas; a combined observational and model study, *Front. Mar. Sci.*, 3, 58, <https://doi.org/10.3389/fmars.2016.00058>, 2016.
- Bayens, W., Gao, Y., Davison, W., Galceran, J., Leermakers, M., Puy, J., Superville, P.-J., and Beguery, L.: In situ measurements of micronutrient dynamics in open seawater show that complex dissociation rates may limit diatom growth, *Sci. Rep.*, 8, 16125, <https://doi.org/10.1038/s41598-018-34465-w>, 2018.
- Barbeau, K., Rue, E. L., Bruland, K. W., and Butler, A.: Photochemical cycling of iron in the surface ocean mediated by microbial iron(III)-binding ligands, *Nature*, 413, 409–413, 2001.
- Barbeau, K.: Photochemistry of organic iron(III) complexing ligands in oceanic systems, *Photochem. Photobiol.*, 82, 1505–1516, 2006.
- Batchelli, S., Muller, F. L. L., Chang, K.-C., and Lee, C.-L.: Evidence for strong but dynamic iron-humic colloidal associations in humic rich coastal waters, *Environ. Sci. Technol.*, 44, 8485–8490, 2010.
- Billler, D. V. and Bruland, K. W.: Analysis of Mn, Fe, Co, Ni, Cu, Zn, Cd, and Pb in seawater using the Nobias-chelate PA1 resin and magnetic sector inductively coupled

- plasma mass spectrometry (ICP-MS), *Mar. Chem.*, 130, 12–20, <https://doi.org/10.1016/j.marchem.2011.12.001>, 2012.
- Boiteau, R. M. and Repeta, D. J.: An extended siderophore suite from *Synechococcus* sp PCC 7002 revealed by LC-ICPMS-ESIMS, *Metallomics*, 7, 877 <https://doi.org/10.1039/c5mt00005j>, 2015.
- Boiteau, R. M., Till, C. P., Coale, T. H., Fitzsimmons, J. N., and Bruland, K. W.: Patterns of iron and siderophore distributions across the California Current System, *Limnol. Oceanogr.*, 9999, 1–14, 2018.
- Boyd, P. W., Watson, A. J., Law, C. S., Abraham, E. R., Trull, T., Murdoch, R., Bakker, D. C. E., Bowie, A. R., Buesseler, K. O., Chang, H., Charette, M., Croot, P., Downing, K., Frew, R., Gall, M., Hadfield, M., Hall, J., Harvey, M., Jameson, G., LaRoche, J., Liddicoat, M., Ling, R., Maldonado, M. T., McKay, R. M., Nodder, S., Pickmere, S., Pridmore, R., Rintoul, S., Safi, K., Sutton, P., Strzepek, R., Tanneberger, K., Turner, S., Waite, A., and Zeldis, J.: A mesoscale phytoplankton bloom in the polar Southern Ocean stimulated by iron fertilization, *Nature*, 407, 695–701, 2000.
- Boyd, P. W. and Ellwood, M. J.: The biogeochemical cycle of iron in the ocean *Nat. Geosci.*, 3, 675–682, 2010.
- Boyd, P. W. and Tagliabue, A.: Using the L* concept to explore controls on the relationship between paired ligand and dissolved iron concentrations in the ocean, *Mar. Chem.*, 173, 52–66, <https://doi.org/10.1016/j.marchem.2014.12.003>, 2015.
- Boye, M., van den Berg, C. M. G., de Jong, J. T. M., Leach, H., Croot, P., de Baar, H. J. W.: Organic complexation of iron in the Southern Ocean, *Deep-Sea Res. Pt. I*, 48, 1477–1497, 2001.
- Boye, M., Nishioka, J., Croot, P. L., Laan, P., Timmermans, K. R., and de Baar, H. J. W.: Major deviations of iron complexation during 22 days of a mesoscale iron enrichment in the open Southern Ocean, *Mar. Chem.*, 96, 257–271, 2005.
- Brickman, T. J. and McIntosh, M. A.: Overexpression and Purification of Ferric Enterobactin Esterase from *Escherichia coli*. Demonstration of enzymatic hydrolysis of enterobactin and its iron complex, *J. Biol. Chem.*, 267, 17, 12350–12355, 1992.
- Buck, K. N., Lohan, M. C., Berger, C. J., and Bruland, K. W.: Dissolved iron speciation in two distinct river plumes and an estuary: Implications for riverine iron supply, *Limnol. Oceanogr.*, 52, 843–855, 2007.
- Buck, K. N., Selph, K. E., and Barbeau, K. A.: Iron-binding ligand production and copper speciation in an incubation experiment of Antarctic Peninsula shelf waters from the Bransfield Strait, Southern Ocean, *Mar. Chem.*, 122, 148–159, <https://doi.org/10.1016/j.marchem.2010.06.002>, 2010.
- Buck, K. N., Moffett, J., Barbeau, K. A., Bundy, R. M., Kondo, Y., and Wu, J.: The organic complexation of iron and copper: an intercomparison of competitive ligand exchange-adsorptive cathodic stripping voltammetry (CLE-ACSV) techniques, *Limnol. Oceanogr.-Meth.*, 10, 496–515, 2012.
- Buck, K. N., Sohst, B., and Sedwick, P. N.: The organic complexation of dissolved iron along the U.S. GEOTRACES (GA03) North Atlantic Section, *Deep Sea Res. Pt. II*, 116, 152–165, <https://doi.org/10.1016/j.dsr2.2014.11.016>, 2015.
- Buck, K. N., Gerringa, L. J. A., and Rijkenberg, M. J. A.: An Intercomparison of Dissolved Iron Speciation at the Bermuda Atlantic Time-series Study (BATS) Site: Results from GEO-TRACES Crossover Station A, *Front. Mar. Sci.*, 3, 262, <https://doi.org/10.3389/fmars.2016.00262>, 2016.
- Buck, K. N., Sedwick, P. N., Sohst, B., and Carlson, C. A.: Organic complexation of iron in the eastern South Pacific: Results from US GEOTRACES Eastern Zonal Transect (GEOTRACES cruise GP16), *Mar. Chem.*, 201, 229–241, doi:10.1016/j.marchem.2017.11007, 2018.
- Buffle, J.: The analytical challenge posed by fulvic and humic compounds, *Anal. Chim. Acta*, 232, 1–2, 1990.
- Buffle, J. and Cominoli, A.: Voltammetric study of humic and fulvic substances: Part IV. Behaviour of fulvic substances at the mercury-water interface, *J. Electroanal. Chem.*, 121, 273–299, 1981.
- Bundy, R. M., Abdulla, H. A. N., Hatcher, P. G., Biller, D. V., Buck, K. N., and Barbeau, K. A.: Iron-binding ligands and humic substances in the San Francisco Bay estuary and estuarine-influenced shelf regions of coastal California, *Mar. Chem.*, 173, 183–194, 2015.
- Bundy, R. M., Biller, D. V., Buck, K. N., Bruland, K. W., and Barbeau, K. A.: Distinct pools of dissolved iron-binding ligands in the surface and benthic boundary layer of the California Current, *Limnol. Oceanogr.*, 59, 769–787, 2014.
- Cabanes, D. J. E., Norman, L., Bowie, A. R., and Strmečki, and Hassler, C. S.: Electrochemical evaluation of iron-binding ligands along the Australian GEOTRACES southwestern Pacific section (GP13), *Mar. Chem.*, 219, 103736, <https://doi.org/10.1016/j.marchem.2019.103736>, 2020.
- Caprara, S., Buck, K. N., Gerringa, L. J. A., Rijkenberg, M. J. A., and Monticelli, D.: A compilation of iron speciation data for open oceanic waters, a data report, *Front. Mar. Sci.*, 3, 221, <https://doi.org/10.3389/fmars.2016.00221>, 2016.
- Cominoli, A., Buffle, J., and Haerdi, W.: Voltammetric study of humic and fulvic substances, Part III, Comparison of the capabilities of the various polarographic techniques for the analysis of humic and fulvic substances, *J. Electroanal. Chem.*, 110, 259–275, doi:10.1016/S0022-0728(80)80378-0, 1980.
- Croot, P. L. and Johansson, M.: Determination of iron speciation by cathodic stripping voltammetry in seawater using the competing ligand 2-(2-Thiazolylazo)-p-cresol (TAC), *Electroanalysis*, 12, 565–576, 2000.
- Croot, P. L. and Heller, M. I.: The importance of kinetics and redox in the biogeochemical cycling of iron in the surface ocean, *Front. Microbiol.*, 3, 219, <https://doi.org/10.3389/fmicb.2012.00219>, 2012.
- Croot, P. L., Andersson, K., Öztürk, M., and Turner, D. R.: The distribution and speciation of iron along 6° E in the Southern Ocean, *Deep-Sea Res. Pt. II*, 51, 2857–2879, <https://doi.org/10.1016/j.dsr2.2003.10.012>, 2004.
- De Baar, H. J. W.: On iron limitation of the Southern Ocean: experimental observations in the Weddell and Scotia Seas, *Mar. Ecol. Prog. Ser.*, 65, 105–122, 1990.
- De Baar, H. J. W. and La Roche, J.: Trace metals in the oceans: evolution, biology and global change, edited by: Wefer, G., Lamy, F., Mantoura, F., *Mar. Sci. Front. Eur.*, Springer Verlag Berlin, 79–105, 2003.
- Dulaquais, G., Waeles, M., Gerringa, L. J. A., Middag, R., Rijkenberg, M. J. A., and Riso, R.: The biogeochemistry of electroactive humic substances and its connection to iron chemistry in the North East Atlantic and the Western Mediterranean Sea, *J. Geo-*

- phys. Res.-Oceans, 123, <https://doi.org/10.1029/2018JC014211>, 2018.
- Fakhari, A. R., Hossainy Davarani, S. S., Ahmar, H., and Makarem, S.: Electrochemical study of catechols in the presence of 2-thiazoline-2-thiol: application to electrochemical synthesis of new 4,5-dihydro-1,3-thiazol-2-ylsulfanyl-1,2-benzenediol Derivatives, *J. Appl. Electrochem.*, 38, 1743–1747, doi:10.1007/s10800-008-9625-0, 2008.
- Gerringa, L. J. A., Rijkenberg, M. J. A., Wolterbeek, H. T., Verburg, T., Boye, M., and de Baar, H. J. W.: Kinetic study reveals weak Fe-binding ligand, which affects the solubility of Fe in the Scheldt estuary, *Mar. Chem.*, 103, 30–45, <https://doi.org/10.1016/j.marchem.2006.06.002>, 2007.
- Gerringa, L. J. A., Rijkenberg, M. J. A., Thuróczy, C.-E., and Maas, L. R. M.: A critical look at the calculation of the binding characteristics and concentration of iron complexing ligands in seawater with suggested improvements, *Environ. Chem.*, 11, 114–136, <https://doi.org/10.1071/EN13072>, 2014.
- Gerringa, L. J. A., Rijkenberg, M. J. A., Schoemann, V., Laan, P., and de Baar, H. J. W.: Organic complexation of iron in the West Atlantic Ocean, *Mar. Chem.*, 177, 434–446, doi:10.1016/j.marchem.2015.04.007, 2015.
- Gerringa, L. J. A., Slagter, H. A., Bown, J., van Haren, H., Laan, P., de Baar, H. J. W., and Rijkenberg, M. J. A.: Dissolved Fe and Fe-binding dissolved organic ligands in the Mediterranean Sea, *Mar. Chem.*, 194, 100–113, <https://doi.org/10.1016/j.marchem.2017.05.012>, 2017.
- Gerringa, L. J. A., Gledhill, M., Laglera, L. M., Ardinarsih, I., and Muntjewerf, N.: Comparing CLE-AdCSV applications using SA and TAC to determine the Fe binding characteristics of model ligands in seawater, NIOZ, V1 [data set], <https://doi.org/10.25850/nioz/7b.b.qb>, 2021.
- Gledhill, M. and van den Berg, C. M. G.: Determination of complexation of iron (III) with natural organic complexing ligands in seawater using cathodic stripping voltammetry, *Mar. Chem.*, 47, 41–54, 1994.
- Gledhill, M. and Buck, K. N.: 2012. The organic complexation of iron in the marine environment: a review, *Front. Microbiol.*, 3, 69, <https://doi.org/10.3389/FMICB.2012.00069>, 2012.
- Gledhill, M. and Gerringa, L. J. A.: The Effect of Metal Concentration on the Parameters Derived from Complexometric Titrations of Trace Elements in Seawater – A Model Study, *Front. Mar. Sci.*, 4, 254, <https://doi.org/10.3389/fmars.2017.00254>, 2017.
- Gledhill, M., McCormack, P., Ussher, S., Achterberg, E. P., Mantoura, R. C., and Worsfold, P. J.: Production of siderophore type chelates by mixed bacterioplankton populations in nutrient enriched seawater incubations, *Mar. Chem.*, 88, 75–83, <https://doi.org/10.1016/j.marchem.2004.03.003>, 2004.
- Gledhill, M., Achterberg, E. P., Li, K., Mohamed, K. N., and Rijkenberg, M. J. A.: Influence of ocean acidification on the complexation of iron and copper by organic ligands in estuarine waters, *Mar. Chem.*, 177, 421–433, <https://doi.org/10.1016/j.marchem.2015.03.016>, 2015.
- Gledhill, M., Basu, S., and Shaked, Y.: Metallophores associated with *Trichodesmium erythraeum* colonies from the Gulf of Aqaba, *Metallomics*, 11, 1547–1557, <https://doi.org/10.1039/C9MT00121B>, 2019.
- Gustafsson, J. P.: The chemical speciation program visual MINTEQ version 3.0, 73 pp., 2012.
- Hansell, D. A., Carlson, C. A., Repeta, D. J., and Schlitzer, R.: Dissolved organic matter in the ocean. A controversy stimulates new insights, *Oceanography*, 22, 202–211, 2009.
- Hassler, C. S., Schoemann, V., Nichols, C. A. M., Butler, E. C. V., and Boyd, P. W.: Saccharides enhance iron bioavailability to Southern Ocean phytoplankton, *P. Natl. Acad. Sci. USA*, 1076–1081, doi:10.1073/pnas.1010963108, 2011.
- Hassler, C. S., Norman, L., Mancuso Nichols, C. A., Clementson, L. A., Robinson, C., Schoemann, V., Watson, R. J., and Daoblin, M. A.: Iron associated with exopolymeric substances is highly bioavailable to oceanic phytoplankton, *Mar. Chem.*, 173, 136–147, 2015.
- Hider, R. C. and Kong, X.: Chemistry and biology of siderophores, *Nat. Prod. Rep.*, 27, 637–657, 2010.
- Hiemstra, T. and van Riemsdijk, W. H.: Biogeochemical speciation of Fe in ocean water, *Mar. Chem.*, 102, 181–197, <https://doi.org/10.1016/j.marchem.2006.03.008>, 2006.
- Hudson, R. J. M.: Which aqueous species control the rates of trace metal uptake by aquatic biota?, Observations and prediction of non-equilibrium effects, *Sci. Total Environ.*, 219, 95–115, 1998.
- Hudson, R. J. M., Rue, E. L., and Bruland, K. W.: Modeling complexometric titrations of natural water samples, *Environ. Sci. Technol.*, 37, 1553–1562, <https://doi.org/10.1021/es025751a>, 2003.
- Hutchins, D. A., Witter, A. E., Butler, A., and Luther III, G. W.: Competition among marine phytoplankton for different chelated iron species, *Nature*, 400, 858–861, 1999.
- Kleint, C., Hawkes, J. A., Sander, S. G., and Koschinsky, A.: Voltammetric Investigation of Hydrothermal Iron Speciation, *Front. Mar. Sci.*, 3, 75, <https://doi.org/10.3389/fmars.2016.00075>, 2016.
- Kondo, Y., Takeda, S., and Furuya, K.: Distinct trends in dissolved Fe speciation between shallow and deep waters in the Pacific Ocean, *Mar. Chem.*, 134, 18–28, 2012.
- Krachler, R., Krachler, R. F., Wallner, G., Hann, S., Laux, M., Cervantes Recalde, M. F., Jirsa, F., Neubauer, E., von der Kammer, F., Hofmann, T., and Keppler, B. K.: River-derived humic substances as iron chelators in seawater, *Mar. Chem.*, 174, 85–93, 2015.
- Lagerström, M. E., Field, M. P., Séguret, M., Fischer, L., Hann, S., and Sherrell, R. M. X.: Automated on-line flow-injection ICP-MS determination of trace metals (Mn, Fe, Co, Ni, Cu and Zn) in open ocean seawater: Application to the GEOTRACES program, *Mar. Chem.*, 155, 71–80, 2015.
- Laglera, L. M., Battaglia, G., and van den Berg, C. M. G.: Determination of humic substances in natural waters by cathodic stripping voltammetry of their complexes with iron, *Anal. Chim. Act.*, 599, 58–66, 2007.
- Laglera, L. M. and van den Berg, C. M. G.: Evidence for geochemical control of iron by humic substances in seawater, *Limnol. Oceanogr.*, 54, 610–619, 2009.
- Laglera, L. M., Battaglia, G., and van den Berg, C. M. G.: Effect of humic substances on the iron speciation in natural waters by CLE/CSV, *Mar. Chem.*, 127, 134–143, 2011.
- Laglera, L. M., Downes, J., and Santos-Echeandí, A.: Comparison and combined use of linear and non-linear fitting for the estimation of complexing parameters from metal titrations of estuarine samples by CLE/AdCSV, *Mar. Chem.*, 155, 102–112, 2013.

- Laglera, L. M. and Filella, M.: The relevance of ligand exchange kinetics in the measurement of iron speciation by CLE-AdCSV in seawater, *Mar. Chem.*, 173, 100–113, 2015.
- Laglera, L. M., Tovar-Sánchez, A., Iversen, M. H., González, H. E., Naik, H., Mangesh, G., Assmy, P., Klaas, C., Mazzocchi, M. G., Montresor, M., Naqvi, S. W. A., Smetacek, V., and Wolf-Gladrow, D. A.: Iron partitioning during LOHAFEX: Copepod grazing as a major driver for iron recycling in the Southern Ocean, *Mar. Chem.*, 196, 148–161, 2017.
- Laglera, L. M., Sukekava, C., Slagter, H. A., Downes, J., Aparicio-Gonzalez, A., and Gerringa, L. J. A.: First direct evidence of humic substances controlling iron transport across the surface of the Arctic Ocean, *Environ. Sci. Technol.*, 53, 13136–13145, <https://doi.org/10.1021/acs.est.9b04240>, 2019.
- Lauderdale, J. M., Braakman, R., Forget, G., Dutkiewicz, S., and Follows, M. J.: Microbial feedbacks optimize ocean iron availability, *P. Natl. Acad. Sci. USA*, 117, 4842–4849, doi:10.1073/pnas.1917277117, 2020.
- Liu, X. and Millero, F. J.: The solubility of iron in seawater, *Mar. Chem.*, 77, 43–54, 2002.
- Lodeiro, P., Rey-Castro, C., David, C., Achterberg, E. P., Puy, J., and Gledhill, M.: Acid-base properties of dissolved organic matter extracted from the marine environment, *Sci. Total Environ.*, 729, 138437, <https://doi.org/10.1016/j.scitotenv.2020.138437>, 2020.
- Luther, G. W., Mullaugh, K. M., Hauser, E. J., Rader, K. J., and Di Toro, D. M.: Determination of ambient dissolved metal ligand complexation parameters via kinetics and pseudo-voltammetry experiments, *Mar. Chem.*, 234, 103998, <https://doi.org/10.1016/j.marchem.2021.103998>, 2021.
- Mawji, E., Gledhill, M., Milton, J. A., Tarran, G. A., Ussher, S., and Thompson, A.: Hydroxamate siderophores: occurrence and importance in the Atlantic Ocean, *Environ. Sci. Tech.*, 42, 8675–8680, <https://doi.org/10.1021/es801884r>, 2008.
- Mawji, E., Gledhill, M., Milton, J. A., Zubkov, M. V., Thompson, A., and Wolff, G. A.: Production of siderophore type chelates in Atlantic Ocean waters enriched with different carbon and nitrogen sources, *Mar. Chem.*, 124, 90–99, <https://doi.org/10.1016/j.marchem.2010.12.005>, 2011.
- Martin, J. H., Fitzwater, S. E., and Gordon, R. M.: Iron deficiency limits phytoplankton growth in Antarctic waters, *Glob. Biogeochem. Cy.*, 4, 5–12, 1990.
- Middag, R., Séférian, R., Conway, T. M., John, S. G., Bruland, K. W., and de Baar, H. J. W.: Intercomparison of dissolved trace elements at the Bermuda Atlantic Time Series station, *Mar. Chem.*, 177, 476–489, 2015.
- Miller, L. A. and Bruland, K. W.: Competitive equilibration techniques for determining transition metal speciation in natural waters: Evaluation using model data, *Anal. Chim. Act.*, 343, 161–181, 1997.
- Morrissey, J. and Bowler, C.: Iron utilization in marine cyanobacteria and eukaryotic algae, *Front. Microbiol.*, 3, 43, <https://doi.org/10.3389/fmicb.2012.00043>, 2012.
- Pham, A. L.-D. and Ito, T.: Ligand binding strength explains the distribution of iron the Atlantic Ocean, *Geophys. Res. Lett.*, 47, 7500–7508, doi:10.1029/2019BL083319, 2019.
- Pižeta, I., Sander, S. G., Hudson, R. J. M., Baars, O., Barbeau, K. A., Buck, K. N., Bundy R. M., Carrasco, G., Croot, P. L., Garnier, C., Gerringa, L. J. A., Gledhill, M., Hirose, K., Kondo, Y., Laglera, L. M., Nuester, J., Omanović, D., Rijkenberg, M. J. A., Takeda, S., Twining, B. S., and Wells, M.: Quantitative analysis of complexometric titration data: An intercomparison of methods for estimating models of metal complexation by mixtures of natural ligands, *Mar. Chem.*, 173, 3–24, 2015.
- Purawatt, S., Siripinyanond, A., and Shiwatana, J.: Flow field-flow fractionation-inductively coupled optical emission spectrometric investigation of the size-based distribution of iron complexed to phytic and tannic acids in a food suspension: implications for iron availability, *Anal. Bioanal. Chem.*, 389, 733–742, 2007.
- Rapp, I., Schlosser, C., Rusiecka, D., Gledhill, M., and Achterberg, E. P.: Automated preconcentration of Fe, Zn, Cu, Ni, Cd, Pb, Co, and Mn in seawater with analysis using high-resolution sector field inductively-coupled plasma mass spectrometry, *Anal. Chim. Act.*, 976, 1–13, <https://doi.org/10.1016/j.aca.2017.05.008>, 2017.
- Rose, A. L. and Waite, T. D.: Kinetics of iron complexation by dissolved natural organic matter in coastal waters, *Mar. Chem.*, 84, 85–103, 2003.
- Rue, E. L. and Bruland, K. W.: Complexation of iron(III) by natural organic ligands in the Central North Pacific as determined by a new competitive ligand equilibration/adsorptive cathodic stripping voltammetric method, *Mar. Chem.*, 50, 117–138, 1995.
- Rue, E. L. and Bruland, K. W.: The role of organic complexation on ambient iron chemistry in the Equatorial Pacific Ocean and the response of a mesoscale iron addition experiment, *Limnol. Oceanogr.*, 42, 901–910, 1997.
- Rijkenberg, M. J. A., Gerringa, L. J. A., Carolus, V. E., Velzeboer, I., and de Baar, H. J. W.: Enhancement and inhibition of iron photoreduction by individual ligands in open ocean seawater, *Geochim. Cosmochim. Acta.*, 70, 2790–2805, <https://doi.org/10.1016/j.gca.2006.03.004>, 2006.
- Rijkenberg, M. J. A., Middag, R., Laan, P., Gerringa, L. J. A., van Aken, H., Schoemann, V., de Jong, J. T. M., and de Baar, H. J. W.: The distribution of dissolved iron in the West Atlantic Ocean, *PLoS ONE*, 9, e101323, <https://doi.org/10.1371/journal.pone.0101323>, 2014.
- Rijkenberg, M. J. A., Slagter, H. A., Rutgers van der Loeff, M., van Ooijen, J., and Gerringa, L. J. A.: Sources of dissolved Fe in the deep and upper Arctic Ocean with a focus on Fe limitation in the Nansen Basin, *Front. Mar. Sci.*, 5, 88, <https://doi.org/10.3389/fmars.2018.00088>, 2018.
- Salmon, T. P., Rose, A. L., Neilan, B. A., and Waite, T. D.: The FeL model of iron acquisition: Non dissociative reduction of ferric complexes in the marine environment, *Limnol. Oceanogr.*, 51, 1744–1754, 2006.
- Schlitzer, R., Anderson, R. F., Masferrer Dodas, E., et al.: The GEOTRACES Intermediate Data Product 2017, *Chem. Geol.*, <https://doi.org/10.1016/j.chemgeo.2018.05.040>, 2018.
- Schijf, J. and Burns, S. M.: Determination of side-reaction coefficient of desferrioxamine B in trace metal free seawater, *Front. Mar. Sci.*, 3, 117, <https://doi.org/10.3389/fmars.201600117>, 2016.
- Schlosser, C. and Croot, P. L.: Application of cross-flow filtration for determining the solubility of iron species in open ocean seawater, *Limnol. Oceanogr.-Meth.*, 6, 630–642, 2008.
- Shaked, Y., Kustka, A. B., and Morel, F. M. M.: A general kinetic model for iron acquisition by eukaryotic phytoplankton, *Limnol. Oceanogr.*, 50, 872–882, 2005.

- Shaked, Y., Buck, K. N., Mellett, T., and Maldonado, M. T.: Insights into the bioavailability of oceanic dissolved Fe from phytoplankton uptake kinetics, *ISME J.*, 14, 1182–1193, <https://doi.org/10.1038/s41396-020-0597-3>, 2020.
- Shaked, Y., Twining, B. S., Tagliabue, A., and Maldonado, M. T.: Probing the bioavailability of dissolved iron to marine eukaryotic phytoplankton using in situ single cell iron quotas, *Biogeochem. Cy.*, 35, 006979, <https://doi.org/10.1029/2021GB006979>, 2021.
- Slagter, H. A., Reader, H. E., Rijkenberg, M. J. A., Rutgers van der Loeff, M., de Baar, H. J. W., and Gerringa, L. J. A.: Fe speciation is related to terrestrial dissolved organic matter in the Arctic Ocean, *Mar. Chem.*, 197, 11–25, <https://doi.org/10.1016/j.marchem.2017.10.005>, 2017.
- Slagter, H. A., Laglera, L. M., Sukekava, C., and Gerringa, L. J. A.: Fe-binding organic ligands in the humic-rich Trans-Polar Drift in the surface Arctic Ocean using multiple voltammetric methods, *J. Geophys. Res.-Oceans*, 124, 14576, <https://doi.org/10.1029/2018JC014576>, 2019.
- Stockdale, A., Tipping, E., and Lofts, S.: Dissolved trace metal speciation in estuarine and coastal waters: Comparison of WHAM/Model VII predictions with analytical results, *Environ. Toxicol. Chem.*, 34, 53–63, <https://doi.org/10.1002/etc.2789>, 2015.
- Su, H., Yang, R., Li, Y., and Wang, X.: Influence of humic substances on iron distribution in the East China Sea, *Chemosphere*, 204, 450–462, 2018.
- Sukekava, C., Downes, J., Slagter, H. A., Gerringa, L. J. A., and Laglera, L. M.: Determination of the contribution of humic substances to iron complexation in seawater by catalytic cathodic stripping voltammetry, *Talanta*, 189, 359–364, <https://doi.org/10.1016/j.talanta.2018.07.021>, 2018.
- Sutten, R. and Sposito, G.: Molecular Structure in Soil Humic Substances: The New View, *Environ. Sci. Technol.*, 39, 9009–9015, 2005.
- Tagliabue, A. and Völker, C.: Towards accounting for dissolved iron speciation in global ocean models, *Biogeosciences*, 8, 3025–3039, <https://doi.org/10.5194/bg-8-3025-2011>, 2011.
- Tagliabue, A., Aumont, O., DeAth, R., Dunne, J.P., Dutkiewicz, S., Galbraith, E., Misumi, K., J., Moore, K. J. K., Ridgwell, A., Sherman, E., Stock, C., Vichi, M., Völker, C., and Yool, A.: How well do global ocean biogeochemistry models simulate dissolved iron distributions?, *Global Biogeochem. Cy.*, 30, 149–174, <https://doi.org/10.1002/2015GB005289>, 2016.
- Thuróczy, C.-E., Gerringa, L. J. A., Klunder, M., Laan, P., le Guitton, M., and de Baar, H. J. W.: Distinct trends in the speciation of iron between the shelf seas and the deep basins of the Arctic Ocean, *J. Geophys. Res.*, 116, C10009, <https://doi.org/10.1029/2010JC006835>, 2011a.
- Thuróczy, C.-E., Gerringa, L. J. A., Klunder, M., Laan, P., and de Baar, H. J. W.: Observation of consistent trends in the organic complexation of dissolved iron in the Atlantic sector of the Southern Ocean, *Deep Sea Res. Pt. II*, 58, 2695–2706, 2011b.
- Thuróczy, C.-E., Alderkamp, A.-C., Laan, P., Gerringa, L. J. A., de Baar, H. J. W., and Arrigo, K. R.: Key role of organic complexation of iron in sustaining phytoplankton blooms in the Pine Island and Amundsen Polynyas (Southern Ocean), *Deep Sea Res. Pt. II*, 71, 49–60, 2012.
- Town, R. M. and Filella, M.: Dispelling the myths: is the existence of L1 and L2 ligands necessary to explain metal ion speciation in natural waters?, *Limnol. Oceanogr.*, 45, 1341–1357, <https://doi.org/10.4319/lo.2000.45.6.1341>, 2000.
- Town, R. M., Chakraborty, P., van Leeuwen, H. P.: Dynamic DGT speciation analysis and applicability to natural heterogeneous complexes, *Environ. Chem.*, 6, 170–177, <https://doi.org/10.1071/EN08091>, 2009.
- Town, R. M. and van Leeuwen, H. P.: Labilities of aqueous nanoparticulate metal complexes in environmental speciation analysis, *Environ. Chem.*, 11, 196–205, <https://doi.org/10.1071/EN13138>, 2014.
- Turoczy, N. J. and Sherwood, J. E.: Modification of the van den Berg–Ruzic method for the investigation of complexation parameters of natural waters, *Anal. Chim. Act.*, 354, 15, [https://doi.org/10.1016/S0003-2670\(97\)00455-8](https://doi.org/10.1016/S0003-2670(97)00455-8), 1997.
- van den Berg, C. M. G.: Determination of copper complexation with natural organic ligands in seawater by equilibration with MnO₂ I., Theory, *Mar. Chem.*, 11, 307–322, 1982.
- van den Berg, C. M. G.: Evidence for organic complexation of iron in seawater, *Mar. Chem.*, 50, 139–157, 1995.
- van den Berg, C. M. G.: Chemical speciation of iron in seawater by cathodic stripping voltammetry with dihydroxynaphthalene, *Anal. Chem.*, 78, 15, 156–163, 2006.
- van den Berg, C. M. G. and Donat, J. R.: Determination and data evaluation of copper complexation by organic ligands in sea water using cathodic stripping voltammetry at varying detection windows, *Anal. Chim. Act.*, 257, 281–291, [https://doi.org/10.1016/0003-2670\(92\)85181-5](https://doi.org/10.1016/0003-2670(92)85181-5), 1992.
- Velasquez, I. B., Ibisami, E., Maas, E. W., Boyd, P. W., Nodder, S., and Sander, S. G.: Ferrioxamine siderophores detected amongst iron binding ligands produced during the remineralization of marine particles, *Front. Mar. Sci.*, 3, 172, <https://doi.org/10.3389/fmars.2016.00172>, 2016.
- Völker, C. and Tagliabue, A.: Modeling organic iron-binding ligands in a three-dimensional biogeochemical ocean model, *Mar. Chem.*, 173, 67–77, 2015.
- Wells, M., Buck, K. N., and Sander, S. G.: New approach to analysis of voltammetric ligand titration data improves understanding of metal speciation in natural waters, *Limnol. Oceanogr.-Meth.*, 11, 450–465, <https://doi.org/10.4319/lom.2013.11.450>, 2013.
- Whitby, H. Planquette, H., Cassar, N., Bucciarelli, E., Osburn, C. L., Janssen, D. J., Cullen, J. T., González, A. G., Völker, C., and Sarthou, G.: A call for refining the role of humic-like substances in the oceanic iron cycle, *Sci. Rep.*, 10, 6144, [doi:10.1038/s41598-020-62266-7](https://doi.org/10.1038/s41598-020-62266-7), 2020.
- Witter, A. E., Hutchins, D. A., Butler, A., and Luther, G. W.: Determination of conditional stability constants and kinetic constants for strong model Fe-binding ligands in seawater, *Mar. Chem.*, 69, 1–17, 2000.
- Wu, J. F. and Luther, G. W.: Complexation of Fe(III) by natural organic-ligands in the northwest Atlantic Ocean by a competitive ligand equilibration method and a kinetic approach, *Mar. Chem.*, 50, 159–177, 1995.
- Wuttig, K., Heller, M. I., and Croot, P. L.: Reactivity of inorganic Mn and Mn desferrioxamine B with O₂, O₂–, and H₂O₂ in seawater, *Environ. Sci. Technol.*, 47, 10257–10265, <https://doi.org/10.1021/es4016603>, 2013.
- Wuttig, K., Townsend, A. T., van der Merwe, P., Gault-Ringold, M., Holmes, T., Schallenberg, C., Latour, P., Tonnard, M., Rijkenberg, M. J. A., Lannuzel, D., and Bovei, A. R.:

- Critical evaluation of a seaFAST system for the analysis of trace metals in marine samples, *Talanta*, 197, 653–668, <https://doi.org/10.1016/j.talanta.2019.01.047>, 2019.
- Yang, R. and Ven den Berg, M. C. G.: Metal complexation by humic substances in seawater, *Environ. Sci. Technol.*, 43, 7192–7197, 2009.
- Yang, R., Su, H., Qu, S., and Wang, X.: Capacity of humic substances to complex with iron at different salinities in the Yangtze River estuary and East China Sea, *Sci. Rep.*, 7, 1381, <https://doi.org/10.1038/s41598-017-01533-6>, 2017.
- Yamashita, Y., Nishioka, J., and Hajime, O.: Shelf humic substances as carriers for basin-scale iron transport in the North Pacific, *Sci. Rep.*, 10, 4505, <https://doi.org/10.1038/s41598-020-61375-7>, 2020.
- Ye, Y. and Völker, C.: On the role of dust-deposited lithogenic particles for iron cycling in the tropical and subtropical Atlantic, *Global Biogeochem. Cy.*, 31, 1543–1558, <https://doi.org/10.1002/2017GB005663>, 2017.
- Ye, Y., Völker, C., and Gledhill, M.: Exploring the iron-binding potential of the ocean using a combined pH and DOC parameterization, *Global Biogeochem. Cy.*, 34, e2019GB006425, <https://doi.org/10.1029/2019GB006425>, 2020.
- Zhang, H. and Davison, W.: Use of diffusive gradients in thin-films for studies of chemical speciation and bioavailability, *Environ. Chem.*, 12, 85–101, <https://doi.org/10.1071/EN14105>, 2015.
- Zhu, K., Hopwood, M. J., Groenenberg, J. E., Engel, A., Achterberg, E. P., and Gledhill, M.: Influence of pH and dissolved organic matter on iron speciation and apparent iron solubility in the Peruvian upwelling region, *Environ. Sci. Technol.*, 55, 9372–9383, 2021.