





Prochlorococcus extracellular vesicles: molecular composition and adsorption to diverse microbes

Steven J. Biller ^{1,2*} Rachel A. Lundeen ³
Laura R. Hmelo,³ Kevin W. Becker ^{4†}
Aldo A. Arellano,¹ Keven Dooley,¹ Katherine R. Heal ^{3‡}
Laura T. Carlson,³ Benjamin A. S. Van Mooy,⁴
Anitra E. Ingalls³ and Sallie W. Chisholm^{1§}

¹Department of Civil and Environmental Engineering, Massachusetts Institute of Technology, Cambridge, MA.

²Department of Biological Sciences, Wellesley College, Wellesley, MA.

³School of Oceanography, University of Washington, Seattle, WA.

⁴Department of Marine Chemistry and Geochemistry, Woods Hole Oceanographic Institution, Woods Hole, MA.

Summary

Extracellular vesicles are small (~50–200 nm diameter) membrane-bound structures released by cells from all domains of life. While vesicles are abundant in the oceans, their functions, both for cells themselves and the emergent ecosystem, remain a mystery. To better characterize these particles – a prerequisite for determining function – we analysed the lipid, protein, and metabolite content of vesicles produced by the marine cyanobacterium *Prochlorococcus*. We show that *Prochlorococcus* exports a diverse array of cellular compounds into the surrounding seawater enclosed within discrete vesicles. Vesicles produced by two different strains contain some materials in common, but also display numerous strain-specific differences, reflecting functional complexity within vesicle populations. The vesicles contain active enzymes, indicating that they can mediate extracellular biogeochemical reactions in the ocean. We further demonstrate that vesicles from *Prochlorococcus* and other bacteria associate with diverse microbes including the most abundant marine bacterium, *Pelagibacter*. Together, our data

point toward hypotheses concerning the functional roles of vesicles in marine ecosystems including, but not limited to, possibly mediating energy and nutrient transfers, catalysing extracellular biochemical reactions, and mitigating toxicity of reactive oxygen species.

Introduction

Many, if not all, bacteria release extracellular vesicles from their surface into the local environment (Deatherage and Cookson, 2012). In exponentially growing Gram-negative bacteria, these structures are thought to derive primarily from the outer membrane, wherein a local region of membrane separates from the cell, carrying with it periplasmic material and other cellular components (Schwechheimer and Kuehn, 2015). A small subset of vesicles from Gram-negative cells may include both outer and inner membrane material as well, further expanding the range of potential vesicle contents (Pérez-Cruz *et al.*, 2015). Vesicles are released constitutively during growth, but release rates can also vary in response to environmental perturbations (MacDonald and Kuehn, 2013; Biller *et al.*, 2014). Extracellular vesicles represent a versatile secretion mechanism for cells (Schwechheimer and Kuehn, 2015; Guerrero-Mandujano *et al.*, 2017), and many classes of cellular compounds, including proteins, nucleic acids, and small molecules, have been identified within bacterial vesicles (Schwechheimer and Kuehn, 2015; Brown *et al.*, 2015). Since they are bounded by a lipid bilayer membrane, vesicles also provide a mechanism for secreting and transporting hydrophobic compounds through aqueous extracellular environments (Mashburn-Warren and Whiteley, 2006).

Extracellular vesicles can shuttle their contents between cells (Kadurugamuwa and Beveridge, 1996; Yaron *et al.*, 2000). This ability enables vesicles to mediate a wide variety of biological functions such as horizontal gene transfer, signalling, pathogenesis, quorum signalling, biofilm development, nutrient exchange, viral interactions, and cellular defence (Kadurugamuwa and Beveridge, 1996; Yaron *et al.*, 2000; MacDonald and Kuehn, 2012; Schwechheimer and Kuehn, 2015; Lynch and Alegado, 2017; Schatz *et al.*, 2017). Although such

Received 18 December, 2020; accepted 29 October, 2021. *For correspondence. E-mail sbiller@wellesley.edu. Present addresses: †GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel; ‡Integral Consulting Inc, Seattle, WA; §Department of Biology, Massachusetts Institute of Technology, Cambridge, MA.

exchanges have been shown to occur both among bacteria and across domains in a few laboratory models, the 'rules' dictating these exchanges are not at all clear, and essentially nothing is known about what occurs between cells and vesicles in complex microbial systems such as those found in the oceans, soils, or within the human microbiome.

Extracellular vesicles can reach concentrations of $>10^5$ and $>10^6$ ml⁻¹ in open ocean and coastal waters, respectively (Biller *et al.*, 2014) and represent an entirely new dimension of dissolved organic carbon pools. Their biological and ecological roles are unknown, but we do know that they are released by diverse taxa of both autotrophic and heterotrophic marine microbes, including the abundant cyanobacterium *Prochlorococcus* (Biller *et al.*, 2014). With a global population of $\sim 3 \times 10^{27}$ cells, *Prochlorococcus* is an important primary producer, responsible for nearly 10% of marine net primary production (Flombaum *et al.*, 2013). This group is known to secrete a number of organic compounds (Bertilsson *et al.*, 2005) that provide marine heterotrophs with a source of carbon and energy (Ottesen *et al.*, 2014; Becker *et al.*, 2019) and contribute to marine dissolved organic carbon pools. *Prochlorococcus* extracellular vesicles represent at least one component of this labile organic photosynthate, as purified vesicles have been shown to support the growth of a marine heterotroph (Biller *et al.*, 2014).

Here, we explore the contents and function of extracellular vesicles produced by *Prochlorococcus*, focusing on their potential contribution to ocean dissolved organic matter and their ability to interact with other marine microbes. We characterized the lipidome, proteome, and metabolome of vesicles released by two ecologically distinct strains and use these inventories to develop hypotheses of vesicle functions. To establish whether *Prochlorococcus* vesicles can chemically interact with surrounding seawater, we determined whether they contain active enzymes. Finally, we investigated the potential for *Prochlorococcus* vesicles to mediate biotic interactions by studying whether they can form specific associations with other strains representative of abundant marine microbes.

Results and discussion

Diverse biomolecules are associated with Prochlorococcus extracellular vesicles

We used a combination of targeted and untargeted 'omics' approaches to examine the contents of vesicles and the cells that released them. Our study focused on vesicles isolated from exponentially growing, asynchronous cultures of two ecologically distinct *Prochlorococcus* – MIT9312, a high-light adapted strain, and MIT9313, a

low-light adapted strain (Biller *et al.*, 2015a,b). We uncovered a vast diversity of biomolecules in these discrete, lipid-bound, colloidal packets, as detailed below.

Lipids and pigments. Lipids from vesicles and cells of both *Prochlorococcus* strains were analysed using an untargeted, high-resolution mass spectrometry-based approach. Vesicles contained a suite of different intact polar lipids (IPLs), various pigments, and plastoquinone. Overall, vesicles were relatively enriched for IPLs versus pigments, as compared with the parent cells (Fig. 1A; Supporting Information Table S1). The carotenoids zeaxanthin and carotene were the most abundant pigments in vesicles from both strains, which contained relatively few chloropigments such as divinyl chlorophylls *a* and *b* (Fig. 1B; Supporting Information Table S1). While plastoquinone was identified in the vesicles (Fig. 1B; Supporting Information Table S1), they contained little thylakoid material overall, suggesting that vesicle carotenoids could come from other cellular locations such as the outer membrane, as has been shown for *Synechocystis* PCC6714 (Jürgens and Weckesser, 1985).

Prochlorococcus cellular membranes are primarily composed of sulfolipids, reflecting an adaptation to the low phosphorous oligotrophic environment in which they live (Van Mooy *et al.*, 2006). This was evident in the lipid composition of its extracellular vesicles as well. The IPL composition of *Prochlorococcus* extracellular vesicles was dominated by sulfoquinovosyl diacylglycerol (SQDG), diglycosyl diacylglycerol (DGDG) and monoglycosyl diacylglycerol (MGDG) – each found in similar relative abundances (as a proportion of all IPLs) in cells and vesicles (Fig. 1C; Supporting Information Table S1). The overall vesicle lipid composition of each strain more closely resembled the composition of the parent cells than the vesicles from the other strain (Figs. 1C and S1), consistent with our previous observations from *Prochlorococcus* strains MIT9313 and MED4 (Biller *et al.*, 2014) suggesting that there is not a universal *Prochlorococcus* vesicle lipidome. Vesicles from the two strains did, however, share some common lipid features that distinguished them from the parental cells. Of note, hydroxylated forms of all major IPLs identified (polar lipids modified by oxidation reactions) were relatively more abundant in vesicle samples as compared with cells (Fig. 1D; Supporting Information Table S1). In addition, the fraction of some hydroxylated IPLs (DGDG in MIT9312; SQDG and MGDG in MIT9313) among biological replicate vesicle samples was significantly more variable than in the corresponding cellular samples (*F*-test, $P < 0.05$; Figs. 1D and S1B–D). Annotation of these hydroxylated lipids revealed that the hydroxylation is site specific, for example, at the $\Delta 9$ -position of a C_{18:2} fatty acid (Supporting Information Fig. S2) regardless of head




Fig. 1. Lipid and pigment content of *Prochlorococcus* cells and vesicles from two strains, MIT9312 and MIT9313.

A. Relative abundance of pigments vs intact polar lipids (IPLs) in vesicles and cells.

B. Relative abundance of specific pigments and plastoquinone.

C. Relative abundance of different IPL groups.

D. Fraction of hydroxylated polar lipids (OH-IPLs) within each IPL class. Values indicate the mean (\pm SD) of three biological replicates of strains MIT9312 (white) and MIT9313 (black). MGDG: monoglycosyl diacylglycerol; DGDG: diglycosyl diacylglycerol; PG: phosphatidylglycerol; SQDG: sulfoquinovosyl diacylglycerol. * indicates significant differences between cells and vesicles (two-tailed *t* test, $P < 0.05$).

group. The exact oxidative mechanism (enzymatic, radical mediated, or non-radical mediated) responsible for generating the hydroxylation is not known, but free-radical (auto) oxidation at a specific double bond would be expected to result in at least four isomers of similar abundance (Rontani and Belt, 2020). While lipid hydroxylation could have occurred either before or after vesicle release, the site-specificity raises the possibility that they could be of biological origin. One potential explanation is that *Prochlorococcus* outer membranes are generally

enriched in hydroxylated lipids relative to other membranes, as has been shown for other Gram-negative bacteria (Schmidt *et al.*, 1980; Volkman *et al.*, 1998). This would be consistent with their enrichment in outer membrane-derived vesicles and perhaps reflect a role for vesicle secretion as a mechanism for removing damaged lipids from the cell. Hydroxy lipids may also be involved in the membrane curvature and bending processes required for vesicle formation, leading to their preferential incorporation and export.

Proteins. Since many of the functional roles attributed to vesicles in other microbial systems are due to the activity of proteins (Schwechheimer and Kuehn, 2015), we explored the global proteomes of the vesicle and cellular fractions of two *Prochlorococcus* strains using a label-free, quantitative shotgun proteomics approach. During extraction and sample preparation, we utilized surfactants compatible with mass spectrometry to improve the recovery of more membrane-bound proteins and better facilitate in-solution protease digestion, which further helped generate more comprehensive vesicle and cellular proteomes. MIT9312 and MIT9313 vesicles contained, respectively, at least 11% and 12.5% of all predicted proteins encoded by the cell's genome (Table 1). This most likely does not represent every protein found in vesicles, but rather what could be detected within the relatively small amount of vesicle biomass we could obtain from 20 l cultures as compared with cellular material (c.f., cellular proteomes for MIT9312 and MIT9313 recovered nearly 52% and 37% of all predicted proteins, respectively; Table 1). That said, these vesicle proteomes yielded notably more protein identifications than our previous study of MIT9313 and MED4 vesicles (in which gel-extracted protein bands were analysed; Biller *et al.*, 2014), and it is noteworthy that 25 of the 27 proteins we previously identified in MIT9313 were also present in this data set. Further, those 25 were among the top 50 most highly abundant proteins in the MIT9313 vesicle proteome (Supporting Information Tables S2 and S3). The concordance of these results shows that *Prochlorococcus* cells grown under similar conditions will reproducibly package and export intact proteins within vesicles.

Vesicles produced by the two *Prochlorococcus* strains contained some proteins in common (Tables 1 and S2), but we also noted strain-specific differences, as has been found in comparisons of vesicles from other closely related microbes (Tandberg *et al.*, 2016; Bitto *et al.*, 2021). The overall *Prochlorococcus* vesicle proteome included proteins associated with many functional categories (Figs. 2A, S3). For instance, both MIT9312 and MIT9313 vesicles contained putative porins and transporters (e.g. PMT9312_1131, Som), peptidases/

hydrolases (e.g. PMT9312_0677, PMT_1636), chaperones (e.g. DnaK and DnaJ), and uncharacterized proteins (Fig. S3; Table S4). The relative abundance of these shared proteins did, however, vary between strains (Fig. 2A; Tables S2–S4). Proteins uniquely found in MIT9312 vesicles included a cAMP phosphodiesterase (PMT9312_0858), an adhesin-like protein (PMT9312_1179), a phosphate ABC transporter substrate-binding protein (PstB), and some ribosomal proteins. Among those found uniquely in MIT9313 were multiple ABC transporter binding proteins (e.g. UrtA, FutA1, PMT_2203), a sulfatase (PMT_1515), and a putative phosphatase (PMT_1619) (Fig. 2A; Tables S2 and S3). Thus, the vesicles released by two relatively similar organisms have distinct functional potentials.

The functional potential of vesicles will also be influenced by the relative abundance of any individual protein within the vesicle population. Indeed, different proteins were packaged at different levels within the vesicles in our samples, with a subset enriched (on a biovolume-normalized basis) in vesicles relative to cells (Figs. 2A, S4; Table S4). To explore what factors might influence the packaging, we examined whether subcellular localization correlated with vesicle export. Consistent with a primarily outer membrane origin for extracellular vesicles, most outer membrane and periplasmic proteins detected in the cellular proteome were identified at some level within vesicles (Fig. S5). As seen in other Gram-negative bacteria (Pérez-Cruz *et al.*, 2015; Zakharchevskaya *et al.*, 2017; Yun *et al.*, 2017), we found proteins from all major cellular regions to be exported within *Prochlorococcus* vesicles (Fig. 2B). We were, however, surprised to find that cytoplasmic proteins made up the largest fraction in strain MIT9312, whereas most proteins in MIT9313 vesicles were predicted to originate from various locations outside the cytoplasm (Fig. 2B). Although a number of abundant cytoplasmic proteins, such as ribosomal proteins, were found in MIT9312 vesicles (Fig. 2A; Tables S2 and S3), there was no general relationship between their abundance in the cells and vesicles (Supporting Information Fig. S6A). Together with the relative lack of thylakoid proteins and chlorophyll in the vesicles, this argues that the presence

Table 1. Number of proteins identified in *Prochlorococcus* cells and vesicles.

Strain	Annotated proteins encoded in genomes (Uniprot)	Annotated proteins shared by both genomes	Proteins detected in cells ^a	Proteins detected in cells of both strains	Proteins detected in vesicles	Proteins detected only in vesicles and not in cells	Proteins detected in vesicles from both strains
MIT9312	1964	1532	1030	714	224	17	111
MIT9313	2830		1055		355	41	

^aThese values reflect the data gathered for the trypsin/Lys-C digestion of *Prochlorococcus* cells and vesicles (Supporting Information Tables S2 and S3); not included here, additional cellular proteins were identified using Glu-C digestion.




Fig. 2. *Prochlorococcus* vesicle proteomes.

A. Relative protein enrichment in *Prochlorococcus* vesicles compared with cells. Points represent the log₂ ratio of relative protein abundance for the top 25% most abundant proteins identified in vesicles of strain MIT9312 (above; orange) or MIT9313 (below; blue), as grouped by NCBI clusters of orthologous groups of proteins (COG) functional categories. The area of each point indicates biovolume-normalized spectral counts (abundance) of that protein within the vesicle proteome. Names or locus tags are indicated for selected proteins; data for all vesicle proteins are found in the Supporting Information Table S4.

B. Relative abundance of proteins found in MIT9312 (orange) and MIT9313 (blue) vesicles, as grouped by predicted subcellular localization (see Methods).

of cytosolic proteins is likely not due to artefacts such as cell lysis. We also saw no clear relationship between a protein's presence in the vesicles and properties such as isoelectric point or molecular weight (Supporting Information Fig. S6B–E).

A number of factors likely contribute to the differences observed between the vesicle proteomes of the two

strains. Some differences are simply attributable to strain-specific genomic differences (Table 1). While the two genomes share over 1500 genes, MIT9313 encodes ~870 more genes than does MIT9312, and its vesicles contained a proportionally more diverse set of proteins. One notable example was the identification of several prochlorosins, or ProcA peptides, within the vesicle

proteome of MIT9313 (Supporting Information Table S2). Prochlorosins are cyclic peptide secondary metabolites encoded by some low-light adapted, but not high-light adapted, *Prochlorococcus* – including 29 diverse *procA* genes in MIT9313 (Li *et al.*, 2010; Cubillos-Ruiz *et al.*, 2017). Strain-specific genome differences do not, however, explain all of the vesicle proteome variation. Even when considering only the proteins shared by both strains, we found that the relative abundance of orthologous proteins was linearly correlated in the whole cell fraction, but that there was no clear relationship in the relative amounts of these proteins within extracellular vesicles (Supporting Information Fig. S4B). We speculate that differences in cellular structure between the strains (Ting *et al.*, 2007) may have influenced the differential incorporation of proteins from the cytosol and other sub-cellular compartments into vesicles.

While we cannot rule out the possibility that some individual proteins were specifically packaged via unknown mechanisms, our data suggest that most proteins are packaged stochastically, with the probability of export influenced by a number of factors including initial cellular protein abundance, subcellular localization, and perhaps strain-specific differences in cellular architecture. Physical constraints, such as the amount of membrane relative to the aqueous lumen of the vesicle, may also influence protein composition and limit the enrichment of some compounds in vesicles relative to cells (Turner *et al.*, 2018). Finally, it is likely that additional sampling would reveal an even longer ‘tail’ of other proteins that are either exported at relatively lower abundances or were perhaps lost through the purification protocols used here. Thus, we suspect that most (or perhaps all) proteins expressed within cells will eventually be found

within vesicle populations. Future work examining vesicles across strains (Tartaglia *et al.*, 2020; Zwarycz *et al.*, 2020) and under different growth conditions (Orench-Rivera and Kuehn, 2016; Zavan *et al.*, 2019) will be needed to advance our understanding of the mechanisms influencing vesicle proteome composition, and how such differences might affect the emergent functional impact of those vesicles on the ecosystem.

Metabolites. A combination of targeted and untargeted metabolomics analysis identified 1662 and 2035 unique mass features in vesicle populations from *Prochlorococcus* MIT9312 and MIT9313, respectively (Supporting Information Table S5). These mass features represented molecules across a range of *m/z* values (~94 to ~800), polarities, and charge states (Supporting Information Table S6). Most vesicle-associated metabolites detected were non-polar (90% and 85% in MIT9312 and MIT9313 vesicles, respectively), whereas non-polar metabolites represented 52%–56% of mass features detected in the cells (Tables S5 and S6). The identification of diverse non-polar molecules in vesicles further supports the idea that extracellular vesicles can serve as vehicles for the secretion of hydrophobic compounds (Mashburn and Whiteley, 2005; Schertzer *et al.*, 2009), facilitating biological activities that depend on extracellular transport of nonpolar molecules within aqueous environments.

While we are not able to identify most of the observed mass features, we could document a number of known compounds within the vesicles (Table 2). For instance, vesicles from both strains contained phyloquinone (Vitamin K1), a compound involved in electron transfer. We also identified a number of oxidized carotenoid

Table 2. Non-intact polar lipid metabolites identified within *Prochlorococcus* vesicles.

Compound name	Fraction	Compound class	MIT9312 vesicles	MIT9313 vesicles	Confidence level	<i>m/z</i>	RT (min)
Triose	HILICAqNeg	Sugar	+	+	3a	503.1629	13.6
Tetraose	HILICAqNeg	Sugar	+	+	3a	665.2164	14.4
Phylloquinone (Vitamin K1)	RPOrgPos	Electron transfer	+	+	1	451.3572	14.6
Carotene	RPOrgPos	Pigment	+	+	1	536.4371	16.1
Protoporphyrin	RPOrgPos	Pigment or cyt-c precursor	+	+	2	563.2656	12.8
1-Lauroyl-glycerol (C12)	RPOrgPos	Lipid	+	+	2	257.2111	11.7
1-Myristoyl-glycerol (C14)	RPOrgPos	Lipid	+	+	2	285.2423	13.2
Beta-Apo-12'-carotenal	RPOrgPos	Pigment oxidation product	+	+	3b	351.2682	13.5
Apo-12'-zeaxanthinal	RPOrgPos	Pigment oxidation product	+	+	3b	367.2633	12.0
Beta-Apo-10'-carotenal	RPOrgPos	Pigment oxidation product	+	+	3b	377.2839	13.8
Apo-10'-zeaxanthinal	RPOrgPos	Pigment oxidation product	+	+	3b	393.2785	12.4

For details on analytical fractions and confidence level assessments, see Methods and Supplementary Information. + = detected.

products (Table 2) in vesicles from both strains, though it is unclear whether these carotenoids were oxidized in the cell and then exported via vesicles or whether they became oxidized following vesicle release. Although we focused our analysis on the compounds present in vesicles, cellular compounds absent in vesicles are also of interest (Table S7). For example, *Prochlorococcus* cells contained potential osmolytes such as sucrose, glycine betaine, or glucosylglycerol, which help maintain cellular osmotic balance in cyanobacteria (Klähn and Hagemann, 2010), but these compounds were undetectable in either set of vesicles (Supporting Information Table S7). This represents a puzzle as to how cytoplasmic material could be exported to vesicles without including (or retaining) the most abundant solutes in the cell and raises questions concerning how vesicles may change in response to osmotic stresses.

Nutrient and energy sources within vesicles

Prochlorococcus vesicles have been shown to serve as sources of organic carbon and/or energy for supporting growth of co-cultured bacteria (Biller et al., 2014). Much of a vesicle's carbon and chemical energy content is found within the lipids, as each 100 nm diameter vesicle will contain ~10 000 lipid molecules in its envelope. The diverse proteins and metabolites exported within vesicles represent potential labile organic carbon sources as well. In one notable example, vesicles from both strains contained triose and tetrose sugars (Table 2), which were not measurable above background levels within cells. While not the focus of this work, *Prochlorococcus* vesicles contain DNA and RNA (Biller et al., 2014) as well, and vesicle-associated nucleic acids may also function as an energy or nutrient source for microbes (Jørgensen and Jacobsen, 1996).

We previously observed nutrient binding proteins, such as those for urea, phosphate, and iron, in the vesicles exported by *Prochlorococcus* strains MED4 and MIT9313 (Biller et al., 2014). This suggested that vesicles might transport nutrients or even scavenge compounds as they diffuse through seawater and, in turn, organisms that encounter these vesicles could gain access to a locally concentrated 'packet' of nutrients. To continue to explore this hypothesis, we looked for annotated substrate-binding proteins and transporters in the proteome data. We again identified putative substrate-binding proteins (many associated with ABC transporters) for urea, phosphate, and iron in the vesicles, along with those for manganese and amino acids (Supporting Information Tables S2 and S3). Vesicles produced by both strains contained putative transporters for sulfate, magnesium, and ammonium (Supporting Information Tables S2 and S3), raising the possibility that – assuming

favourable energetics and proper protein orientation in the membrane – vesicles could accumulate these compounds following cellular release.

As a first step toward determining whether the nutrients that the proteomic data suggest might be transported by vesicles are actually present, we measured vesicle phosphate content and compared it with that of the average cell. Vesicles from MIT9312 and MIT9313 contained 3.4 ± 3.2 and 3.7 ± 2.1 femtomoles phosphate per 10^6 vesicles, respectively. Given that a *Prochlorococcus* MED4 cell (similar in size to MIT9312) contains ~10 attomoles phosphorus (P) (Bertilsson et al., 2003) and *Pelagibacter* HTCC7211 cells contain ~16 attomoles P (White et al., 2019) under P-limited conditions, an encounter with a single *Prochlorococcus* vesicle would supply only ~0.01% of these cells' P quota. These are averages, however, and absent an understanding of the distribution of P per vesicle and vesicle-cell encounter rates it is impossible to speculate further; we simply flag this as an interesting question to pursue. We also note that the presence of nucleic acids in vesicles represents a possible source of organic phosphorous.

As has been observed in gram-negative pathogens (Pérez-Cruz et al., 2015), vesicles from both *Prochlorococcus* strains contained ATP, with more released by MIT9312 than MIT9313 (c.f., $3.6 \pm 1.4 \times 10^{-3}$ and $0.13 \pm 0.097 \times 10^{-3}$ femtomoles ATP per 10^6 vesicles, respectively). This has relevance for interpreting measurements of 'dissolved' ATP in the <0.2 µm fraction of seawater (Azam and Hodson, 1977) – which we now know contains vesicles. This dissolved ATP can be utilized by marine microbes (Azam and Hodson, 1977) and has been proposed to arise largely from cellular excretion during active growth and/or grazing activity, as opposed to lysis of dead cells (Nawrocki and Karl, 1989; Björkman and Karl, 2001). Our results suggest that at least some fraction of this dissolved ATP excreted by marine microbes is likely associated with extracellular vesicles.

Prochlorococcus vesicles contain catalytically active enzymes

Many of the most abundant proteins in the vesicles were putative enzymes (Supporting Information Table S2) which have the potential to mediate exoenzymatic activities. We conducted a set of *in vitro* biochemical assays with intact, purified vesicles and showed that they exhibit lipase, phosphatase, protease, and sulfatase activity (Fig. 3). Vesicle-associated enzymes are thus able to access substrates in the media, likely either via leakage of compounds across the membrane, transport into the vesicle interior, or by acting directly on substrates outside of the vesicle. We also observed that substrate




Fig. 3. Enzymatic activity in purified *Prochlorococcus* vesicles. Activity of (A) lipases, (B) phosphatases, (C) proteases, and (D) sulfatases in samples of vesicles from *Prochlorococcus* MIT9312 (white) and MIT9313 (grey) are compared to that of cells and media background. Values represent the mean (\pm SD) of two to three biological replicates, standardized per population of either 10⁹ vesicles or 10⁶ cells. *nd* = not detected.

accessibility can differ between vesicles and cells: *Prochlorococcus* MIT9313 vesicles displayed measurable sulfatase activity, while intact cells did not (Fig. 3D). Thus, vesicle-associated enzymes have the potential to act upon extracellular substrates that they might not encounter within the cell (Ebner and Götz, 2019), suggesting that many proteins typically thought to be cytosolic could contribute to extracellular biogeochemical processes *in situ*.

We speculate that exporting enzymes in vesicles could be a means of protecting the proteins from environmental damage and degradation, allow for simultaneous co-secretion of multiple proteins, and/or provide an environment where substrates could be maintained in close proximity in the otherwise dilute oligotrophic ocean (Bonnington and Kuehn, 2014). Given that osmolytes can play important roles in maintaining enzyme activity within cells (Yancey, 2005), it is also intriguing that these

vesicle-associated enzyme activities are maintained in the absence of the most abundant detectable cellular osmolytes. While cyanobacteria and many other microbes specifically secrete certain exoenzymes into their local environment (Christie-Oleza *et al.*, 2015), the broad diversity of the vesicle proteome seen here indicates that most cellular enzymes may be exported at some level within vesicles and thus have the potential to function extracellularly.

Can marine extracellular vesicles interact with cells from different organisms?

We know from studies of terrestrial and host-associated microbes that many biological functions of vesicles are attributable to the transfer of vesicle contents between cells (Schwechheimer and Kuehn, 2015; Brown

et al., 2015). Vesicle-mediated biotic ‘interactions’ may encompass a range of mechanisms, from vesicle-associated activities occurring in proximity to a cell (e.g. enzymatic activities or phage defence; MacDonald and Kuehn, 2012; Biller et al., 2014; Rakoff-Nahoum et al., 2014) to the delivery of vesicle contents into another cell (Kadurugamuwa and Beveridge, 1999). To begin to explore interactions between *Prochlorococcus* vesicles and cells, we looked for physical associations between fluorescently labelled vesicles and six different marine microbes from the Proteobacteria, Cyanobacteria, and Bacteroidetes – the three most abundant bacterial phyla in the oceans (Fig. 4). We found that vesicles from *Prochlorococcus* MIT9313 associated with cells of both *Prochlorococcus* MIT9312 and MIT9313, along with representatives of the marine Gammaproteobacteria and Alphaproteobacteria – including *Candidatus Pelagibacter ubique* HTCC7211, a member of the numerically dominant SAR11 group of marine heterotrophs (Figs. 4 and S7). The breadth of interactions was not specific to *Prochlorococcus* vesicles, as vesicles purified from the marine heterotroph *Alteromonas* MIT1002, which was co-isolated with *Prochlorococcus* (Biller et al., 2015b), were also able to interact with these strains (Figs. 4 and S7). To ensure that the observed vesicle-cell associations were not an artefact of the fluorescent dye used, we incubated the same set of marine microbes with vesicles from an *E. coli* strain (Dinh and Bernhardt, 2011) which instead expressed GFP within vesicles, and found that they were also able to interact with multiple microbes (Figs. 4B and S7). While we further verified that our

fluorescent vesicle labelling approach did not significantly influence the ability of *E. coli* vesicles to interact with cells (Fig. S8), we cannot rule out a potential influence of vesicle surface modifications on the pairwise interactions tested.

Four of the strains examined interacted with all of the vesicles tested (Fig. 4B) while two others exhibited apparent specificity toward vesicles from different sources. That is, vesicles from both *Alteromonas* and *Prochlorococcus* MIT9312 did not associate with *Prochlorococcus* MIT9313 cells, and none of the vesicles interacted with *Polaribacter* MED152, a marine *Bacteroidetes* (González et al., 2008), at our detection levels (Fig. 4B). This confirms that our observations are not simply due to experimental conditions or random ‘sticking’ of vesicles to cells. Specificity in vesicle-cell associations has been shown in some heterotrophic bacteria (Tashiro et al., 2017; Toyofuku et al., 2017) and likely underlies variation in vesicle-mediated DNA transfer rates among species (Tran and Boedicker, 2017). Our data are consistent with the hypothesis that different vesicle-cell pairs may vary in the strength of their interactions (Supporting Information Fig. S7). For example, the propensity of a vesicle to associate with a particular cell may depend on its specific surface composition, which would likely be obscured by the averaging effect of our population-scale analysis. The apparent lack of vesicle associations with *Polaribacter* in our experiments is striking and invites an exploration of what factors, such as surface charge (Tashiro et al., 2017), cell envelope structure, and surface hydrophobicity (MacDonald and Beveridge, 2002),




Fig. 4. Association of extracellular vesicles with diverse microbial cells.

A. Example flow cytometry plots of fluorescently labelled *Prochlorococcus* MIT9313 vesicles (left), *Alteromonas* MIT1002 and *Pelagibacter ubique* (SAR11) cell populations (centre), and the same cells following a 2 h incubation with vesicles (right). Internal reference beads are labelled.

B. Interaction of labelled vesicles with cells (+), or lack thereof (nd = not detected), for different ‘source’ strains (vesicle producers) and ‘recipient’ strains (cells exposed to labelled vesicles). Positive interactions were determined based on two criteria: an increase in normalized median cellular 530 nm fluorescence when mixed with labelled vesicles (as in a; see also Fig. S7), and a statistically significant shift in the cellular population distribution (χ^2 T(x) test, $P < 0.01$ in two to three biological replicates; see Supplementary Methods). *Alteromonas* and *Prochlorococcus* vesicles were covalently labelled with an amine-reactive Alexa 488 dye; *E. coli* vesicles contained GFP.

might define interaction boundaries. More generally, these results raise new questions concerning the degree to which vesicles should be considered 'public goods' in the oceans.

The vesicle-cell associations could be disrupted by repeated rounds of centrifugation and washing of the cells (Supporting Information Fig. S9), suggesting that the majority of labelled vesicle material was not fully integrated into the outer membrane of the cells (at least at this timepoint) but was attached via non-covalent bonding between the two surfaces. Although these observations do not directly establish a particular biologically relevant function, we speculate that 'captured' vesicles could serve many functions, including delivering cargo through membrane fusion, 'flipping' of molecules from the vesicle membrane into the cell (Remis *et al.*, 2014), or extracellular degradation and subsequent uptake of vesicle contents. Alternatively, the presence of vesicles nearby could serve defensive roles for cells (Manning and Kuehn, 2011) or influence cellular processes via enzymatic activities.

Potential functional roles for Prochlorococcus vesicles in marine ecosystems

The diversity of extracellular vesicle contents observed here is consistent with the idea that vesicles do not have one dedicated function but instead may play a variety of roles within microbial ecosystems. Our detailed inventory of *Prochlorococcus* extracellular vesicle contents indicates potential functional impacts of vesicles specific to dilute ocean ecosystems that merit future exploration from both conceptual and quantitative perspectives. There are a broad range of compounds within vesicles that could be exchanged between microbes or serve other roles in these particular ecosystems. For instance, the triose and tetrose sugars could serve as an additional energy or carbon source for organisms, act as antioxidants, or possibly serve as the primary osmolyte within vesicles (Goh *et al.*, 2010). While there may be only ~2 molecules of ATP within any individual vesicle on a bulk-average basis, the aggregate impact of vesicle-associated ATP exchange among cells remains to be established; further, the presence of proteins and ATP within the same vesicle could facilitate ATP-requiring extracellular reactions or enable other types of roles such as signalling. Many additional possible ecological functions for vesicles are likely reflected in the other proteins and metabolites found in our inventory of *Prochlorococcus* vesicles, which we have not specifically discussed here.

The identification of diverse oxidized compounds in *Prochlorococcus* vesicles (Table 2 and Fig. 1D) raises additional questions concerning the role vesicles may

play in dissipating oxidative stress – a particular challenge for *Prochlorococcus*, which lacks enzymes typically used for detoxification of hydrogen peroxide such as catalase (Scanlan *et al.*, 2009). Vesicles could be acting as a mechanism for removing damaged compounds, complementing other cellular metabolite degradation and repair pathways (Linster *et al.*, 2013), or as a 'sink' for reactive oxygen species (ROS) generated within the cell, within vesicles, or in the extracellular seawater environment (Morris *et al.*, 2016). Regardless of the oxidative mechanism(s) involved, suppression of ROS toxicity and/or removal of oxidized metabolites may be another way in which vesicles serve a protective role for cells (Manning and Kuehn, 2011). Our identification of vesicle-associated carotenoids, which have also been observed in vesicles from *Synechocystis* PCC6803 (Pardo *et al.*, 2015), raise broader questions as to whether these compounds might serve a photoprotective role for vesicle contents.

Given the number of proteins and small molecules identified in the vesicles, it seems implausible that these materials are uniformly distributed among each member of the vesicle population. We have previously shown heterogeneity of DNA fragment incorporation into vesicles from multiple marine bacteria (Biller *et al.*, 2017), and vesicle subpopulations containing both outer and inner membrane material from Gram-negative cells have been reported (Pérez-Cruz *et al.*, 2015). Furthermore, distinct vesicle subpopulations with unique origins and epitopes have been noted for some time in eukaryotic cells (Kowal *et al.*, 2016). Thus, the functional potential of one vesicle is likely different from the next – similar to our perspective on the cells from which vesicles are derived, whose ecological role can only be understood by viewing them as populations of diverse entities, which may display a diverse range of functions in different contexts (Biller *et al.*, 2015a).

Conclusions

The abundance of vesicles in seawater, and the diversity of their chemical contents, prompt a reassessment of how we conceptualize and interpret the pool of dissolved organic matter that is so critical to the function of marine ecosystems. While many cellular compounds are secreted in the form of truly 'dissolved' individual molecules, it is now clear that microbes also release organic molecules in locally structured, colloidal vesicles which may influence the accessibility, activity, and half-lives of their contents. As such, vesicles blur the boundaries between definitions of 'particulate' and 'dissolved', or 'living' and 'detrital', in marine biogeochemistry – distinctions that are critical for understanding and modelling the system. Extracellular vesicles represent a new frontier for

understanding the flow of energy, materials, and information in ocean ecosystems.

Experimental procedures

Culturing conditions and sampling for lipidomics, proteomics, and metabolomics

To minimize media background signals, axenic cultures of *Prochlorococcus* MIT9312 and MIT9313 used for lipidomics, proteomics, and metabolomics analyses were grown in chemically defined artificial AMP1 media (Moore *et al.*, 2007) supplemented with 10 mM (final concentration) filter-sterilized sodium bicarbonate. All 20 l cultures were grown in polycarbonate carboys (ThermoFisher Nalgene, Waltham, MA, USA) with gentle stirring (60 rpm), under constant light flux (10–20 $\mu\text{mol Q m}^{-2} \text{ s}^{-1}$ for MIT9313; 30–40 $\mu\text{mol Q m}^{-2} \text{ s}^{-1}$ for MIT9312) at 24°C.

All cell and vesicle samples were collected during mid-to-late exponential growth phase. Cell pellets were obtained by gently centrifuging cells (7500 $\times g$) for 10 min at 4°C; vesicle isolation is detailed below. For lipidomics, proteomics, and metabolomics analyses, a total of seven 20 l cultures were grown in AMP1 media for each of the two *Prochlorococcus* strains, providing three replicates for the lipid and small metabolite analysis and an additional sample for proteomics analysis. *Prochlorococcus* AMP1 growth media blanks and vesicle suspension buffer [phosphate buffered saline (PBS)] blanks served as controls for lipidomics, proteomics, and metabolomics analysis; they were extracted and analysed alongside all cell and vesicle samples. Since *Prochlorococcus* was grown under continuous light, the cultures were not synchronized. Thus, all cell samples represent a bulk average population of cells at all stages of the cell cycle; vesicle samples similarly integrate material released across both the lag and exponential growth (steady state) phases.

Vesicle isolation

Vesicles were isolated as described previously (Biller *et al.*, 2014). Briefly, cultures were first gravity filtered through a 0.2 μm capsule filter (Polycap 150TC; GE Life Sciences/Whatman, Maidstone, UK). The filtrate was then concentrated using a 100 kDa tangential flow filter (Ultrasette with Omega membrane; Pall, Port Washington, NY, USA) and re-filtered through a 0.2 μm syringe filter. Vesicles were pelleted from the sample by ultracentrifugation at $\sim 100\,000 \times g$ (Beckman-coulter SW32Ti rotor; 32 000 rpm, 1.5 h, 4°C), purified on an OptiPrep gradient (Biller *et al.*, 2014), then washed and resuspended in 0.2 μm filtered $1 \times$ PBS.

Vesicle concentrations were measured with a NanoSight LM10HS instrument equipped with a LM14 blue laser module using NTA software V3.1 (NanoSight/Malvern, Westborough, MA, USA). Samples were diluted such that the average number of particles per field was between 20 and 60, per the manufacturer's guidelines. Three replicate videos were collected from each sample at a camera level of 11 and analysed at a detection threshold of 1. The sample chamber was thoroughly flushed with 18.2 $\text{M}\Omega \text{ cm}^{-1}$ water (Milli-Q; Millipore, Burlington, MA, USA) between samples and visually examined to ensure that no particles were carried over.

Lipidomics

Lipids were extracted from triplicate cell pellets and vesicles from *Prochlorococcus* MIT9312 or MIT9313 using a modified Bligh and Dyer protocol using a 2:1:0.8 ratio of methanol:dichloromethane:phosphate buffered saline (Popendorf *et al.*, 2013). The total lipid extract was analysed on an Agilent 1200 reversed phase high-performance liquid chromatograph (HPLC) coupled to a ThermoFisher Exactive Plus Orbitrap high-resolution mass spectrometer (HRMS; ThermoFisher, Waltham, MA, USA) which scanned across a range of 100–1500 m/z . HPLC and MS conditions are based on methods described by Collins *et al.*, 2016 (modified after Hummel *et al.*, 2011) and are detailed in the Supporting Information. For the identification and peak area integration, we used LOBSTAHS, an open-source lipidomics software workflow based on adduct ion abundances and several other orthogonal criteria (Collins *et al.*, 2016). Lipid peaks identified using the LOBSTAHS software were integrated from MS data after pre-processing with XCMS (Smith *et al.*, 2006) and CAMERA (Kuhl *et al.*, 2012) and corrected for differences in response factors among different lipids using commercially available standards as described by Becker *et al.* (2018). Additional details are available in the Supplementary Information.

Proteomics

Proteins were extracted from cell biomass and vesicles from a single 20 l batch culture of *Prochlorococcus* MIT9313 or MIT9312. Filters containing cell biomass or vesicle suspensions were extracted using bead-beating and freeze–thaw cycles. Proteins from lysed cells or vesicles were suspended in RapiGest SF (Waters, Milford, MA, USA) to facilitate protein solubilization and then underwent disulfide reduction with tris(2-carboxyethyl)phosphine, alkylation with iodoacetamide, and in-solution protease digestion using either a trypsin/Lys-C mix or Glu-C (Promega, Madison, WI, USA). Following RapiGest hydrolysis and

desalting, samples were resuspended in a solution containing an internal standard of synthetic peptides (Hi3 *Escherichia coli* standard, Waters), spiked with iRT retention time standard (Biognosys, Boston, MA, USA), and immediately analysed on a Waters ACQUITY M-class LC coupled to a Thermo QExactive HF HRMS equipped with a nano-electrospray source. Data-dependent acquisition was performed on the top 10 ions and data analysis were conducted using the software from the trans-proteomic pipeline (TPP v.5.1.0) (Nesvizhskii *et al.*, 2007).

Label-free comparison of relative protein abundances was facilitated by normalizing protein spectral counts to the Hi3 internal standard to account for differences in ionization of peptides due to sample matrix effects and then normalizing to the amount of biomass injected onto the LC-HRMS. To compare protein enrichments within *Prochlorococcus* cell and vesicle fractions, 'per cell' or 'per vesicle' spectral counts were then normalized on the basis of size, or biovolume, differences between a cell and a vesicle, herein referred to as 'biovolume-normalized protein spectral counts' (Supporting Information Tables S2 and S3). Unless otherwise indicated, proteome comparisons across cells and vesicles were conducted using biovolume-normalized spectral counts from only trypsin/Lys-C-based proteomes (Fig. 2). Relative protein enrichments in *Prochlorococcus* vesicles as compared with cells was computed using \log_2 ratios (Supporting Information Table S4) and displayed in Fig. 2A using only the proteins with spectral counts in the top 25% of those identified in vesicles of strain MIT9312 or MIT9313. Subcellular localization assignments are based on a combination of predictions from the Uniprot database (<https://www.uniprot.org>), results from the PSORTb algorithm (V3.0.2) (Yu *et al.*, 2010), amended with data from TMHMM 2.0 (Krogh *et al.*, 2001) and SignalP (V4.1) (Nielsen, 2017), based on the Gram-negative bacteria model. Proteins with a putative signal peptide but no other significant localization predictions are noted as being 'unclassified non-cytoplasmic'. Additional details are available in Supplementary Information.

Metabolomics

Cell pellets and vesicles originating from triplicate 20 l batch cultures of *Prochlorococcus* MIT9312 or MIT9313 were extracted using a modified Bligh and Dyer technique with cold 1:1 methanol/water (aqueous phase) and cold dichloromethane (organic phase) (Bligh and Dyer, 1959; Boysen *et al.*, 2018). Cell pellets were manually disrupted by bead beating during the extraction, as described by Boysen *et al.* (2018); vesicles were extracted without bead beating. Metabolite separations were achieved using reversed-phase (RP, for aqueous and organic extracts) and hydrophilic interaction liquid

chromatography (HILIC, for aqueous extracts only), as detailed in Table S8 and Boysen *et al.*, 2018. Data were collected in positive ion mode for RP analyses with a scan range of 90–900 *m/z*. For HILIC analyses, data were obtained in positive and negative mode (using polarity switching) with a scan range of 80–900 *m/z*. For every sample, data were processed in four subgroups, defined by phase, chromatography, and ion mode: RP-organic-positive (RPOrgPos), RP-aqueous-positive (RPAqPos), HILIC-aqueous-positive (HILICAqPos), and HILIC-aqueous-negative (HILICAqNeg). Data were collected using Thermo QExactive HF HRMS using same settings as in the study by Boysen *et al.* (2018).

For targeted data, individual metabolite features were integrated using Skyline Daily (MacLean *et al.*, 2010) and subjected to an in-house quality control protocol (Boysen *et al.*, 2018). Untargeted data were processed using MS-DIAL software (Tsugawa *et al.*, 2015) using parameters reported in Table S9. MS-DIAL was used to pick, align, and integrate mass features from raw datasets. Identification of mass features within untargeted data sets was accomplished via dereplication of the mass feature list against several databases, yielding identifications of variable confidence as described in the study by Heal *et al.* (2019). We ranked confidence in mass feature identifications according to existing literature (Sumner *et al.*, 2007). All mass features were searched against an Ingalls Lab in-house database of authenticated standards, LOBSTAHs output, and MassBank (Horai *et al.*, 2010). Additional details are available in Supplementary Information.

Vesicle enzyme activity assays

All nutrient and enzymatic assays were carried out using vesicles collected from 20 l exponentially growing *Prochlorococcus* cultures grown as above. Vesicles were purified using an OptiPrep (Iodixanol) density gradient as described in the study by Biller *et al.* (2014) and washed in clean 1 × PBS. ATP was measured using a standard luminescence-based assay with the BacTiter-Glo kit (Promega) according to the manufacturer's directions. Vesicle phosphate was measured using the Sigma-Aldrich Phosphate Assay kit (MilliporeSigma, St. Louis, MO, USA). Protease measurements were made using the Sigma-Aldrich Protease Fluorescent Detection Kit following the manufacturer's instructions with the following modifications: 10 μ l of a 2 × incubation buffer was used in each 50 μ l reaction, which allowed us to add 20 μ l of sample to each reaction and increase the number of vesicles included in the reaction. The reaction was carried out for 18–20 h at 28 °C in the dark. Sulfatase activity was measured using the BioVision Sulfatase Activity Assay (BioVision, Milpitas, CA, USA), following manufacturer's instructions.

Phosphatase and lipase activity were measured by following the hydrolysis of fluorescent substrates 4-methylumbelliferyl phosphate and 4-methylumbelliferyl oleate (both from Sigma-Aldrich) following standard methods (Hoppe, 1993). Briefly, $\sim 10^{10}$ vesicles in $1 \times$ PBS were incubated with $100 \mu\text{M}$ substrate (dissolved in ethylene glycol monomethyl ether) and natural seawater from the Sargasso Sea in a $200 \mu\text{l}$ reaction within a black-walled 96-well plate at room temperature for 16 h. Boiled vesicle samples and substrate in seawater lacking any added vesicles were used as controls.

Vesicle interaction experiments

All cultures used for the interaction experiments were grown in media based on $0.2 \mu\text{m}$ filtered natural water from Vineyard Sound, MA except where noted. Vesicles were purified from 20 l cultures of *Prochlorococcus* MIT9312 or MIT9313 grown as above but in Pro99 media (Moore et al., 2007), and 10 l cultures of *Alteromonas* strain MIT1002 (Biller et al., 2015b) were grown at 24°C in ProMM media (Berube et al., 2015) (Pro99 media, as above, plus lactate, pyruvate, glycerol, acetate, and $1 \times$ Va vitamin mix; Waterbury and Willey, 1988). The 10 l cultures of *E. coli* strain TB28(*attHKT*B263) (Dinh and Bernhardt, 2011; this strain expresses the Superfolder variant of GFP and targets it to the periplasm) were grown at 30°C in M9 maltose media supplemented with $250 \mu\text{M}$ isopropyl β -D-1-thiogalactopyranoside and $100 \mu\text{g m}^{-1}$ ampicillin. Purified vesicles were covalently labelled with an amine-reactive Alexa Fluor 488 5-SDP ester dye (Molecular Probes/ThermoScientific).

To examine the ability of vesicles to associate with representative phylogenetically distinct marine microbes, cells of *Prochlorococcus* MIT9312 and MIT9313 (in Pro99 media), *Alteromonas* MIT1002 and *Thalassospira* MIT1004 (Biller et al., 2017) (in ProMM media), and *Polaribacter* MED152 (González et al., 2008) (in Pro99 media, supplemented with 5 g l peptone and 1 g l yeast extract) were all grown to mid-exponential growth phase at 24°C . *Pelagibacter* HTCC7211 was grown in defined AMS1 media (Carini et al., 2013) at 22°C . Approximately 10^9 – 10^{10} vesicles (labelled or unlabelled), or an equivalent volume of the PBS Alexa 488 labelling control, were added to 2 ml of culture at $\sim 10^5$ cells ml^{-1} (final ratio: $\sim 1000:1$ vesicles:cells). Cultures with vesicles were incubated for 2 h at the normal growth temperatures and examined on an Influx flow cytometer (Cytocopia/BD, Franklin Lakes, NJ). Cells were excited using a blue 488 nm laser and monitored for chlorophyll (692/40 nm emission) and Alexa 488/GFP (530/40 nm emission) fluorescence. All flow cytometry data were analysed using FlowJo (V10.5). Additional details are available in Supplementary Information.

Miscellaneous

All data were analysed in R (V3.5.1). Plots were created using ggplot2 (Wickham, 2009).

Data availability

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE (Perez-Riverol et al., 2019) partner repository with the dataset identifier PXD013602. Metabolite data are available at the NIH Common Fund's National Metabolomics Data Repository (NMDR) website, the Metabolomics Workbench, <https://www.metabolomicsworkbench.org> (Sud et al., 2016), under Study ID # ST001524.

Acknowledgements

This work was funded by grants from the National Science Foundation (OCE-1356460 to S.W.C.) and the Simons Foundation (SCOPE Award ID 329108 to B.A.S.V.M., A.E.I., S.W.C.; Life Sciences Project Award ID 337262, S.W.C.; Simons Award ID 385428 to A.E.I. and 598819 to K.R.H.). K.W.B was supported by the Postdoctoral Scholarship Programme at the Woods Hole Oceanographic Institution. R.A.L was partially supported by a postdoctoral fellowship from the Swiss National Science Foundation. The authors thank Thomas Bernhardt (Harvard Medical School) for providing the sfGFP-expressing *E. coli* strain, Jarone Pinhassi (Linnaeus U.) for sharing *Polaribacter* MED152, and Stephen Giovannoni (U. Oregon) for *Pelagibacter* HTCC7211. The authors also thank Daniel Sher (U. Haifa) for providing helpful suggestions on the manuscript.

References

- Azam, F., and Hodson, R.E. (1977) Dissolved ATP in the sea and its utilisation by marine bacteria. *Nature* **267**: 696–698.
- Becker, J.W., Hogle, S.L., Rosendo, K., and Chisholm, S.W. (2019) Co-culture and biogeography of *Prochlorococcus* and SAR11. *ISME J* **13**: 1506–1519.
- Becker, K.W., Collins, J.R., Durham, B.P., Groussman, R.D., White, A.E., Fredricks, H.F., et al. (2018) Daily changes in phytoplankton lipidomes reveal mechanisms of energy storage in the open ocean. *Nat Commun* **9**: 5179.
- Bertilsson, S., Berglund, O., Karl, D., and Chisholm, S. (2003) Elemental composition of marine *Prochlorococcus* and *Synechococcus*: implications for the ecological stoichiometry of the sea. *Limnol Oceanogr* **48**: 1721–1731.
- Bertilsson, S., Berglund, O., Pullin, M., and Chisholm, S. (2005) Release of dissolved organic matter by *Prochlorococcus*. *Vie et Milieu* **55**: 225–232.
- Berube, P.M., Biller, S.J., Kent, A.G., Berta-Thompson, J.W., Roggensack, S.E., Roache-Johnson, K.H., et al. (2015) Physiology and evolution of nitrate acquisition in *Prochlorococcus*. *ISME J* **9**: 1195–1207.

- Billar, S.J., Berube, P.M., Lindell, D., and Chisholm, S.W. (2015a) *Prochlorococcus*: the structure and function of collective diversity. *Nat Rev Microbiol* **13**: 13–27.
- Billar, S.J., Coe, A., Martin-Cuadrado, A.-B., and Chisholm, S.W. (2015b) Draft genome sequence of *Alteromonas macleodii* strain MIT1002, isolated from an enrichment culture of the marine cyanobacterium *Prochlorococcus*. *Genome Announc* **3**: e00967–e00915.
- Billar, S.J., McDaniel, L.D., Breitbart, M., Rogers, E., Paul, J. H., and Chisholm, S.W. (2017) Membrane vesicles in sea water: heterogeneous DNA content and implications for viral abundance estimates. *ISME J* **11**: 394–404.
- Billar, S.J., Schubotz, F., Roggensack, S.E., Thompson, A. W., Summons, R.E., and Chisholm, S.W. (2014) Bacterial vesicles in marine ecosystems. *Science* **343**: 183–186.
- Bitto, N.J., Cheng, L., Johnston, E.L., Pathirana, R., Phan, T. K., Poon, I.K.H., et al. (2021) *Staphylococcus aureus* membrane vesicles contain immunostimulatory DNA, RNA and peptidoglycan that activate innate immune receptors and induce autophagy. *J Extracell Vesicles* **10**: e12080.
- Björkman, K.M., and Karl, D.M. (2001) A novel method for the measurement of dissolved adenosine and guanosine triphosphate in aquatic habitats: applications to marine microbial ecology. *J Microbiol Methods* **47**: 159–167.
- Bligh, E.G., and Dyer, W.J. (1959) A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* **37**: 911–917.
- Bonnington, K.E., and Kuehn, M.J. (2014) Protein selection and export via outer membrane vesicles. *Biochim Biophys Acta* **1843**: 1612–1619.
- Boysen, A.K., Heal, K.R., Carlson, L.T., and Ingalls, A.E. (2018) Best-matched internal standard normalization in liquid chromatography-mass spectrometry metabolomics applied to environmental samples. *Anal Chem* **90**: 1363–1369.
- Brown, L., Wolf, J.M., Prados-Rosales, R., and Casadevall, A. (2015) Through the wall: extracellular vesicles in Gram-positive bacteria, mycobacteria and fungi. *Nat Rev Microbiol* **13**: 620–630.
- Carini, P., Steindler, L., Beszteri, S., and Giovannoni, S.J. (2013) Nutrient requirements for growth of the extreme oligotroph “*Candidatus Pelagibacter ubique*” HTCC1062 on a defined medium. *ISME J* **7**: 592–602.
- Christie-Oleza, J.A., Armengaud, J., Guerin, P., and Scanlan, D.J. (2015) Functional distinctness in the exoproteomes of marine *Synechococcus*. *Environ Microbiol* **17**: 3781–3794.
- Collins, J.R., Edwards, B.R., Fredricks, H.F., and Van Mooy, B.A.S. (2016) LOBSTAHS: an adduct-based lipidomics strategy for discovery and identification of oxidative stress biomarkers. *Anal Chem* **88**: 7154–7162.
- Cubillos-Ruiz, A., Berta-Thompson, J.W., Becker, J.W., van der Donk, W.A., and Chisholm, S.W. (2017) Evolutionary radiation of lanthipeptides in marine cyanobacteria. *Proc Natl Acad Sci* **114**: E5424–E5433.
- Deatherage, B.L., and Cookson, B.T. (2012) Membrane vesicle release in Bacteria, Eukaryotes, and Archaea: a conserved yet underappreciated aspect of microbial life. *Infect Immun* **80**: 1948–1957.
- Dinh, T., and Bernhardt, T.G. (2011) Using Superfolder green fluorescent protein for periplasmic protein localization studies. *J Bacteriol* **193**: 4984–4987.
- Ebner, P., and Götz, F. (2019) Bacterial excretion of cytoplasmic proteins (ECP): occurrence, mechanism, and function. *Trends Microbiol* **27**: 176–187.
- Flombaum, P., Gallegos, J.L., Gordillo, R.A., Rincón, J., Zabala, L.L., Jiao, N., et al. (2013) Present and future global distributions of the marine cyanobacteria *Prochlorococcus* and *Synechococcus*. *Proc Natl Acad Sci* **110**: 9824–9829.
- Goh, F., Barrow, K.D., Burns, B.P., and Neilan, B.A. (2010) Identification and regulation of novel compatible solutes from hypersaline stromatolite-associated cyanobacteria. *Arch Microbiol* **192**: 1031–1038.
- González, J.M., Fernandez-Gomez, B., Fernández-Guerra, A., Gómez-Consarnau, L., Sánchez, O., Coll-Lladó, M., et al. (2008) Genome analysis of the proteorhodopsin-containing marine bacterium *Polaribacter* sp. MED152 (Flavobacteria). *Proc Natl Acad Sci* **105**: 8724–8729.
- Guerrero-Mandujano, A., Hernández-Cortez, C., Ibarra, J.A., and Castro-Escarpulli, G. (2017) The outer membrane vesicles: secretion system type zero. *Traffic* **18**: 425–432.
- Heal, K.R., Kellogg, N.A., Carlson, L.T., Lionheart, R.M., and Ingalls, A.E. (2019) Metabolic consequences of cobalamin scarcity in the diatom *Thalassiosira pseudonana* as revealed through metabolomics. *Protist* **170**: 328–348.
- Hoppe, H.-G. (1993) . In *Use of Fluorogenic Model Substrates for Extracellular Enzyme Activity (EEA) Measurement of Bacteria*. In *Handbook of Methods in Aquatic Microbial Ecology*, Kemp, P.F., Sherr, B.F., Sherr, E.B., and Cole, J.J. (eds). Boca Raton, FL: Lewis Publishers, pp. 423–431.
- Horai, H., Arita, M., Kanaya, S., Nihei, Y., Ikeda, T., Suwa, K., et al. (2010) MassBank: a public repository for sharing mass spectral data for life sciences. *J Mass Spectrom* **45**: 703–714.
- Hummel, J., Segu, S., Li, Y., Irgang, S., Jueppner, J., and Giavalisco, P. (2011) Ultra performance liquid chromatography and high resolution mass spectrometry for the analysis of plant lipids. *Front Plant Sci* **2**: 54.
- Jørgensen, N.O., and Jacobsen, C.S. (1996) Bacterial uptake and utilization of dissolved DNA. *Aquat Microb Ecol* **11**: 263–270.
- Jürgens, U.J., and Weckesser, J. (1985) Carotenoid-containing outer membrane of *Synechocystis* sp. strain PCC6714. *J Bacteriol* **164**: 384–389.
- Kadurugamuwa, J.L., and Beveridge, T.J. (1996) Bacteriolytic effect of membrane vesicles from *Pseudomonas aeruginosa* on other bacteria including pathogens: conceptually new antibiotics. *J Bacteriol* **178**: 2767–2774.
- Kadurugamuwa, J.L., and Beveridge, T.J. (1999) Membrane vesicles derived from *Pseudomonas aeruginosa* and *Shigella flexneri* can be integrated into the surfaces of other gram-negative bacteria. *Microbiology* **145**: 2051–2060.
- Klähn, S., and Hagemann, M. (2010) Compatible solute biosynthesis in cyanobacteria. *Environ Microbiol* **13**: 551–562.
- Kowal, J., Arras, G., Colombo, M., Jouve, M., Morath, J.P., Primdal-Bengtson, B., et al. (2016) Proteomic comparison defines novel markers to characterize heterogeneous populations of extracellular vesicle subtypes. *Proc Natl Acad Sci* **113**: E968–E977.

- Krogh, A., Larsson, B., von Heijne, G., and Sonnhammer, E. L. (2001) Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *J Mol Biol* **305**: 567–580.
- Kuhl, C., Tautenhahn, R., Böttcher, C., Larson, T.R., and Neumann, S. (2012) CAMERA: an integrated strategy for compound spectra extraction and annotation of liquid chromatography/mass spectrometry data sets. *Anal Chem* **84**: 283–289.
- Li, B., Sher, D., Kelly, L., Shi, Y., Huang, K., Knerr, P.J., et al. (2010) Catalytic promiscuity in the biosynthesis of cyclic peptide secondary metabolites in planktonic marine cyanobacteria. *Proc Natl Acad Sci U S A* **107**: 10430–10435.
- Linster, C.L., Van Schaftingen, E., and Hanson, A.D. (2013) Metabolite damage and its repair or pre-emption. *Nat Chem Biol* **9**: 72–80.
- Lynch, J.B., and Alegado, R.A. (2017) Spheres of hope, packets of doom: the good and bad of outer membrane vesicles in interspecies and ecological dynamics. *J Bacteriol* **199**: e00012–e00017.
- MacDonald, I.A., and Kuehn, M.J. (2012) Offense and defense: microbial membrane vesicles play both ways. *Res Microbiol* **163**: 607–618.
- MacDonald, I.A., and Kuehn, M.J. (2013) Stress-induced outer membrane vesicle production by *Pseudomonas aeruginosa*. *J Bacteriol* **195**: 2971–2981.
- MacDonald, K.L., and Beveridge, T.J. (2002) Bactericidal effect of gentamicin-induced membrane vesicles derived from *Pseudomonas aeruginosa* PAO1 on gram-positive bacteria. *Can J Microbiol* **48**: 810–820.
- MacLean, B., Tomazela, D.M., Shulman, N., Chambers, M., Finney, G.L., Frewen, B., et al. (2010) Skyline: an open source document editor for creating and analyzing targeted proteomics experiments. *Bioinformatics* **26**: 966–968.
- Manning, A.J., and Kuehn, M.J. (2011) Contribution of bacterial outer membrane vesicles to innate bacterial defense. *BMC Microbiol* **11**: 258.
- Mashburn, L.M., and Whiteley, M. (2005) Membrane vesicles traffic signals and facilitate group activities in a prokaryote. *Nature* **437**: 422–425.
- Mashburn-Warren, L.M., and Whiteley, M. (2006) Special delivery: vesicle trafficking in prokaryotes. *Mol Microbiol* **61**: 839–846.
- Moore, L., Coe, A., Zinser, E.R., Saito, M., Sullivan, M., Lindell, D., et al. (2007) Culturing the marine cyanobacterium *Prochlorococcus*. *Limnol Oceanogr: Methods* **5**: 353–362.
- Morris, J.J., Johnson, Z.I., Wilhelm, S.W., and Zinser, E.R. (2016) Diel regulation of hydrogen peroxide defenses by open ocean microbial communities. *J Plankton Res* **38**: 1103–1114.
- Nawrocki, M.P., and Karl, D. (1989) Dissolved ATP turnover in the Bransfield Strait, Antarctica during a spring bloom. *Mar Ecol Prog Ser* **57**: 35–44.
- Nesvizhskii, A.I., Vitek, O., and Aebersold, R. (2007) Analysis and validation of proteomic data generated by tandem mass spectrometry. *Nat Methods* **4**: 787–797.
- Nielsen, H. (2017) Predicting secretory proteins with SignalP. *Methods Mol Biol* **1611**: 59–73.
- Orench-Rivera, N., and Kuehn, M.J. (2016) Environmentally controlled bacterial vesicle-mediated export. *Cell Microbiol* **18**: 1525–1536.
- Ottesen, E.A., Young, C.R., Gifford, S.M., Eppley, J.M., Marin, R., Schuster, S.C., et al. (2014) Multispecies diel transcriptional oscillations in open ocean heterotrophic bacterial assemblages. *Science* **345**: 207–212.
- Pardo, Y.A., Florez, C., Baker, K.M., Schertzer, J.W., and Mahler, G.J. (2015) Detection of outer membrane vesicles in *Synechocystis* PCC 6803. *FEMS Microbiol Lett* **362**: fnv163.
- Pérez-Cruz, C., Delgado, L., López-Iglesias, C., and Mercade, E. (2015) Outer-inner membrane vesicles naturally secreted by Gram-negative pathogenic bacteria. *PLoS ONE* **10**: e0116896.
- Perez-Riverol, Y., Csordas, A., Bai, J., Bernal-Llinares, M., Hewapathirana, S., Kundu, D.J., et al. (2019) The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic Acids Res* **47**: D442–D450.
- Popendorf, K.J., Fredricks, H.F., and Van Mooy, B.A.S. (2013) Molecular ion-independent quantification of polar glycerolipid classes in marine plankton using triple quadrupole MS. *Lipids* **48**: 185–195.
- Rakoff-Nahoum, S., Coyne, M.J., and Comstock, L.E. (2014) An ecological network of polysaccharide utilization among human intestinal symbionts. *Curr Biol* **24**: 40–49.
- Remis, J.P., Wei, D., Gorur, A., Zemla, M., Haraga, J., Allen, S., et al. (2014) Bacterial social networks: structure and composition of *Myxococcus xanthus* outer membrane vesicle chains. *Environ Microbiol* **16**: 598–610.
- Rontani, J.-F., and Belt, S.T. (2020) Photo- and autoxidation of unsaturated algal lipids in the marine environment: an overview of processes, their potential tracers, and limitations. *Org Geochem* **139**: 103941.
- Scanlan, D.J., Ostrowski, M., Mazard, S., Dufresne, A., Garczarek, L., Hess, W.R., et al. (2009) Ecological genomics of marine picocyanobacteria. *Microbiol Mol Biol Rev* **73**: 249–299.
- Schatz, D., Rosenwasser, S., Malitsky, S., Wolf, S.G., Feldmesser, E., and Vardi, A. (2017) Communication via extracellular vesicles enhances viral infection of a cosmopolitan alga. *Nat Microbiol* **2**: 1485–1492.
- Schertzer, J.W., Boulette, M.L., and Whiteley, M. (2009) More than a signal: non-signaling properties of quorum sensing molecules. *Trends Microbiol* **17**: 189–195.
- Schmidt, W., Drews, G., Weckesser, J., Fromme, I., and Borowiak, D. (1980) Characterization of the lipopolysaccharides from eight strains of the cyanobacterium *Synechococcus*. *Arch Microbiol* **127**: 209–215.
- Schwechheimer, C., and Kuehn, M.J. (2015) Outer-membrane vesicles from Gram-negative bacteria: biogenesis and functions. *Nat Rev Microbiol* **13**: 605–619.
- Smith, C.A., Want, E.J., O'Maille, G., Abagyan, R., and Siuzdak, G. (2006) XCMS: processing mass spectrometry data for metabolite profiling using nonlinear peak alignment, matching, and identification. *Anal Chem* **78**: 779–787.
- Sud, M., Fahy, E., Cotter, D., Azam, K., Vadivelu, I., Burant, C., et al. (2016) Metabolomics workbench: an international repository for metabolomics data and

- metadata, metabolite standards, protocols, tutorials and training, and analysis tools. *Nucleic Acids Res* **44**: D463–D470.
- Sumner, L.W., Amberg, A., Barrett, D., Beale, M.H., Beger, R., Daykin, C.A., *et al.* (2007) Proposed minimum reporting standards for chemical analysis Chemical Analysis Working Group (CAWG) metabolomics standards initiative (MSI). *Metabolomics* **3**: 211–221.
- Tandberg, J.I., Lagos, L.X., Langlete, P., Berger, E., Rishovd, A.-L., Roos, N., *et al.* (2016) Comparative analysis of membrane vesicles from three *Piscirickettsia salmonis* isolates reveals differences in vesicle characteristics. *PLoS ONE* **11**: e0165099–e0165027.
- Tartaglia, N.R., Nicolas, A., Ossemond, J., Nguyen, M.T., FLR, d.C., Gilot, D., *et al.* (2020) Extracellular vesicles produced by human and animal *Staphylococcus aureus* strains share a highly conserved core proteome. *Sci Rep* **10**: 1–13.
- Tashiro, Y., Hasegawa, Y., Shintani, M., Takaki, K., Ohkuma, M., Kimbara, K., and Futamata, H. (2017) Interaction of bacterial membrane vesicles with specific species and their potential for delivery to target cells. *Front Microbiol* **8**: 813–873.
- Ting, C.S., Hsieh, C., Sundararaman, S., Mannella, C., and Marko, M. (2007) Cryo-electron tomography reveals the comparative three-dimensional architecture of *Prochlorococcus*, a globally important marine Cyanobacterium. *J Bacteriol* **189**: 4485–4493.
- Toyofuku, M., Morinaga, K., Hashimoto, Y., Uhl, J., Shimamura, H., Inaba, H., *et al.* (2017) Membrane vesicle-mediated bacterial communication. *ISME J* **11**: 1504–1509.
- Tran, F., and Boedicker, J.Q. (2017) Genetic cargo and bacterial species set the rate of vesicle-mediated horizontal gene transfer. *Sci Rep* **7**: 8813.
- Tsugawa, H., Cajka, T., Kind, T., Ma, Y., Higgins, B., Ikeda, K., *et al.* (2015) MS-DIAL: data-independent MS/MS deconvolution for comprehensive metabolome analysis. *Nat Methods* **12**: 523–526.
- Turner, L., Bitto, N.J., Steer, D.L., Lo, C., D'Costa, K., Ramm, G., *et al.* (2018) *Helicobacter pylori* outer membrane vesicle size determines their mechanisms of host cell entry and protein content. *Front Immunol* **9**: 1466.
- Van Mooy, B.A.S., Rocap, G., Fredricks, H.F., Evans, C.T., and Devol, A.H. (2006) Sulfolipids dramatically decrease phosphorus demand by picocyanobacteria in oligotrophic marine environments. *Proc Natl Acad Sci* **103**: 8607–8612.
- Volkman, J.K., Barrett, S.M., Blackburn, S.I., Mansour, M.P., Sikes, E.L., and Gelin, F. (1998) Microalgal biomarkers: a review of recent research developments. *Org Geochem* **29**: 1163–1179.
- Waterbury, J.B., and Willey, J.M. (1988) Isolation and growth of marine planktonic cyanobacteria. *Methods Enzymol* **167**: 100–105.
- White, A.E., Giovannoni, S.J., Zhao, Y., Vergin, K., and Carlson, C.A. (2019) Elemental content and stoichiometry of SAR11 chemoheterotrophic marine bacteria. *Limnol Oceanogr* **4**: 44–51.
- Wickham, H. (2009) *ggplot2: Elegant Graphics for Data Analysis*. New York, NY: Springer.
- Yancey, P.H. (2005) Organic osmolytes as compatible, metabolic and counteracting cytoprotectants in high osmolarity and other stresses. *J Exp Biol* **208**: 2819–2830.
- Yaron, S., Kolling, G.L., Simon, L., and Matthews, K.R. (2000) Vesicle-mediated transfer of virulence genes from *Escherichia coli* O157:H7 to other enteric bacteria. *Appl Environ Microbiol* **66**: 4414–4420.
- Yu, N.Y., Wagner, J.R., Laird, M.R., Melli, G., Rey, S., Lo, R., *et al.* (2010) PSORTb 3.0: improved protein subcellular localization prediction with refined localization subcategories and predictive capabilities for all prokaryotes. *Bioinformatics* **26**: 1608–1615.
- Yun, S.H., Lee, S.-Y., Choi, C.W., Lee, H., Ro, H.-J., Jun, S., *et al.* (2017) Proteomic characterization of the outer membrane vesicle of the halophilic marine bacterium *Novosphingobium pentaromativorans* US6-1. *J Microbiol* **55**: 56–62.
- Zakharzhevskaya, N.B., Vanyushkina, A.A., Altukhov, I.A., Shavarda, A.L., Butenko, I.O., Rakitina, D.V., *et al.* (2017) Outer membrane vesicles secreted by pathogenic and nonpathogenic *Bacteroides fragilis* represent different metabolic activities. *Sci Rep* **7**: 16–65.
- Zavan, L., Bitto, N.J., Johnston, E.L., Greening, D.W., and Liaskos, M.K. (2019) *Helicobacter pylori* growth stage determines the size, protein composition, and preferential cargo packaging of outer membrane vesicles. *Proteomics* **19**: e1800209–e1800223.
- Zwarycz, A.S., Livingstone, P.G., and Whitworth, D.E. (2020) Within-species variation in OMV cargo proteins: the *Myxococcus xanthus* OMV pan-proteome. *Mol Omics* **16**: 387–397.

Supporting Information

Additional Supporting Information may be found in the online version of this article at the publisher's web-site:

Appendix S1: Supporting Information

Supplementary Table S1: Complete dataset of membrane lipids, pigments, and plastoquinone found within *Prochlorococcus* MIT9312 and MIT9313 cells and vesicles.

Supplementary Table S2: Complete proteomic data for *Prochlorococcus* MIT9312 and MIT9313 cells and vesicles, including protein and peptide identification and raw spectral counts.

Supplementary Table S3: Relative protein abundances, or biovolume-normalized spectral counts, for *Prochlorococcus* MIT9312 and MIT9313 cells and vesicles.

Supplementary Table S4: Relative enrichment of proteins in *Prochlorococcus* vesicles vs. cells.

Supplementary Table S5: Metabolite diversity in *Prochlorococcus* vesicle and cell samples.

Supplementary Table S6: Full biovolume normalized peak areas for *Prochlorococcus* vesicles and cells.

Supplementary Table S7: Targeted metabolites detected in vesicles and cells from *Prochlorococcus* MIT9312 and MIT9313.

Supplementary Table S8: Experimental conditions used to analyse metabolite fractions.

Supplementary Table S9: Parameters for MS-DIAL analysis for each analytical fraction.