

Formulation of a Target Plastic Model to estimate critical plastic burdens of toxicants.

Deedar Nabi^{1,2}, Aaron J. Beck¹, Eric P. Achterberg¹

¹ GEOMAR Helmholtz Centre for Ocean Research Kiel

Wischhofstr. 1-3, 24148 Kiel, Germany

²Institute of Environmental Science and Engineering (IESE), School of Civil and Environmental Engineering (SCEE), National University of Sciences and Technology (NUST), H-12,


Islamabad, Pakistan.

Abstract

Plastic pollution has become a widespread problem affecting multiple environmental compartments, with associated chemicals having harmful effects on living organisms. Here, we developed a Target Plastic Model (TPM) to estimate the critical plastic burden of various toxicants in five types of plastics, namely polydimethylsiloxane (PDMS), polyoxymethylene (POM), polyacrylate (PA), low-density polyethylene (LDPE), and polyurethane ester (PU), following the Target Lipid Model (TLM) framework. The critical plastic burdens of baseline (n=115), less-inert (n=73), and reactive (n=75) toxicants ranged from 0.17-51.33, 0.04-26.62, and 1.00×10^{-6} - 6.78×10^{-4} mmol/kg of plastic, respectively. While critical plastic burdens were also estimated for other plastic phases, such as polypropylene (PP), polystyrene (PS), polyvinyl chloride (PVC), ultra-high molecular weight polyethylene (UHMWPE), and high-density polyethylene (HDPE), the findings were less reliable due to a lack of experimental data. Our study showed that PDMS, PA, POM, PE, and PU are similar to biomembranes in mimicking the exchange of chemicals with the water phase. Using the TPM, median lethal concentration (LC₅₀) values for fish exposed to baseline toxicants were predicted, and the results agreed with experimental values, with RMSE ranging from 0.311-0.538 log unit. For less inert chemicals, predictions were within a factor of 5 of experimental values. The TPM's performance was comparable to other widely used models, such as the TLM, ECOSAR, and Abraham Solvation Model. However, like other models, TPM was not effective in predicting the toxicities of reactive toxicants, with RMSE exceeding 1 log unit. TPM can provide valuable insights into the toxicities of chemicals associated with environmental plastic phases, assisting in selecting the best polymeric phase for passive sampling and designing better passive dosing techniques for toxicity experiments. Moreover, TPM can assist in selecting the best plastic phase for developing animal alternatives for toxicity measurement and determining the toxicity of complex mixtures such as those arising during oil spills.

Graphical abstract

FROM TARGET LIPID MODEL TO TARGET PLASTIC MODEL



Introduction

Plastic has undoubtedly brought numerous benefits to modern society. Nevertheless, the plastic revolution has also inflicted significant impacts on global ecosystems. Plastic pollution has become ubiquitous, infiltrating our air,^{1,2} water,³ soil,⁴ biota,⁵ and even our food,⁶ as well as human blood⁷ and fetal fluids.⁸ Disentangling the complexity of plastic pollution is a daunting task, but at its simplest, environmental issues associated from plastic arise from three key factors: the particulate nature of plastic itself,⁹ which includes macro-, micro-, and nanoplastics; the harmful microorganisms that can harbor on plastic debris,^{10,11} and the chemicals associated with plastic.¹²

Chemicals associated with plastic can be categorized as native chemicals, which are the additives added during manufacturing,¹³ and non-native chemicals that plastic materials accumulate once released into the environment.¹⁴ As a result, plastic plays a crucial role in the transportation of both native and non-native chemicals in the environment. These chemicals can potentially leach from the plastic phase into various environmental waters, such as freshwater, marine water, and soil pore water.¹³⁻¹⁵

The leaching of chemicals from plastic is influenced by several factors including the properties of the chemicals, the properties of the plastic materials, water chemistry, and fluid dynamics. The properties of chemicals that affect the uptake and release of chemicals from plastic materials are intermolecular interactions such as polarity, polarizability, hydrogen bonding, and London dispersion interactions.¹⁶ The properties of plastic materials include interaction terms that correspond to the intermolecular interactions of molecules mentioned above.¹⁶ Furthermore, factors such as crystallinity and the

weathering age of plastic materials also play a role in the uptake and release of chemicals, both at equilibrium and under kinetic conditions.¹⁷

Chemical exchange between water and the rubbery (amorphous) fraction of plastic is primarily governed by partitioning (absorption) processes, whereas the crystalline fraction of plastic generally follows adsorption processes.¹⁸ The impact of weathering age on the exchange of chemicals between water and plastic materials has not been extensively studied.¹⁹ Water chemistry, such as pH, dissolved organic matter concentration, and salinity, can influence the exchange of chemicals between plastics and water, particularly under kinetic conditions.¹⁹ Finally, flow regimes, including turbulent, laminar, and stagnant conditions, can affect the thickness of diffusion boundary layer between the water phase and plastic phase, which, in turn, has an impact on the kinetic exchange of chemicals between the two phases. For instance, under turbulent conditions, the thickness of the aqueous boundary layer is thinner compared to stagnant conditions, resulting in faster kinetics.²⁰

The amorphous or rubbery portion of plastic phases mimics organisms in terms of the passive uptake of chemicals.²¹ These plastic phases only take up the freely dissolved fraction, which also defines the bioavailable fraction, of the total concentration of chemicals.²² The total concentration, also referred to as nominal concentration, consists of the freely dissolved fraction as well as fractions that are associated with particulate matter and dissolved organic matter. However, it is the freely dissolved fraction of the chemical concentration that truly defines its exposure to organisms.²³ Plastic phases can help to isolate and quantify this fraction, which can be a valuable tool for studying the behavior of chemicals in aquatic environments and their potential impacts on organisms.

As a result, organic environmental chemists have begun to use plastic phases as passive sampling²⁴ and dosing devices.²⁵ Polymeric phases such as polydimethylsiloxane (PDMS), polyoxymethylene (POM), polyacrylate (PA), ethylene–vinyl acetate, high-density polyethylene (HDPE), and low-density polyethylene (LDPE) are used to monitor the truly dissolved concentration of organic pollutants in air, water,²⁴ sediments, biotic media, and humans.²⁶

Furthermore, scientists are increasingly favoring passive dosing methods to determine exposure,²⁷ toxicity,²⁸ bioconcentration,²⁹ and speciation and fractionation³⁰ of hydrophobic organic compounds in complex systems. These methods utilize the same polymeric phases as passive sampling and provide precise control over exposure concentrations of hydrophobic chemicals in laboratory experiments involving multiple phases.²⁵ In addition to providing precise measurements of dissolved concentration, these passive samplers offer the benefit of easy storage and archiving, as well as cleaner extracts, which leads to improved detection limits of contaminants.³¹

Biomembranes are considered the first line of defense against xenobiotics in organisms, making them the primary target site for these chemicals.³² These membranes are primarily composed of phospholipids,³³ and the partition coefficients of these chemicals between the phospholipids and external media, such as water ($K_{lipid-water}$), drive their uptake through the biomembrane (equation 1).

$$K_{lipid-water} = \frac{C_{lipid}}{C_w} \quad (1)$$

When the truly dissolved concentration of a chemical C_w is equal to or exceeds the median lethal concentration (LC_{50}) in the external medium, the chemical can reach a critical level on the biomembrane that can cause toxic injury to the membrane's functioning. This concentration on the biomembrane is called the critical membrane burden or critical lipid burden (C_{lipid}^{crit}).

Hence, $C_w \rightarrow LC_{50}$, $C_{lipid} \rightarrow C_{lipid}^{crit}$, equation 1 can be rearranged in the following logarithmic form:

$$\log C_{lipid}^{crit} = \log LC_{50} + \log K_{lipid-water} \quad (2)$$

For many inert chemicals that interact with biomembranes nonspecifically, it has been observed that the critical membrane burden is fairly constant, with a median value of around 100 mmol/kg (-1 mol/kg in log units) for many aquatic species, such as fish.³⁴ In this case, equation 2 can be rewritten as.

$$-\log LC_{50} = 1 + \log K_{lipid-water} \quad (3)$$

A linear regression plot of $-\log LC_{50}$ against $\log K_{lipid-water}$ should result in intercept and slope close to unity. The Target Lipid Model (TLM) is based on this concept and assumes that the critical burden is primarily driven by the partition coefficient between the phospholipids and the water phase.³⁵

In this study, we borrowed the TLM framework to formulate the Target Plastic Model (TPM) as follows. We assume that the lipid in the TLM can be substituted with plastic, given its biomimetic nature, to create the TPM. In this case, the partition coefficient ($K_{plastic-water}$) is a ratio of concentration of a chemical in the plastic phase ($C_{plastic}$) to that in the water phase (C_w) at the equilibrium.

$$K_{plastic-water} = \frac{C_{plastic}}{C_w} \quad (4)$$

Assuming, when $C_w \rightarrow LC_{50}$, $C_{plastic} \rightarrow C_{plastic}^{crit}$, equation 4 becomes

$$K_{plastic-water} = \frac{C_{plastic}^{crit}}{LC_{50}} \quad (5)$$

Equation 5 can be rearranged in the following logarithmic form.

$$\log C_{plastic}^{crit} = \log LC_{50} + \log K_{plastic-water} \quad (6)$$

Critical plastic burden ($\log C_{plastic}^{crit}$) of chemicals can be calculated if the $\log LC_{50}$ and $\log K_{plastic-water}$ data of chemicals are available.

In this study, our hypothesis is that our new target plastic model can be parameterized with the $\log C_{plastic}^{crit}$, which is expected to remain relatively constant for a particular type of plastic, similar to its counterpart, $\log C_{lipid}^{crit}$. This parameterization will enable reliable predictions of LC_{50} with comparable performance to other widely used models, including the target lipid model. We propose three tests to evaluate this hypothesis:

1. First, we will explore the extent of similarity between the biotic phases involved in determining chemical toxicity and various types of plastic. We will evaluate the similarity between these phases based on the intermolecular interactions they experience while

- interacting with the chemicals, such as polarity, polarizability, hydrogen bonding, and dispersion interactions. This similarity will be inspected by dimensionality analysis, pair-wise correlation and linear regression analyses between $\log K_{plastic-water}$ and $\log K_{biotic\ phase-water}$ (partition coefficient between the biotic phases and water).
2. Second, we will estimate $\log C_{plastic}^{crit}$ values for various plastic phases using equation 6, and then calculate the median of the resulting distribution. We will also estimate $\log C_{plastic}^{crit}$ as the intercept obtained by linear regression of $\log K_{plastic-water}$ against $-\log LC_{50}$ values for a diverse set of chemicals. The variance in $\log C_{plastic}^{crit}$ compared to $\log C_{lipid}^{crit}$ will be examined to perform a plastic sensitivity distribution (PSD) analysis, similar to the concept of species sensitivity distribution (SSD) analysis.
 3. Third, we will compare the predicted LC_{50} values obtained by putting the $\log C_{plastic}^{crit}$ value into the TPM with the experimental values and predictions of other widely used models.

Successful testing of the hypothesis would provide a viable means for environmental scientists to establish a direct link between quantification of native and non-native pollutants on plastic phases in the environment and chemical risk assessment. This would also enhance the ability of scientists to design their passive sampling and dosing experiments in laboratory and field settings.

Material and Method

Experimental acute toxicity data of diverse chemicals reported for the fish were taken from the compilations available in the literature.³⁶ The compilation comprises experimental LC_{50} values for 949 chemicals. Experimental Abraham solute descriptors were obtained for these chemicals from the freely available online UFZ-LSER database,³⁷ resulting in complete sets of experimental Abraham solute parameters for 587 chemicals. Due to the lack of experimental Abraham solute parameters, the remaining 488 chemicals were not considered for further analysis. Based on their toxic mode of action, the final set of 587 chemicals were categorized into three groups: baseline toxicants or nonpolar narcotics, less-inert toxicants or polar narcotics, and reactive toxicants. To evaluate and validate the TPM, the following five sets were created from these three groups.

The first set, called the Nonpolar Narcotics or Baseline Evaluation Set, comprised 115 chemicals that are known to act via baseline or nonpolar narcotic mode of toxic action.³⁸ These chemicals belong to chemical families such as alkanes, alcohols, ketones, ethers, alkyl benzenes, and their chlorinated derivatives. The critical lipid and plastic burden values were derived using this set. This set is available as Table S1 in Supporting Information (SI).

The second set, called the Baseline Validation Set (Table S2), contains 132 chemicals that were predicted to follow baseline mode of toxic action according to the baseline model (BL).³⁶ These chemicals belong to diverse chemical families such as alkyl halides, alkenes, fluoroalcohol, chloroalcohol, diols, triols, alcohols, ethers, esters, carboxylic acids, amines, amides, carbamates, triazine, sulfides, disulfides, sulfoxides,

organophosphates, aromatic aldehydes, phthalates, halogenated phenols, nitrobenzenes, and pyridines. While some of these chemicals belong to chemical classes that are typically considered out of the domain of baseline toxicity, many of them are moderately to highly hydrophobic in nature and have been found to act via baseline modes of toxic action, despite containing polar functional groups, as reported in the literature.^{36,39} The critical plastic and lipid burden values derived using the Baseline Evaluation Set were used to predict the LC₅₀ values for this set, which was done to independently validate the TPM.

The third set, called the Polar Narcotics or Less-Inert Evaluation Set (Table S3), comprised 73 chemicals that are known to exert slightly higher toxicity than the baseline compounds via polar narcotic mode of toxic action.³⁶ These chemicals belong to classes such as phenols, halogenated phenols, and anilines. The critical plastic and lipid burdens of polar narcotic chemicals were obtained by analyzing this set.

The fourth set, called the Less-Inert Validation Set (Table S4), consisted of 128 chemicals, for which the critical lipid and plastic burden values obtained using the Polar Narcotic Evaluation Set were used to predict the LC₅₀ values. The predictions were then compared with the experimental LC₅₀ values to validate the TPM for polar narcotic chemicals. These chemicals were selected because the Less-Inert Model (LIM)'s prediction of LC₅₀ values deviated from the experimental LC₅₀ within 1 log unit.³⁶

The fifth set, called the Reactive Chemical Set (Table S5), comprised 75 chemicals and belonged to chemical families such as aldehydes, benzaldehydes, halogenated benzaldehydes, α,β -unsaturated esters, diamines, dinitrobenzenes, and their hydroxy derivatives. Reactive chemicals are known to covalently react with proteins or DNA in organisms and exert toxicities far above those predicted by the nonpolar and polar narcosis.⁴⁰

The partition coefficients of 587 chemicals were estimated for various biotic phases (phospholipid,⁴¹ storage lipid,⁴² muscle protein,⁴³ and blood protein⁴⁴), technical solvents (octanol⁴⁵ and triolein⁴⁶) and plastic phases (PDMS,⁴⁷ PA,⁴⁸ POM,⁴⁹ and PE¹⁶) using Abraham Solvation Model (ASM) equations, with the input of experimental Abraham solute parameters. The ASM was only available for the polyurethane ester (PU)-air system, and not for the PU-water system. Therefore, partition coefficient for the PU-water values were obtained by using a thermodynamic cycle between ASM estimated partition coefficients for the PU-air⁵⁰ and air-water⁵¹ systems. These data are available in Table S1-S5 in the SI.

The ASM equations for five other plastic types, namely polypropylene (PP), polystyrene (PS), polyvinyl chloride (PVC), ultra-high molecular weight polyethylene (UHMWPE), and high-density polyethylene (HDPE), were not available to estimate the partition coefficient between water phases and these plastic phases. Although some experimental plastic-water partition coefficient data for these plastic types were available for small sets of chemicals,⁵² the experimental toxicity data for these chemicals were very sparse (Table S6). The Target Plastic Model was also evaluated for these plastic phases. However, the evaluations for these plastic types could not be considered as reliable as those for the following five plastic types: PDMS, PA, POM, LDPE, and PU phases. This is mainly due to the limited experimental partitioning and toxicity data available for the former plastic types. In this study, these plastic phases (PP, PS, PVC, UHMWPE, and HDPE) are collectively referred to as "other plastics".

The critical burdens for three groups of chemicals, namely nonpolar narcotics, polar narcotics, and reactive toxicants, were estimated using three different methods. As a starting point, the first method involved assuming a critical burden of 100 mmol/kg of lipid or plastic for all groups, a value that is widely supported in the literature for lipids.⁵³ This method will be referred to as "100 mmol method" in the

subsequent text. In the second method, which will be referred to as the median method henceforth, the critical burden of each chemical group was calculated by using Equation 2 and Equation 6 to respectively calculate the critical lipid and critical plastic burden for each chemical in the Baseline Evaluation Set, Polar Narcotic Evaluation Set, and Reactive Chemical Set. The median of the distribution of these burden values was used to represent the critical lipid and plastic burdens for each group of chemicals. In the third method, hereafter referred to as the intercept method, $-\log LC_{50}$ values were linearly regressed against $\log K_{plastic-water}$ (in the case of plastics) and $\log K_{lipid-water}$ (in the case of lipid) for chemicals in each group. The intercept obtained from this linear regression respectively represents the critical lipid and plastic burdens according to Equation 2 and Equation 6. The slopes obtained in these cases indicate the strengths or sensitivity of plastic phases or lipid phases in deriving the toxicity of chemicals, which is similar to species sensitivity analysis towards toxicants using the target lipid model.³⁵ We evaluated these three methods of estimating the critical burdens to determine which one produces the best results for predicting toxicities.

To evaluate and validate the TPM, the critical plastic burden values obtained using the three methods described above and the partition coefficients for plastic-water were inserted into Equation 6 to predict the LC_{50} of chemicals in the five chemical sets. The resulting predictions were then compared with experimental values. The same procedure was repeated for the TLM (equation 2), allowing for a comparative analysis. In addition to experimental values, predictions of other widely used models were also used to compare the performance of the TPM. To estimate LC_{50} values for the chemicals, the ASM calibrated for several fish species⁵⁴ was used. Additionally, the US-EPA's ECOSAR module⁵⁵ was utilized, which categorizes the chemicals into ECOSAR classes based on their structure and functional groups. This categorization helps to identify whether the chemical follows a baseline mode of toxic action or a specific mode of action. Moreover, two previously published models,³⁶ BL and LIM, were used to predict LC_{50} values. These models are based on linear regression of $-\log LC_{50}$ against $\log K_{ow}$ and were respectively calibrated using baseline chemicals and less-inert chemicals.

In this study, diverse datasets were used, consisting of chemicals with varying chemical structures, including branched/unbranched/cyclic aliphatic and aromatic compounds, and several different types of functional groups. These chemicals exhibited a broad range of toxicities, hydrophobicities, and intermolecular interactions. Baseline Evaluation and Validation Sets covered a range of $-\log LC_{50}$ spanning six and seven orders of magnitude, respectively. The hydrophobicity of these sets ranged over six and eight orders of magnitude, respectively. Polar Narcotic Evaluation and Validation Sets encompassed toxicities with a range spanning more than nine and seven orders of magnitude, respectively, and their hydrophobicities covered more than six orders of magnitude. The Reactive Chemical Set consisted of chemicals with toxicities ranging from 1.79-7.34 log units and octanol-water partition coefficients ranging from -1.88 to 5.05 log units. The wide range of Abraham solute descriptors for the chemicals in these sets indicates their diversity in terms of intermolecular interactions.

Results and Discussion

Biomimetic nature of plastic phases

How closely the plastic phases mimic organisms depends on the extent to which the intermolecular interactions governing $\log K_{plastic-water}$ of chemicals resemble those controlling the $\log K_{biotic\ phase-water}$. This can be investigated using approaches based on dimensionality analysis, pairwise correlation and linear regression between $\log K_{plastic-water}$ and $\log K_{biotic\ phase-water}$. The results of these analyses are presented below.

Dimensionality analyses were performed to assess similarities among the biotic phases (phospholipid, storage lipid, muscle protein, and blood protein), technical solvents (octanol and triolein), plastic phases (PDMS, PA, POM, and LDPE), and toxicity endpoint (LC_{50}). The published system coefficients of the ASM equations available for these phases were used for the analysis. The first two dimensions obtained from the PCA test on the standardized system coefficients of ASM equations for toxicity, biotic phases, technical solvents, and plastic polymers represent 81.6% of the total information encoded in the ASM equations (Figure 1a). The ASM equations for these phases are calibrated using experimental datasets that are diverse both in terms of intermolecular interactions and chemical structures, making the similarity among these phases more representative than if they were based solely on the evaluation sets.

The Euclidean distance found between $-\log LC_{50}$ and $\log K_{PA-water}$ is the lowest compared to distances between $-\log LC_{50}$ and other phases observed on the PCA biplot representing the first two dimensions. The next closest plastic phase to $-\log LC_{50}$ is the POM polymer. This indicates the $\log K_{PA-water}$ and $\log K_{POM-water}$ are the closest allies of $-\log LC_{50}$ in terms of intermolecular interactions such as polarizability, polarity, hydrogen bonding interaction and dispersion forces. The PDMS and PE phases tend to respectively cluster with storage lipid and triolein phase indicating the chemical similarities between these phases, suggesting the PDMS and PE phases can be considered appropriate phases for estimating bioaccumulation. Octanol depicts good proximity to the phospholipid, which corroborates the previous success of TLM where octanol was taken as a proxy for phospholipid. Protein phases were found to be loners in this analysis.




Figure 1. Overlap in information between the biotic and plastic phases in terms of intermolecular interactions and partition coefficients. Panel (a) illustrates a cluster biplot obtained by performing principal component analysis on the system coefficients of ASM equations for biotic, technical solvents, plastic phases and toxicity endpoint (LC_{50}). PC 1 and PC 2, representing Principal Component 1 and 2 respectively, collectively account for 81.6% of the information, and Square cosine (\cos^2) reflects the quality of phase representation on the biplot. Panels (b-d) showcase the Pearson's pairwise correlation between the biotic and plastic phases for baseline ($n=115$), less-inert ($n=73$), and reactive ($n=75$) groups of chemicals.

The biomimetic nature of plastic phases can further be discerned by the linear relationship between the toxicity, plastic materials and biotic phases. For the three groups of chemicals, $\log K_{p-water}$ for the five types of plastics (PDMS, PA, POM, LDPE, and PU) exhibited a strong pairwise correlation with the partition coefficients for the biotic phases (phospholipid, storage lipid, muscle protein and blood proteins), and technical solvent (octanol) (Figure 1b-d). The degree of correlation of $-\log LC_{50}$ with the partition coefficients for biotic phases were in the same neighborhood as was found with the $\log K_{p-w}$. This supports the notion that the uptake of chemicals by these plastic phases mimics the uptake by the biotic phases. The degree of correlation between $-\log LC_{50}$ and $\log K_{p-water}$ for the five types of plastics was found to be very strong ($r > 0.95$) for baseline toxicants (Figure 1b), strong ($r > 0.816$) for less-inert toxicants (Figure 1c) and moderate ($r > 0.62$) for reactive toxicants (Figure 1d). This corroborates the

fact that the toxicity is mainly driven by the partitioning properties of baseline toxicants; whereas the contribution of partitioning in describing the toxicity for reactive chemicals decreases significantly.

The regression statistics such as R^2 and RMSE observed for the linear relationships between $-\log LC_{50}$ and $\log K_{plastic-water}$ were similar to the ones found for the relationships between $-\log LC_{50}$ and $\log K_{biotic\ phase-water}$ (Table 1). For these linear relationships, the intercept represents the $-\log C_{plastic}^{crit}$ (mol/kg of plastic) and slope represents the partitioning sensitivity of the plastic compared to the biomembrane. For the baseline toxicants ($n=115$), a linear regression of $-\log LC_{50}$ against the $\log K_{plastic-water}$ for the five types of plastics resulted in equations with intercept (1.43-3.80) and slope (0.78-0.971) with R^2 and RMSE in ranges of 0.947-0.908 and 0.307-0.406, respectively (Table 1). The intercept and slope obtained by linear regression of $-\log LC_{50}$ against the $\log K_{phospholipid-water}$ and $\log K_{ow}$ were 1.07 and 1.13, and 0.96 and 0.90, respectively. The linear relationship is similar to the TLMs for various species reported in the literature.³⁵ Phospholipid is considered a more accurate phase to use for calculations of the critical lipid burden of narcotic chemicals.³³ As evident by the comparisons of the fitting coefficients of the equations obtained by regressing $-\log LC_{50}$ against the $\log K_{storage\ lipid-water}$ and against $\log K_{phospholipid-water}$, the critical lipid burden is overestimated by a factor of 3.6, if the storage lipid, instead of the phospholipid, is considered as the target lipid. However, the phospholipid shows stronger partitioning sensitivity towards the baseline chemicals than the storage lipid. In many cases, the total lipid pool is used to normalize the toxicity endpoints.³³ For instance, taking into account the total lipid pool of an organism, rather than solely the phospholipid portion, may result in an overestimation of the critical burden by a factor of 2 for baseline toxicants.

Among plastic phases, PA demonstrated the best fit with $R^2=0.947$ and $RMSE=0.307$ log unit for baseline toxicants. The slopes observed for the relationships of $-\log LC_{50}$ against the $\log K_{plastic-water}$ of each PA and POM phases were in close agreement with slopes observed for TLMs based on phospholipid-water and octanol-water. This indicates the partitioning property of baseline toxicants for the PA and POM materials is similar to that for the target lipid. The critical plastic burden — as indicated by the intercepts — of baseline toxicants for the plastic phases were lower than the ones found for the target lipids.

In the case of less-inert toxicants ($n=73$), the fit statistics for the linear relationships $-\log LC_{50}$ and $\log K_{biotic\ phases-water}$ and between $-\log LC_{50}$ and $\log K_{plastic-water}$ are satisfactory, although not as good as those found for baseline toxicants. For the five types of plastics, values of RMSE and R^2 ranged from 0.26 to 0.49 log units and from 0.67 to 0.902, respectively, with PU showing the best fits and PDMS showing the least good fits. For the five types of biotic phases, values of RMSE and R^2 were in the ranges of 0.28 to 0.42 log units and 0.67 to 0.902, respectively. Regression of $-\log LC_{50}$ against $\log K_{ow}$ depicted $R^2 = 0.83$ and $RMSE = 0.344$ log unit. The relatively inferior statistics for the less-inert toxicants indicate that factors other than partitioning are important to account for the toxicity variability for this category of toxicants. The response sensitivities (slopes) are lower for less-inert chemicals than those found for baseline toxicants, and the critical burdens are lower for less-inert chemicals than those for baseline toxicants, which is expected as less-inert chemicals are more toxic than the baseline toxicants.

The regression of $-\log LC_{50}$ against $\log K_{biotic\ phases-water}$ and $\log K_{plastic-water}$ for reactive toxicants ($n=75$) yielded unsatisfactory fit statistics, with R^2 values ranging from 0.380-0.429 and RMSE values spanning a range of 0.886-0.923 log units. These results suggest that the reactive mode of toxic action for these chemicals is not well modeled by the partitioning models. Reactive toxicants are known to react

covalently with cellular components, such as proteins and DNA, and to form adducts that alter their structure and function, leading to toxicity. As a result, TPM and TLM, which have limitations of capturing such covalent interactions are expected to be ineffective for such chemicals.

Overall, these results support previous findings that plastic phases behave similarly to biotic phases in terms of exchanging baseline toxicants with the aqueous phases. This similarity remains adequate for less-inert chemicals but cannot be reliably established for reactive toxicants. Therefore, the biomimetic properties of plastic can be utilized to formulate TPM for the baseline and less-inert toxicants.

Table 2. Regression coefficients for the equation^a $-\log LC_{50} = -\log C_{biotic\ or\ plastic\ phase}^{crit} + m \log K_{biotic\ or\ plastic\ phase-water}$, based on data fitting from the baseline, less-inert and reactive sets.

Group	$\log K_{biotic\ or\ plastic\ phase-water}$	$-\log C_{biotic\ or\ plastic\ phase}^{crit}$	m	RMSE	R ²	n
Baseline Toxicants	Phospholipid-water	1.071±0.069	0.962 ± 0.022	0.322	0.942	115
	Storage lipid-water	1.627± 0.060	0.762 ± 0.018	0.331	0.939	115
	Pooled lipid-water	1.369 ± 0.061	0.854 ± 0.019	0.314	0.945	115
	Muscle Protein-water	1.800 ± 0.055	1.121 ± 0.026	0.325	0.941	115
	Serum protein-water	0.881 ± 0.070	1.162 ± 0.026	0.310	0.946	115
	Octanol-water	1.127±0.078	0.900 ± 0.024	0.367	0.925	115
	PDMS-water	1.933±0.066	0.780 ± 0.023	0.406	0.908	115
	PA-water	1.403±0.059	0.954 ± 0.021	0.307	0.947	115
	POM-water	1.774±0.059	0.971 ± 0.024	0.344	0.934	115
	PE-water	2.034±0.061	0.808 ± 0.023	0.387	0.916	115
	Polyurethane-water	3.803±0.030	0.828 ± 0.019	0.319	0.943	115
Less-Inert Toxicants	Phospholipid-water	2.153 ± 0.111	0.737 ± 0.032	0.292	0.880	73
	Storage lipid-water	3.144 ± 0.109	0.552 ± 0.038	0.420	0.751	73
	Pooled lipid-water	2.677 ± 0.111	0.646 ± 0.035	0.353	0.824	73
	Muscle Protein-water	2.767 ± 0.089	0.806 ± 0.037	0.302	0.872	73
	Serum protein-water	2.013 ± 0.109	0.857 ± 0.035	0.275	0.894	73
	Octanol-water	2.529 ± 0.115	0.628 ± 0.033	0.344	0.833	73
	PDMS-water	3.660 ± 0.095	0.575 ± 0.048	0.486	0.666	73
	PA-water	2.317 ± 0.096	0.739 ± 0.030	0.270	0.897	73
	POM-water	2.574± 0.093	0.806 ± 0.035	0.289	0.882	73
	PE-water	3.352 ± 0.096	0.662 ± 0.045	0.422	0.749	73
	Polyurethane-water	4.396 ± 0.032	0.674 ± 0.026	0.264	0.902	73
Reactive toxicants	Phospholipid-water	3.623 ± 0.1 ₅₀	0.609 ± 0.083	0.890	0.423	75
	Storage lipid-water	4.098 ± 0.113	0.439 ± 0.061	0.899	0.412	75
	Pooled lipid-water	3.887 ± 0.125	0.522 ± 0.071	0.888	0.426	75
	Muscle Protein-water	4.064 ± 0.116	0.694 ± 0.099	0.904	0.404	75
	Serum protein-water	3.528 ± 0.167	0.696 ± 0.101	0.912	0.394	75
	Octanol-water	3.723 ± 0.144	0.568 ± 0.081	0.905	0.404	75
	PDMS-water	4.374 ± 0.107	0.412 ± 0.062	0.923	0.380	75
	PA-water	3.816 ± 0.135	0.565 ± 0.081	0.907	0.401	75
	POM-water	3.995 ± 0.122	0.546 ± 0.079	0.912	0.395	75
	PE-water	4.339 ± 0.103	0.484 ± 0.065	0.886	0.429	75
	Polyurethane-water	4.396 ± 0.032	0.674 ± 0.026	0.907	0.401	75

^a The term $-\log C_{biotic\ or\ plastic\ phase}^{crit}$ denotes the critical burden of chemicals on the biotic or plastic phase, obtained as the intercept of a plot of $-\log LC_{50}$ against the partition coefficient for the biotic or plastic phase ($\log K_{biotic\ or\ plastic\ phase-water}$). The slope of the equation is represented by m .

Critical Plastic Burden Vis-à-vis Critical Lipid Burden

The critical burden of the baseline toxicants (n=115) on phospholipid was calculated using equation 2, resulting in a value of 108.5 mmol (-0.96 log unit). This value is close to the critical burden calculated for octanol (-0.88 log unit) using the same group of chemicals. However, the critical burden values for octanol were found to have a more dispersed distribution compared to those observed for phospholipid (Figure 2). Overall, the calculated critical burden values for both phospholipid and octanol fall within the range of literature-reported values.^{35,53} The critical plastic burden is lower than the critical lipid and octanol burden and it varies from 0.17-51.33 mmol for five different types of plastics, as shown in Figure 2a. Among five plastic types, PU is the most sensitive plastic, with a critical burden value more than two orders of magnitude lower than that of phospholipid and octanol phases. On the other hand, PA exhibits the highest critical plastic burden value.

As anticipated, the critical lipid and plastic burden of the polar narcotics (n=73) was found to be lower than that of nonpolar narcotics. This is attributed to the higher toxicity of polar narcotics compared to nonpolar narcotics. Specifically, the critical burden of polar narcotics on phospholipid (46.3 mmol/kg) and octanol (41.7 mmol/kg) was approximately half of the critical burdens observed for nonpolar narcotics on these phases. Moreover, the critical plastic burden for the five types of plastics ranges from 0.04 to 6.90 mmol/kg, which is smaller than the critical lipid burden for polar narcotics. Notably, PU was identified as the most sensitive plastic, requiring only a burden of 0.04 mmol/kg of PU to correspond to the median lethal concentration in the water phase. Conversely, PA exhibited the least sensitivity as a plastic phase, with a critical burden of 26.6 mmol/kg of PA.

Reactive toxicants are known to be highly toxic, and this is reflected in their critical burdens for the lipid and plastic phases. The difference in critical burden values between reactive toxicants and nonpolar narcotics is significant, with a difference of approximately two orders of magnitude observed for the lipid and plastic phases, except for PU. The PU phase, in particular, shows a much larger sensitivity to reactive toxicants. Conversely, the PA phase is the least sensitive plastic phase towards reactive toxicants, with a critical burden value of 6.8×10^{-4} mmol/kg.

The comparison of critical burdens for biotic and plastic phases estimated by the median and intercept methods is presented here. The intercept values are summarized in Table 1. The median values are depicted in Figure 2. For baseline toxicants, the critical burden values for phospholipid and octanol phases were comparable between the two methods. Furthermore, the median method produced critical burden values for PA, POM, and PU that were in good agreement with those obtained by the intercept method (Table 1). However, differences of up to 0.50 log units were observed for PDMS and PE phases when comparing the two methods. For less-inert chemicals, there were significant differences of more than one order of magnitude between the two methods for the critical octanol and lipid burdens. On the other hand, the differences between the two methods for the plastic phases except PE were not as large as those found for the lipid. For reactive toxicants, the critical burdens obtained from the two methods were similar in magnitude for these phases, except for the PU phase, which showed a difference of 1.6 log units.

These results show that while the median and intercept methods produce similar results for some phases and chemicals, there may be discrepancies for others. Overall, which method is more effective can be ascertained further by putting the values of critical burdens from both methods into the TLM and TPM to predict LC_{50} and comparing these predictions with the experimental values. This comparative test was performed and the results are presented in the next section of the paper.




Figure 2: Distribution of critical burdens for various toxicants on lipid, octanol, and plastic phases. Boxplots are shown for (a) baseline toxicants, (b) less inert toxicants, and (c) reactive toxicants, on PDMS, PA, POM, PE, and PU. Panel (d) shows the distributions for other plastic phases (PP, PS, PVC, UHMWPE, and HDPE), for which evaluation data were limited. The red colored + symbols and the black horizontal lines within the boxes indicate the mean and median values of the critical burden distributions, respectively.

Prediction of Acute Toxicity from Critical Plastic Burden

The critical burdens for the three groups of chemicals were estimated using three different methods, and their effects on the accuracy of LC₅₀ predictions were evaluated. For the baseline toxicants (n=115), the TLM showed good agreement between experimental and predicted values of LC₅₀ (Figure 3a). The input of critical burden values using three different methods into the TLM did not significantly affect the accuracy of the model (Figure 3a, S1 and S2). As expected, the TLM based on phospholipid performed better than the TLM based on octanol, as phospholipids are better representatives of membrane lipids.

The target plastic model exhibited a close agreement between its predicted values and experimental values for the five types of plastics, except for PU, which showed systematic deviations of 2.84 log units from the experimental values when using the 100 mmol method of critical burden. However, for the median and intercept methods, PU also demonstrated good agreement between predicted and experimental values (RMSE=0.42 log unit). Overall, the input of critical burden values estimated using the median method into the TPM performed better than the 100 mmol and intercept methods. Thus, the median method of estimating critical burden is recommended for input to the TPM.

Among plastic phases, the target PA and POM models performed the best with the lowest RMSE values of 0.311 and 0.343 log units for median method, respectively. The prediction accuracy was on par with the target phospholipid model. The performance of target PE model and target PU model was similar to the target octanol model. The target PDMS model exhibited RMSE = 0.538 log units when its predictions were compared with the experimental values.

The performance of the TPM was also compared with other models such as ASM, ECOSAR, BL, and LIM using the Baseline Evaluation Set (n=115). These models exhibited RMSE values ranging from 0.349-0.306 log units (Figure 3a). The performance of these models was similar to the TLM and TPM. It should be noted that the experimental dataset utilized to evaluate the TLM and TPM had already been employed to train regression models such as the BL and LIM. When comparing the predictions of two models against experimental data using the same training data, the model that has been trained on that data is expected to perform better than the non-fitted model, such as the TLM and TPM.

The Baseline Validation Set comprising 132 chemicals was used to evaluate the performance of the TPM compared to the TLM, ASM, ECOSAR, BL and LIM (Figure 3b). The chemicals in the validation set were predicted to follow a baseline mode of toxic action, as shown by the toxic ratio or residual (experimental LC₅₀ minus predicted LC₅₀ by the baseline regression model) values < 1 log unit. Unlike the test set of 115 chemicals, the validation set was not used to compute the critical burdens for plastic, lipid, and octanol using median and intercept methods, thereby providing an unbiased evaluation.

The target phospholipid and octanol models showed a good agreement between experimental and predicted values, with RMSE values of 0.42 and 0.44 log units, respectively (Figure 3b). The performance of the target PA model and POM model was similar to that of the target phospholipid and octanol models, with RMSE values of 0.45 and 0.49 log units, respectively. However, the target PU model and target PE model exhibited higher RMSE values of 0.76 and 0.93 log units, respectively. The poorest performing model was the target PDMS model, with an RMSE value of 1.11 log units (Figure 3b).

In comparison, the ASM and ECOSAR models had RMSE values of 0.42 and 0.53 log units, respectively, for the Baseline Validation Set. Notably, predicted LC50 values obtained by the Target PDMS and Target PE Models for chemicals belonging to various classes, such as halogenated alcohols, diols, α,β -unsaturated alcohols, alcohol-ethers, diol-ethers, amines, amides, sulfoxides, and benzoic acids, differed from the experimental values by more than one order of magnitude. Many of these chemicals were hydrophilic and polar in nature. To investigate the performance of the models for chemicals with $\log K_{ow} > 3$, a subset of the validation set with 47 chemicals was analyzed. The results showed that the performance of the PDMS, PE, and PU models was improved by 0.48, 0.31, and 0.24 log units, respectively, for this subset.

The predictive performance of TLM and TPM for polar narcotics is slightly inferior to that of nonpolar narcotics (Figure 3c). Other models, such as ASM, ECOSAR, and BL, also demonstrate relatively poor performance. This was expected because polar narcotics or less-inert toxicants may have specific interactions with the target organ that are not adequately represented by partitioning processes. The target phospholipid model and target octanol model have RMSE values of 0.40 and 0.57 log units, respectively. The target phospholipid model is more accurate than the target octanol model because it better represents membrane lipids. The target plastic model has an RMSE range of 0.347-0.697 log units, with the POM being the most accurate and the PDMS being the least accurate. The LIM outperforms all other models with an RMSE of 0.335. However, it should be noted that unlike TLM and TPM, LIM is a fitted model trained on the same dataset used for comparison, favoring its performance.

The Less-Inert Validation Set of 128 chemicals was used to validate the TPM for polar narcotics. These chemicals were predicted to follow a less inert mode of toxic action based on the predictions of LIM. Like the Baseline Validation Set, the chemicals in the Less-Inert Validation Set were not used to calculate the critical plastic, lipid, and octanol burdens, which helped ensure an unbiased evaluation of the models. The results showed that the predictions of the target phospholipid model and target octanol model were in good agreement with the experimental values for the 128 chemicals, with RMSE values of 0.46 and 0.44 log unit, respectively (Figure 3d). However, the TPM exhibited a wider range of RMSE values, ranging from 0.56 to 1.11 log unit, with PA performing the best and PDMS performing the worst when compared to the experimental values of LC_{50} . In comparison, ASM and ECOSAR had RMSE values of 0.55 and 0.46 log units, respectively, for the same set of chemicals. Furthermore, the residuals for chemicals belonging to classes such as unsaturated alkenes, amine-alcohols, halogenated nitrobenzenes, and nitrogen-containing biphenyls were significantly higher for the PDMS and PE plastics.

Finally, the performance of various models, including the TLM, TPM, ASM, ECOSAR, BL, and LIM, were evaluated for reactive toxicants ($n=75$). However, none of these models performed well for these chemicals (Figure S3). When the predicted values from these models were compared with experimental values for 75 reactive chemicals, the resulting RMSE values were all over 1 log unit. This poor performance was anticipated since reactive toxicities are influenced by specific interaction parameters that are not accounted for in any of the models studied.




Figure 3: Comparison of LC_{50} predictions by the Target Plastic Model based on five plastic types (PDMS, PA, POM, PE, and PU) with experimental LC_{50} values for fish. The figure also displays predictions from other models, such as the Target Lipid Model based on phospholipid and octanol, as well as ASM, ECOSAR, BL, and LIM. Panels (a) and (c) present model evaluation using evaluation sets of baseline toxicants ($n=115$) and less-inert toxicants ($n=73$), respectively. Panels (b) and (d) show model validation using validation sets of baseline toxicants ($n=132$) and less-inert toxicants ($n=128$), respectively. RMSE values in log units are provided for each model in each panel, obtained by comparing predicted LC_{50} values with experimental values. The dotted line in the middle of each panel represents 1:1 agreement, while the upper and lower dotted lines indicate 1:2 agreement between the experimental and predicted LC_{50} values.

Evaluation of Target Model for Other Plastics

The critical plastic burdens of chemicals on additional plastic types, such as polypropylene (PP), polystyrene (PS), polyvinyl chloride (PVC), ultra-high-molecular-weight polyethylene (UHMWPE), and high-density polyethylene (HDPE), were estimated tentatively. The values for the critical plastic burden on these plastic types fell within the range of 0.01 – 63.89 mmol/kg of plastic (Fig. 2d), with PS demonstrating the lowest and HDPE exhibiting the highest chemical critical burden. PVC demonstrated the highest variance in the distribution of critical burden, covering a range of more than seven orders of magnitude for 32 chemicals. This variance may be due to the fact that these chemicals belong to different chemical classes such as PCBs, PAHs, and pharmaceuticals, which follow different modes of toxic actions. The critical burden of chemicals on HDPE (15.65 mmol/kg) was found to be similar to that on LDPE (24.95

mmol/kg), indicating that the density of polyethylene does not strongly affect the partitioning behavior of chemicals considered here.

It is important to note that the ASM estimated LC₅₀ values were used to compute the critical burdens of chemicals on these plastic types due to a lack of experimental values. Additionally, the sample size and structural diversity of the chemicals used to evaluate these additional plastic types were limited compared to the plastic types used previously to evaluate and validate the target plastic model. Hence, the results for these additional plastic types should be interpreted with caution as they were based on estimated LC₅₀ values due to the lack of experimental data and the limited sample size and structural diversity of chemicals for these plastic types.

The critical plastic burdens calculated for each plastic type were used to predict LC₅₀ values using TPM. The predicted LC₅₀ values were then compared to the ASM estimated values instead of experimental values due to the limited data available. The results showed a good agreement between the predicted LC₅₀ values by the target PP model and by the ASM, with an RMSE of 0.45 log units. For the same set of chemicals (n=9), the target phospholipid and octanol models showed RMSEs of 0.11 and 0.17 log units, respectively. ECOSAR's predictions also matched well with ASM predictions, with an RMSE of 0.28 log units. However, the BL showed an RMSE of 0.61 log units with respect to the ASM predicted values for these chemicals. In this comparison, PCB 187 and 128 showed the highest residuals for the target PP model, which may be attributed to the poor data quality of measured partition coefficients of these strongly hydrophobic chemicals between PP and the water phase.

LC₅₀ values predicted by the target PS model did not match favorably with the ASM predicted LC₅₀ values based on 8 chemicals, showing an RMSE of 1.52 log units. In particular, nonylphenol showed a deviation of more than 2.75 log units from the ASM predicted value. Interestingly, the ECOSAR class for this chemical is phenols, which generally exhibit excess toxicity compared to the baseline toxicity of neutral organics. Additionally, significant deviations between the predictions of the two models were observed for very hydrophobic PCB congeners such as PCB 171, 200, and 206. Although other models worked well for these chemicals, the deviation for the TPM may be attributed to the poor quality of the reported experimental plastic-water partition coefficients or to factors other than partitioning responsible for the exchange of chemicals between the polystyrene and water phases.

The performance of the target PVC model and ASM in predicting LC₅₀ values for compounds from various chemical families was evaluated. Overall, the comparison of the predictions showed that the agreement between the models was not favorable, with an RMSE of 1.67 log units. However, this comparison included strongly hydrophobic chemicals such as PCB congeners and DEHP (di-2-ethylhexyl phthalate), which showed the highest deviations, not only for the target PVC model but also for other models used in this study. In contrast, the target PVC model's performance was satisfactory for moderately hydrophobic PCB congeners. For DEHP, even the ASM failed, as the difference between the experimental and ASM-predicted LC₅₀ values was more than five orders of magnitude. For PCB 209, with a log *K*_{ow} of 8.27, the deviations for the PVC model were also more than five orders of magnitude. For most of the pharmaceutical drugs, there was good agreement between the predicted LC₅₀ values of the PVC model and ASM. However, there were exceptions, such as flunitrazepam and chlorpromazine, which showed significantly higher residuals among the pharmaceutical drugs. Huge deviations between the predictions of two models for certain chemicals may be rationalized by considering specific toxic nature of the chemicals and/or the uncertain data quality stemming from the experimental challenges of measuring

such chemicals in partitioning and toxicity experiments. For example, the ECOSAR class identified for DEHP is esters, implying it might be following a mode of toxic action other than baseline toxicity, and with a $\log K_{ow}$ value of 7.6, it also belongs to the strongly hydrophobic category. Measuring physicochemical properties of strongly hydrophobic chemicals free from experimental artifacts is a challenging task. Similarly, flunitrazepam and chlorpromazine follow a mode of toxic action other than baseline, as indicated by their ECOSAR classes as amides and aliphatic amines, respectively. Such specific toxic modes of actions are difficult to account for by partitioning processes alone.

The performance of the target HDPE model in predicting LC_{50} values for chemicals was also evaluated. The comparison of the predictions revealed that the predicted LC_{50} by target HDPE model compared favorably with the ASM-predicted LC_{50} , as shown by its RMSE of 0.38 log units. Other models also performed well for this chemical set. However, the dataset used for this evaluation comprised only hydrocarbons with moderate hydrophobicities ($\log K_{ow}$ ranging from 2.73-5.81). These chemicals are known to follow the baseline toxic mode of action, and their measured partition coefficient values between the plastic and water phases are expected to not suffer too much from experimental artifacts. Therefore, the better performance of HDPE for such chemicals is not surprising. It is worth noting that the dataset used to evaluate the UHMWPE was not quite meaningful, as it only comprised three chemicals, with two belonging to the strongly hydrophobic category and one belonging to a mode of toxic action other than baseline toxicity.

Implications and Outlook

The framework of TPM developed in this study has several potential benefits for environmental scientists, which are described below.

Firstly, like TLM, the TPM can be applied to estimate the toxic unit. The toxic unit is a convenient way of calculating the toxicity of mixtures if components of the mixtures follow the same mode of toxic action. It is defined as.

$$TU = \frac{C_w}{LC_{50}} \quad (7)$$

Equation 4 and 5 can be rearranged in favor of C_w and LC_{50} for further insertion into equation 7 to obtain the following simplified form.

$$TU = \frac{C_{plastic}}{C_{plastic}^{crit}} \quad (8)$$

We can quantify the chemicals detected on environmental plastics or passive samplers and normalize these quantities to the critical plastic burden. This allows calculation of toxic units, which can be additive for baseline toxicants and used to determine mixture toxicity using equation 9 and 10.

$$\sum_1^n TU = TU_1 + TU_2 + TU_3 + \dots TU_n \quad (9)$$

$$\sum_1^n TU = \frac{C_{plastic_1} + C_{plastic_2} + C_{plastic_3} + \dots C_{plastic_n}}{C_{plastic}^{crit}} \quad (10)$$

This approach offers a significant advantage over existing methods for calculating mixture toxicity, as it does not rely on the availability of LC₅₀ and plastic-water partition coefficient data, which can be limited, particularly for emerging chemicals. Only the knowledge of critical plastic burden is required to calculate the toxic unit. For instance, while several hydrophobic micropollutants have been quantified in environmental plastic samples from Swiss surface waters,⁵⁶ more insight could be gained from these studies by comparing the quantities with critical plastic burdens to calculate toxic units for risk assessment. Given the success of TLM in predicting the toxicity of oil spills,³⁵ TPM can be considered a promising method for estimating the toxicity of complex mixtures resulting from oil spills before and after weathering.

Secondly, the TPM can be used as an animal-alternative technique for finding LC₅₀ values for new chemicals in the laboratory. From the plastic phases considered in this study, it is clear that PA and POM are the most appropriate phases to be used as an alternative to fish in determining LC₅₀ values for baseline toxicants. The target plastic model based on these plastic phases was able to predict the LC₅₀ values for a wide range of chemicals within the range of experimental error.

Thirdly, the TPM can be used in passive sampling-based field studies. Passive sampling is becoming increasingly popular among environmental scientists and regulatory authorities, as it provides more insights into pollution risks by simultaneously determining the environmental levels and toxicities of detected chemicals. This approach can be particularly useful in situations such as marine oil spills, with complex mixtures of hydrocarbons.

Finally, the TPM may be a useful tool for designing passive dosing-based toxicity experiments, as passive dosing techniques provide precise control over exposure concentrations. By utilizing the TPM, scientists can pre-select appropriate passive doses that will encompass the critical plastic burden, leading to expected responses and resulting in a well-defined dose-response curve.

The target plastic model developed in this study has several limitations that must be considered. Firstly, the model is not applicable to ionizable or reactive chemicals because they can undergo chemical reactions or ionization that affect their behavior and distribution in plastic phases, which are not fully explainable through equilibrium partitioning theory. Secondly, the model does not account for non-persistent chemicals that undergo metabolism or physical or chemical transformation, which can change over time, making it difficult to predict their behavior accurately in plastic phases. Thirdly, the model only considers passive exposure to chemicals and disregards active exposure, which may limit its applicability in certain environmental settings. Finally, the model does not consider adsorption of chemicals on plastics, only their absorption (partitioning), which may limit its accuracy for some plastic materials as adsorption could be a crucial mechanism for their behavior in the environment. Overall, although the TPM shows potential for environmental chemistry applications, its limitations should be carefully evaluated when interpreting results and applying it to various scenarios.

This study presents promising research avenues for the further development of the target plastic model. Firstly, the model can be extended beyond fish to other aquatic species, following the success of the target lipid model for several aquatic species. Future studies could investigate the applicability of the

target plastic model to a wider range of species, which could enhance its usefulness in environmental risk assessments.

Secondly, the target plastic model could be extended beyond acute toxicity to chronic toxicity levels. Similar to the target lipid model, the target plastic model can be applied to derive concentrations above which 95% of species should be protected (HC5 values) for organic chemicals. These HC5 values can then be used to estimate predicted no-effect concentrations (PNECs), which could improve the accuracy and reliability of environmental risk assessments.

Overall, these research avenues have the potential to enhance the utility of the target plastic model in environmental chemistry, providing new insights into the toxicity and behavior of organic chemicals associated with the plastic. Further research in these areas could lead to the development of more effective and efficient risk assessment methods, contributing to the protection of human and ecological health.

In summary, this study has shown that plastic phases exhibit a behavior similar to biotic phases, allowing the development of a target plastic model based on the theoretical framework of the target lipid model. The target plastic model, specifically based on PA and POM plastic types, successfully predicted the acute toxicity endpoint for fish within the range of experimental errors. Environmental chemists can utilize the critical plastic burdens presented in this study for polar and nonpolar toxicants to rapidly estimate the toxicity of hundreds or thousands of chemicals associated with plastic.

Supporting Information

The Supporting Information for this manuscript is available in the form of an Excel spreadsheet that contains information about the evaluation and validation sets used to develop the target plastic model.

Acknowledgement

The authors acknowledge funding of the project P-LEACH (<https://www.ufz.de/P-LEACH>) from the Helmholtz Association, Innovation Pool of the Research Field Earth and Environment in support of the Research Program “Changing Earth – Sustaining our Future.”

References

- (1) Zhu, X.; Huang, W.; Fang, M.; Liao, Z.; Wang, Y.; Xu, L.; Mu, Q.; Shi, C.; Lu, C.; Deng, H.; et al. Airborne Microplastic Concentrations in Five Megacities of Northern and Southeast China. *Environ. Sci. Technol.* **2021**, *55* (19).
- (2) Zhang, Q.; Zhao, Y.; Du, F.; Cai, H.; Wang, G.; Shi, H. Microplastic Fallout in Different Indoor Environments. *Environ. Sci. Technol.* **2020**, *54* (11).
- (3) Issac, M. N.; Kandasubramanian, B. Effect of Microplastics in Water and Aquatic Systems. *Environmental Science and Pollution Research*. 2021, pp 19544–19562.
- (4) Guo, J. J.; Huang, X. P.; Xiang, L.; Wang, Y. Z.; Li, Y. W.; Li, H.; Cai, Q. Y.; Mo, C. H.; Wong, M. H. Source, Migration and Toxicology of Microplastics in Soil. *Environment International*. 2020.
- (5) Anbumani, S.; Kakkar, P. Ecotoxicological Effects of Microplastics on Biota: A Review. *Environmental Science and Pollution Research*. 2018, pp 14373–14396.
- (6) Mamun, A. Al; Prasetya, T. A. E.; Dewi, I. R.; Ahmad, M. Microplastics in Human Food Chains: Food Becoming a Threat to Health Safety. *Science of the Total Environment*. 2023.
- (7) Leslie, H. A.; van Velzen, M. J. M.; Brandsma, S. H.; Vethaak, A. D.; Garcia-Vallejo, J. J.; Lamoree, M. H. Discovery and Quantification of Plastic Particle Pollution in Human Blood. *Environ. Int.* **2022**, *163*.
- (8) Ragusa, A.; Svelato, A.; Santacroce, C.; Catalano, P.; Notarstefano, V.; Carnevali, O.; Papa, F.; Rongioletti, M. C. A.; Baiocco, F.; Draghi, S.; et al. Plasticenta: First Evidence of Microplastics in Human Placenta. *Environment International*. 2021.
- (9) Kögel, T.; Bjørøy, Ø.; Toto, B.; Bienfait, A. M.; Sanden, M. Micro- and Nanoplastic Toxicity on Aquatic Life: Determining Factors. *Science of the Total Environment*. 2020.
- (10) Junaid, M.; Siddiqui, J. A.; Sadaf, M.; Liu, S.; Wang, J. Enrichment and Dissemination of Bacterial Pathogens by Microplastics in the Aquatic Environment. *Science of the Total Environment*. 2022.
- (11) Pham, D. N.; Clark, L.; Li, M. Microplastics as Hubs Enriching Antibiotic-Resistant Bacteria and Pathogens in Municipal Activated Sludge. *J. Hazard. Mater. Lett.* **2021**, *2*.
- (12) Zimmermann, L.; Göttlich, S.; Oehlmann, J.; Wagner, M.; Völker, C. What Are the Drivers of Microplastic Toxicity? Comparing the Toxicity of Plastic Chemicals and Particles to *Daphnia Magna*. *Environ. Pollut.* **2020**, *267*.
- (13) Bridson, J. H.; Gaugler, E. C.; Smith, D. A.; Northcott, G. L.; Gaw, S. Leaching and Extraction of Additives from Plastic Pollution to Inform Environmental Risk: A Multidisciplinary Review of Analytical Approaches. *Journal of Hazardous Materials*. 2021.
- (14) Iqbal, R.; Khan, M. T.; Bilal, H.; Aslam, M. M.; Khan, I. A.; Raja, S.; Arslan, M.; Nguyen, P. M. Microplastics as Vectors of Environmental Contaminants: Interactions in the Natural Ecosystems. *Hum. Ecol. Risk Assess.* **2022**, *28* (9), 1022–1042.
- (15) Scopetani, C.; Cincinelli, A.; Martellini, T.; Lombardini, E.; Ciofini, A.; Fortunati, A.; Pasquali, V.;

- Ciattini, S.; Ugolini, A. Ingested Microplastic as a Two-Way Transporter for PBDEs in *Talitrus Saltator*. *Environ. Res.* **2018**, *167*, 411–417.
- (16) Khawar, M. I.; Nabi, D. Relook on the Linear Free Energy Relationships Describing the Partitioning Behavior of Diverse Chemicals for Polyethylene Water Passive Samplers. *ACS Omega* **2021**.
- (17) Müller, A.; Becker, R.; Dorgerloh, U.; Simon, F. G.; Braun, U. The Effect of Polymer Aging on the Uptake of Fuel Aromatics and Ethers by Microplastics. *Environ. Pollut.* **2018**, *240*.
- (18) Yao, S.; Cao, H.; Arp, H. P. H.; Li, J.; Bian, Y.; Xie, Z.; Cherubini, F.; Jiang, X.; Song, Y. The Role of Crystallinity and Particle Morphology on the Sorption of Dibutyl Phthalate on Polyethylene Microplastics: Implications for the Behavior of Phthalate Plastic Additives. *Environ. Pollut.* **2021**, *283*.
- (19) Liu, P.; Zhan, X.; Wu, X.; Li, J.; Wang, H.; Gao, S. Effect of Weathering on Environmental Behavior of Microplastics: Properties, Sorption and Potential Risks. *Chemosphere*. 2020.
- (20) Huckins, J. N.; Petty, J. D.; Booij, K. *Monitors of Organic Chemicals in the Environment: Semipermeable Membrane Devices*; Springer Science & Business Media, 2006.
- (21) Lohmann, R.; Booij, K.; Smedes, F.; Vrana, B. Use of Passive Sampling Devices for Monitoring and Compliance Checking of POP Concentrations in Water. *Environ. Sci. Pollut. Res.* **2012**, *19* (6), 1885–1895.
- (22) Leslie, H. A.; Oosthoek, A. J. P.; Busser, F. J. M.; Kraak, M. H. S.; Hermens, J. L. M. Biomimetic Solid-Phase Microextraction to Predict Body Residues and Toxicity of Chemicals That Act by Narcosis. *Environ. Toxicol. Chem.* **2002**, *21* (2), 229–234.
- (23) Escher, B. I.; Hermens, J. L. M. Peer Reviewed: Internal Exposure: Linking Bioavailability to Effects. *Environ. Sci. Technol.* **2004**, *38* (23), 455A–462A.
- (24) Taylor, A. C.; Fones, G. R.; Vrana, B.; Mills, G. A. Applications for Passive Sampling of Hydrophobic Organic Contaminants in Water—A Review. *Critical Reviews in Analytical Chemistry*. 2021, pp 20–54.
- (25) Smith, K. E. C.; Oostingh, G. J.; Mayer, P. Passive Dosing for Producing Defined and Constant Exposure of Hydrophobic Organic Compounds during *In Vitro* Toxicity Tests. *Chem. Res. Toxicol.* **2010**, *23* (1), 55–65.
- (26) O’Connell, S. G.; Kerkvliet, N. I.; Carozza, S.; Rohlman, D.; Pennington, J.; Anderson, K. A. *In Vivo* Contaminant Partitioning to Silicone Implants: Implications for Use in Biomonitoring and Body Burden. *Environ. Int.* **2015**, *85*, 182–188.
- (27) Bera, G.; Parkerton, T.; Redman, A.; Turner, N. R.; Renegar, D. A.; Sericano, J. L.; Knap, A. H. Passive Dosing Yields Dissolved Aqueous Exposures of Crude Oil Comparable to the CROSERF (Chemical Response to Oil Spill: Ecological Effects Research Forum) Water Accommodated Fraction Method. *Environ. Toxicol. Chem.* **2018**, *37* (11), 2810–2819.
- (28) Smith, K. E. C.; Dom, N.; Blust, R.; Mayer, P. Controlling and Maintaining Exposure of Hydrophobic Organic Compounds in Aquatic Toxicity Tests by Passive Dosing. *Aquat. Toxicol.* **2010**, *98* (1), 15–24.
- (29) Adolffson-Erici, M.; Åkerman, G.; Mclachlan, M. S. Measuring Bioconcentration Factors in Fish

- Using Exposure to Multiple Chemicals and Internal Benchmarking to Correct for Growth Dilution. *Environ. Toxicol. Chem.* **2012**, *31* (8), 1853–1860.
- (30) ter Laak, T. L.; Durjava, M.; Struijs, J.; Hermens, J. L. M. Solid Phase Dosing and Sampling Technique to Determine Partition Coefficients of Hydrophobic Chemicals in Complex Matrixes. *Environ. Sci. Technol.* **2005**, *39* (10), 3736.
- (31) Harman, C.; Allan, I.; Thomas, K. V. Passive Sampling of Organic Contaminants in Waters. In *Comprehensive Sampling and Sample Preparation*; 2012.
- (32) Van Wezel, A. P.; Opperhuizen, A. Narcosis Due to Environmental Pollutants in Aquatic Organisms: Residue-Based Toxicity, Mechanisms, and Membrane Burdens. *Crit. Rev. Toxicol.* **1995**, *25* (3), 255–279.
- (33) Endo, S.; Escher, B. I.; Goss, K. U. Capacities of Membrane Lipids to Accumulate Neutral Organic Chemicals. *Environ. Sci. Technol.* **2011**, *45* (14), 5912–5921.
- (34) Bittermann, K.; Spycher, S.; Goss, K. U. Comparison of Different Models Predicting the Phospholipid-Membrane Water Partition Coefficients of Charged Compounds. *Chemosphere* **2016**, *144*, 382–391.
- (35) Kipka, U.; Di Toro, D. M. Technical Basis for Polar and Nonpolar Narcotic Chemicals and Polycyclic Aromatic Hydrocarbon Criteria. III. A Polyparameter Model for Target Lipid Partitioning. *Environ. Toxicol. Chem.* **2009**, *28* (7), 1429–1438.
- (36) Wang, X. H.; Yu, Y.; Huang, T.; Qin, W. C.; Su, L. M.; Zhao, Y. H. Comparison of Toxicities to *Vibrio Fischeri* and Fish Based on Discrimination of Excess Toxicity from Baseline Level. *PLoS One* **2016**, *11* (2).
- (37) Endo, S.; Watanabe, N.; Ulrich, N.; Bronner, G.; Goss, K.-U. UFZ-LSER database v 2.1.
- (38) Verhaar, H. J. M.; van Leeuwen, C. J.; Hermens, J. L. M. Classifying Environmental Pollutants. 1: Structure-Activity Relationships for Prediction of Aquatic Toxicity. *Chemosphere* **1992**, *25* (4), 471–491.
- (39) Zhao, Y. H.; Zhang, X. J.; Wen, Y.; Sun, F. T.; Guo, Z.; Qin, W. C.; Qin, H. W.; Xu, J. L.; Sheng, L. X.; Abraham, M. H. Toxicity of Organic Chemicals to *Tetrahymena Pyriformis*: Effect of Polarity and Ionization on Toxicity. *Chemosphere* **2010**, *79* (1), 72–77.
- (40) Enoch, S. J.; Ellison, C. M.; Schultz, T. W.; Cronin, M. T. D. A Review of the Electrophilic Reaction Chemistry Involved in Covalent Protein Binding Relevant to Toxicity. *Critical Reviews in Toxicology*. 2011, pp 783–802.
- (41) Endo, S.; Escher, B. I.; Goss, K.-U. Capacities of Membrane Lipids to Accumulate Neutral Organic Chemicals. *Environ. Sci. Technol.* **2011**, *45* (14), 5912–5921.
- (42) Geisler, A.; Endo, S.; Goss, K. U. Partitioning of Organic Chemicals to Storage Lipids: Elucidating the Dependence on Fatty Acid Composition and Temperature. *Environ. Sci. Technol.* **2012**, *46* (17), 9519–9524.
- (43) Endo, S.; Bauerfeind, J.; Goss, K. U. Partitioning of Neutral Organic Compounds to Structural Proteins. *Environ. Sci. Technol.* **2012**, *46* (22), 12697–12703.
- (44) Endo, S.; Goss, K. U. Serum Albumin Binding of Structurally Diverse Neutral Organic Compounds:

- Data and Models. *Chem. Res. Toxicol.* **2011**, *24* (12).
- (45) Abraham, M. H.; Chadha, H. S.; Whiting, G. S.; Mitchell, R. C. Hydrogen Bonding. 32. An Analysis of Water-octanol and Water-alkane Partitioning and the $\Delta \log p$ Parameter of Seiler. *J. Pharm. Sci.* **1994**, *83* (8), 1085–1100.
- (46) Abraham, M. H.; Acree, W. E. Equations for Water-Triolein Partition Coefficients for Neutral Species; Comparison with Other Water-Solvent Partitions, and Environmental and Toxicological Processes. *Chemosphere* **2016**, *154*, 48–54.
- (47) Sprunger, L.; Proctor, A.; Acree, W. E.; Abraham, M. H. Characterization of the Sorption of Gaseous and Organic Solutes onto Polydimethyl Siloxane Solid-Phase Microextraction Surfaces Using the Abraham Model. *J. Chromatogr. A* **2007**, *1175* (2), 162–173.
- (48) Endo, S.; Droge, S. T. J.; Goss, K. U. Polyparameter Linear Free Energy Models for Polyacrylate Fiber-Water Partition Coefficients to Evaluate the Efficiency of Solid-Phase Microextraction. *Anal. Chem.* **2011**, *83* (4), 1394–1400.
- (49) Endo, S.; Hale, S. E.; Goss, K. U.; Arp, H. P. H. Equilibrium Partition Coefficients of Diverse Polar and Nonpolar Organic Compounds to Polyoxymethylene (POM) Passive Sampling Devices. *Environ. Sci. Technol.* **2011**, *45* (23), 10124–10132.
- (50) Schneider, M.; Goss, K. U. Systematic Investigation of the Sorption Properties of Tenax TA, Chromosorb 106, Porapak n, and Carboxpak F. *Anal. Chem.* **2009**, *81* (8), 3017–3021.
- (51) Goss, K. U. Predicting the Equilibrium Partitioning of Organic Compounds Using Just One Linear Solvation Energy Relationship (LSER). *Fluid Phase Equilib.* **2005**, *233* (1), 19–22.
- (52) O'Connor, I. A.; Golsteijn, L.; Hendriks, A. J. Review of the Partitioning of Chemicals into Different Plastics: Consequences for the Risk Assessment of Marine Plastic Debris. *Marine Pollution Bulletin*. 2016, pp 17–24.
- (53) Bittermann, K.; Goss, K. U. Assessing the Toxicity of Ionic Liquids – Application of the Critical Membrane Concentration Approach. *Chemosphere* **2017**, *183*, 410–418.
- (54) Hoover, K. R.; Acree, W. E.; Abraham, M. H. Chemical Toxicity Correlations for Several Fish Species Based on the Abraham Solvation Parameter Model. *Chem. Res. Toxicol.* **2005**, *18* (9), 1497–1505.
- (55) United States Environmental Protection Agency. Ecological Structure Activity Relationships (ECOSAR) Predictive Model. 2017.
- (56) Faure, F.; Demars, C.; Wieser, O.; Kunz, M.; De Alencastro, L. F. Plastic Pollution in Swiss Surface Waters: Nature and Concentrations, Interaction with Pollutants. *Environ. Chem.* **2015**, *12* (5), 582–591.