

# *Vibrio syngnathi* sp. nov., a fish pathogen, isolated from the Kiel Fjord

Cynthia Maria Chibani<sup>1,2</sup>, Robert Hertel<sup>1</sup>, Meina Neumann-Schaal<sup>3</sup>, Henry Goehlich<sup>4,5</sup>, Kim Wagner<sup>4,5</sup>, Boyke Bunk<sup>3</sup>, Cathrin Spröer<sup>3</sup>, Jörg Overmann<sup>3</sup>, Michael Hoppert<sup>1</sup>, Silke Mareike Marten<sup>4,5</sup>, Olivia Roth<sup>4,5</sup>, Heiko Liesegang<sup>1</sup> and Carolin C. Wendling<sup>4,6,\*</sup>

## Abstract

A new *Vibrio* strain, K08M4<sup>T</sup>, was isolated from the broad-nosed pipefish *Syngnathus typhle* in the Kiel Fjord. Infection experiments revealed that K08M4<sup>T</sup> was highly virulent for juvenile pipefish. Cells of strain K08M4<sup>T</sup> were Gram-stain-negative, curved rod-shaped and motile by means of a single polar flagellum. The strain grew aerobically at 9–40°C, at pH 4–10.5 and it tolerated up to 12% (w/v) NaCl. The most prevalent (>10%) cellular fatty acids of K08M4<sup>T</sup> were C<sub>16:1</sub> ω7c and C<sub>16:0</sub>. Whole-genome comparisons revealed that K08M4<sup>T</sup> represents a separate evolutionary lineage that is distinct from other *Vibrio* species and falls within the *Splendidus* clade. The genome is 4,886,292 bp in size, consists of two circular chromosomes (3,298,328 and 1,587,964 bp) and comprises 4,178 protein-coding genes and 175 RNA genes. In this study, we describe the phenotypic features of the new isolate and present the annotation and analysis of its complete genome sequence. Based on these data, the new isolate represents a new species for which we propose the name *Vibrio syngnathi* sp. nov. The type strain is K08M4<sup>T</sup> (=DSM 109818<sup>T</sup>=CECT 30086<sup>T</sup>).

## INTRODUCTION

The genus *Vibrio* comprises 137 validly published species (<https://lpsn.dsmz.de/genus/vibrio>) which have been grouped into 16 monophyletic clades [1, 2]. The *Splendidus* clade, which contains 17 closely related *Vibrio* species, constitutes the largest clade within the genus *Vibrio* [3]. The phenotypic variability of additional species that did not fit exactly the description of *Vibrio splendidus* led to the proposal of the term *Vibrio splendidus*-like, which is now widely used for isolates that cannot unequivocally be assigned to the species *V. splendidus*.

Bacteria of the *Splendidus* clade are dominant members of the coastal bacterioplankton, in sediments and various marine organisms, for a review see [4]. Various *V. splendidus*-like strains have been found to cause vibriosis. Vibriosis is one of the most prevalent diseases affecting marine life and it accounts for major economic losses in aquaculture [4, 5]. Vibriosis causes high rates of mortality in aquaculture animals such as turbot, scallops, clams and oysters [6–9], and can even spread to humans through the consumption of infected seafood [10, 11]. Virulence of these strains is multifactorial and regulated by genetic determinants, e.g. prophages [12] or pathogenicity islands [13] as well as extrinsic factors, such as temperature [14] or salinity [15]. Furthermore, a recent epidemiological survey of the *Splendidus* clade suggests an unexplored diversity of virulence factors and a massive lateral transfer of virulence genes within this clade [16].

Despite a significant body of empirical research, our understanding of the factors that contribute to the virulence of *V. splendidus*-like strains is far from being complete. Whole-genome sequencing can provide further insights into potential virulence mechanisms

**Author affiliations:** <sup>1</sup>Georg-August University Goettingen, Institute of Microbiology and Genetics, 37077 Goettingen, Germany; <sup>2</sup>Institute for General Microbiology, University of Kiel, 24118 Kiel, Germany; <sup>3</sup>Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures, Inhoffenstr. 7B, 38124 Braunschweig, Germany; <sup>4</sup>GEOMAR, Helmholtz Centre for Ocean Research, Marine Evolutionary Ecology, Düsternbrooker Weg 20, 24105 Kiel, Germany; <sup>5</sup>Marine Evolutionary Biology, Zoological Institute, Kiel University, 24118 Kiel, Germany; <sup>6</sup>ETH Zürich, Institute of Integrative Biology, Universitätsstraße 16, 8092 Zürich, Switzerland.

\*Correspondence: Carolin C. Wendling, carolin.wendling@env.ethz.ch

**Keywords:** *Splendidus* clade; genomic comparison; fish pathogen; virulence factors; *Vibrio*; complete genome sequence.

**Abbreviations:** ANI, average nucleotide identity; dDDH, digital DNA–DNA hybridization; TCBS, thiosulphate–citrate–bile–sucrose; TYGS, Type Strain Genome Server.

The complete genome sequence is available at NCBI GenBank under the accession numbers CP017916 and CP017917. The 16S rRNA sequence can be found at NCBI GenBank under the accession number OP359305. The raw reads can be found under BioProject accession number PRJNA345286. One supplementary figure and six supplementary tables are available with the online version of this article.

005951 © 2023 The Authors



of this pathogen. At the time of writing, 819 complete *Vibrio* genome sequences and 13 *V. splendidus*-like strains were publicly available (NCBI, date of access 22 May 2022).

In this study, we sequenced and annotated the complete genome of strain K08M4<sup>T</sup>, which was found to represent a new *Vibrio* species, *Vibrio syngnathi* sp. nov. We follow the proposed minimal standards for the use of genome sequence data for taxonomic purposes [17] and provide first insight into the genomic basis of its pathogenicity.

## ORIGIN AND ISOLATION

Strain K08M4<sup>T</sup> was isolated during a previous study from the intestine of a healthy adult pipefish collected in the Kiel Fjord in 2010 (for a detailed description see [18]) and was shown to be pathogenic to juvenile pipefish in laboratory experiments. Based on a multi-locus sequencing approach, K08M4<sup>T</sup> was initially affiliated to the *Splendidus* clade [19]. The whole-genome comparison in the present study supports the consideration that K08M4<sup>T</sup> represents a distinct species within the *Splendidus* clade.


## PHENOTYPIC CHARACTERIZATION

K08M4<sup>T</sup> was routinely cultivated on *Vibrio*-selective thiosulphate–citrate–bile–sucrose (TCBS) agar plates or in liquid culture using medium 101 (medium 101: 0.5 % (w/v) peptone, 0.3 % (w/v) meat extract, 1.5 % (w/v) NaCl in MilliQ water) at 25 °C.

Cell morphology of strain K08M4<sup>T</sup> was determined by electron microscopy of negatively stained cells using a JEOL 1011 transmission electron microscope (Eching, Munich, Germany) as described by Chibani *et al.* [20]. Briefly, cells were grown in liquid medium 101 for 12 h overnight at 25 °C with constant shaking at 230 r.p.m. as pre-culture. The main culture was inoculated with 1.5% v/v grown for 3 h and adsorbed onto Formvar-carbon grids (Plano, Wetzlar, Germany) prior to staining. Phosphotungstic acid, dissolved in pure water (1 % resp 0.5% w/v final concentration) and adjusted to pH 7 with sodium hydroxide served as a staining solution. Images were captured using a Gatan Orius SC 1000A camera and processed with the Gatan Digital Micrograph software package (version 1.84.1177). Cells of strain K08M4<sup>T</sup> were slightly curved (1.5–2.5 µm long and 0.5–1 µm wide) with a monotrichous flagellum (Fig. 1).

Salinity and temperature ranges were determined by generating growth curves over 24 h time intervals of 1 : 100 dilutions of overnight cultures using a microplate reader (Tecan Infinite M200) at different NaCl concentrations ranging from 0 to 15 % (w/v) at 25 °C with 0.5% intervals, as well as across a temperature gradient of 4–45 °C with 5 °C intervals at 1.5% (w/v) NaCl and different pH levels ranging from pH 3.5 to 11 at intervals of 1.0 pH unit. Strain K08M4<sup>T</sup> produced yellow-coloured, round colonies on TCBS agar plates and cream-coloured, round colonies on medium 101 plates after incubation for 24 h at 25 °C. The strain did not grow below 9 °C and showed optimum growth between 25 and 35 °C at 0.4–12% NaCl, with optimum growth occurring at 1.5% NaCl and between pH 4 and 10.5.

We performed a comparative phenotypic analysis between strain K08M4<sup>T</sup> and three closely related species obtained from the German Collection of Microorganisms and Cell Cultures DSMZ: *Vibrio tasmaniensis* DSM 17182<sup>T</sup>, *Vibrio splendidus* DSM 19640<sup>T</sup> and *Vibrio splendidus* DSM 26178<sup>T</sup>. The Gram reaction was determined by KOH and aminopeptidase using the Bactident Aminopeptidase (Merck, 113301) tests [21]. Oxidase activity was tested by the method of Kovacs *et al.* [22]. Catalase tests were performed by mixing freshly grown bacterial cells with 10 % H<sub>2</sub>O<sub>2</sub>, followed by examining gas bubble formation. Chitinase activity was confirmed using a chitinase assay kit (Sigma Aldrich, CS0980). Activities of constitutive enzymes and other physiological properties were determined using API 20E, API ZYM and API 50CH (bioMérieux) kits according to the instructions of the manufacturer. Susceptibility to the vibriostatic agent 2,4-diamino-6,7-diisopropyl pteridine phosphate salt (O/129; Sigma) was



**Fig. 1.** Left: broad-nosed pipefish, *Syngnathus typhe* (by Uli Kunz), from which K08M4<sup>T</sup> was isolated from; right: transmission electron microscope image of K08M4<sup>T</sup> after 3 h of growth in medium 101.

**Table 1.** Differential characteristics between strain K08M4<sup>T</sup> and the reference strains of some phylogenetically related species

Strains: 1, DSM 109818<sup>T</sup> (this study); 2, *Vibrio tasmaniensis* DSM 17182<sup>T</sup> (this study); 3, *Vibrio splendidus* DSM 19640<sup>T</sup> (this study); 4, *Vibrio splendidus* DSM 26178<sup>T</sup> (this study); 5, *Vibrio pomeroyi* LMG 20537<sup>T</sup> [43]; 6, *Vibrio gallaecicus* LMG 24045<sup>T</sup> [44, 45]; 7, *Vibrio cholerae* ATCC 14035<sup>T</sup> [46]. +, Positive reaction; -, negative reaction; ±, variable outcomes; ND, no data available. A full table containing all results from API ZYM, API 20E and API 50CHE assays can be found in Tables S1–S3.

Characteristic	1	2	3	4	5	6	7
Arginin dihydrolase	-	+	+	+	+	-	-
Voges-Proskauer test	-	-	-	-	-	-	±
ONPG β-Galactosidase	-	-	+	+	+	-	ND
O/129 susceptibility	+	-	+	-	+	+	-
Gelatin hydrolysis	-	-	+	+	+	+	ND
Acid from:							
Galactose	-	-	+	+	ND	ND	+
Mannitol	-	+	+	+	+	+	+
Amygdalin	+	+	+	+	+	-	ND
Melibiose	-	-	+	+	-	-	-
Sucrose	+	-	-	-	+	-	+
2-Ketogluconate	-	-	+	-	ND	ND	-
Mannose	-	+	+	+	ND	ND	+

determined by means of discs containing 150 µg O/129. Fatty acid analysis was performed using the standard protocol of the Sherlock Microbial Identification (system package version 6.4 with the TSBA6.0 database), according to the technical instructions provided by the manufacturer [23]. Combined analysis by gas chromatography coupled to a mass spectrometer was used to confirm the identity of the fatty acids based on retention time and mass spectral data [24]. Major characteristics and unique phenotypic characteristics differentiating K08M4<sup>T</sup> from its close relatives *V. tasmaniensis* DSM 17182<sup>T</sup>, *V. splendidus* DSM 19640<sup>T</sup> and *V. splendidus* DSM 26178<sup>T</sup> are summarized in Table 1 and full results are given in Table S1–S3. The cellular fatty acid profile of K08M4<sup>T</sup> in comparison to *V. tasmaniensis* DSM 17182<sup>T</sup> and *V. splendidus* DSM 19640<sup>T</sup> is listed in Table 2, full results for strain K08M4<sup>T</sup> only are given in Table S4.

**Table 2.** Fatty acid profile of strain K08M4<sup>T</sup> (=DSM 109818<sup>T</sup>) in comparison with *Vibrio tasmaniensis* DSM 17182<sup>T</sup> and *Vibrio splendidus* DSM 19640<sup>T</sup>

All data in the table are from the present study. Fatty acids present in amounts lower than 1 % are not shown. Quantification by GC-FID, identity was confirmed by mass spectrometry. A full table can be found in Table S4.

Fatty acid	K08M4 <sup>T</sup>	<i>V. tasmaniensis</i> DSM 17182 <sup>T</sup>	<i>V. splendidus</i> DSM 19640 <sup>T</sup>
C <sub>12:0</sub>	3.9%	4.0%	5.0%
C <sub>12:0</sub> 3OH	2.7%	2.4%	2.5%
C <sub>14:0</sub>	3.8%	4.1%	5.7%
C <sub>14:0</sub> 3OH	1.8%	1.7%	1.9%
C <sub>16:0</sub> iso	7.6%	3.4%	3.2%
C <sub>16:1</sub> ω7c	39.9%	43.8%	40.7%
C <sub>16:1</sub> ω7t	3.3%	4.3%	5.3%
C <sub>16:0</sub>	19.2%	21.6%	19.4%
C <sub>18:1</sub> ω7c	8.7%	7.0%	11.1%
C <sub>18:0</sub>	1.0%	1.1%	0.5%


## GENOMIC CHARACTERIZATION

The Qiagen Genomic Tip/100 G was used to extract high molecular weight DNA from cells of strain K08M4<sup>T</sup>. The extracted DNA was then used to generate a SMRTbell™ template library, according to the instructions from Pacific Biosciences, Menlo Park, CA, USA, following the Procedure & Checklist - 10 kb Template Preparation Using BluePippin™ Size-Selection System.

SMRT genome sequencing was carried out on the PacBio RSII (Pacific Biosciences, Menlo Park, CA, USA), taking one 240 min movie for a single SMRT cell using P6 chemistry at the Leibniz Institute DSMZ-German Collection of Microorganisms and Cell Cultures in Braunschweig, Germany. Genome finishing and annotation were performed at the Institute for Microbiology and Genetics at the Georg-August University of Göttingen, Germany.

We obtained a total of 64,464 post-filtered reads with a mean read length of 13,396 bp representing a genome coverage of 141.45-fold. Genome assembly was performed with the RS\_HGAP\_Assembly.3 protocol included in SMRT Portal version 2.3.0. The assembly resulted in two large contigs that could be associated with chromosomes 1 and 2. The genome has a size of 4,886,292 bp (chromosome 1, 3.2 Mb; chromosome 2, 1.5 Mb) and the G+C content of strain K08M4 is 44.06 mol%. The completeness and contamination levels of strain K08M4<sup>T</sup> were computed using CheckM [25]. Based on the assembly of the genome without gaps or ambiguities, the completeness level (99.69%) and the contamination level (1.69%) of both chromosomes, a metagenome assembled genome would be labelled as 'Finished' [26]. The genome project has been deposited at the European Nucleotide Archive under the ID number 345286 and accession number PRJNA345286. The accession number for chromosome 1 is CP017916 and for chromosome 2 is CP017917. The raw reads were deposited under the BioProject accession number PRJNA345286.

Automatic annotation, gene prediction, and rRNA and tRNA gene identification were carried out with Prokka (version 1.11) [27]. Functional annotations were done by searching against InterPro [28] and COG databases [29]. A total of 4,353 genes were predicted, of which 4178 encode proteins, 175 encode rRNAs, 43 encode tRNAs and at least 46 encode ncRNAs. Twelve



**Fig. 2.** Phylogenetic tree inferred with FastME 2.1.6.1 [34] from GBDP distances calculated from genome sequences. The branch lengths are scaled in terms of GBDP distance formula  $d_s$ . The numbers above branches are GBDP pseudo-bootstrap support values >60 % from 100 replications, with an average branch support of 97.7 %. The tree was rooted at the midpoint [35].

pseudogenes were found, with eight located on chromosome 1 and four located on chromosome 2. Of the predicted CDS, a functional prediction was made for 77.1 %. A COG prediction could be retrieved for 70.0%, with the remaining annotated as hypothetical proteins. We found a total of 148 insertion elements: 72 on chromosome 1 and 76 on chromosome 2. A putative 48.2 kb intact prophage was identified on chromosome 2 using PHASTER [30] and this region is predicted to encode 45 proteins, of which most are phage-related, whereas 23 are hypothetical or uncharacterized. In addition, one incomplete prophage was found on chromosome 2.

Whole genome based taxonomic analysis was done using the Type (Strain) Genome Server (TYGS) [31], a free bioinformatics platform provided by the DSMZ (<https://tygs.dsmz.de>) following the standard pipeline including recent updates [32]. The 11 closest type strain genomes were determined following the standard pipeline.

For phylogenomic inference, all pairwise comparisons among the set of genomes were conducted using GBDP and accurate intergenomic distances inferred under the algorithm 'trimming' and distance formula  $d_5$  [33]. 100 distance replicates were calculated each. Digital DNA–DNA hybridization (dDDH) values and confidence intervals were calculated using the recommended settings of the Genome-to-Genome Calculator (GGDC) 3.0.

The resulting intergenomic distances were then used to infer a balanced minimum-evolution tree with branch support via FastME 2.1.6.1 including SPR postprocessing [34]. Branch support was inferred from 100 pseudo-bootstrap replicates each. The tree was rooted at the midpoint [35] and visualized using FigTree version 1.4.2 (<http://tree.bio.ed.ac.uk/software/figtree/>).

The type-based species clustering using a 70 % dDDH radius around each of the 11 type strains was done as previously described [31]. Subspecies clustering was done using a 79 % dDDH threshold as previously introduced [36]. The clustering yielded 12 species clusters and the provided query strains were assigned to one of these, K08M4<sup>T</sup> was located in one of 12 subspecies clusters (Fig. 2). The dDDH values between the complete genome sequence of strain K08M4<sup>T</sup> and the genome sequences of the 11 most closely related species from the *Splendidus* clade are significantly below the species delimitation cut-off of 70 % (Table 3).

**Table 3.** Genome-to-genome distance calculation for strain K08M4<sup>T</sup> using the DSMZ GGDC server


Distances have been calculated using formula 4, for details see [32, 33].

Species	DDH	Model CI	Distance	G+C difference (mol%)	Accession No. Chr. 1
<i>Vibrio synnathi</i> K08M4 <sup>T</sup>	–	–	–	44.06	CP017916.1
<i>Vibrio tasmaniensis</i> LMG 20012 <sup>T</sup>	58.9	[56.0–61.6%]	0.0536	0.03	NZ_AP025510.1
<i>Vibrio atlanticus</i> LGP32	54.0	[51.3–56.7%]	0.0629	0.15	NC_011753.2
<i>Vibrio atlanticus</i> CECT 7223 <sup>T</sup>	54.1	[51.4–56.8%]	0.0628	0.09	NZ_AP025460.1
<i>Vibrio cyclitrophicus</i> ED008	39.2	[36.8–41.8%]	0.1020	0.12	NZ_CP064172.1
<i>Vibrio crassostreae</i> LMG 22240 <sup>T</sup>	37.8	[35.4–40.3%]	0.1071	0.44	NZ_AP025476.1
<i>Vibrio gigantis</i> ACE001	35.8	[33.4–38.3%]	0.1149	0.23	NZ_CP092384.1
<i>Vibrio crassostreae</i> ED395	39.4	[36.9–41.9%]	0.1015	0.31	NZ_CP064170.1
<i>Vibrio cyclitrophicus</i> ED287	39.3	[36.8–41.8%]	0.1018	0.01	NZ_CP065366.1
<i>Vibrio gigantis</i> LMG 22741 <sup>T</sup>	35.7	[33.3–38.3%]	0.1151	0.25	NZ_AP025492.1
<i>Vibrio splendidus</i> LMG 19031 <sup>T</sup>	43.6	[41.1–46.2%]	0.0881	0.15	NZ_AP025508.1
<i>Vibrio pomeroyi</i> LMG 20537 <sup>T</sup>	34.9	[32.5–37.4%]	0.1185	0.46	NZ_AP025506.1
<i>Vibrio lentus</i> LMG 21034 <sup>T</sup>	38.6	[36.1–41.1%]	0.1043	0.09	NZ_AP025499.1
<i>Vibrio chagasii</i> LMG 21353 <sup>T</sup>	30.4	[28.0–32.9%]	0.1402	0.41	NZ_AP025465.1
<i>Vibrio artabrorum</i> CECT 7226 <sup>T</sup>	31.8	[29.4–34.3%]	0.1327	0.16	NZ_AP025458.1
<i>Vibrio gallaecicus</i> CECT 7244 <sup>T</sup>	24.4	[22.1–26.8%]	0.1790	2.44	NZ_AP025490.1
<i>Vibrio cholerae</i> El Tor N16961	20.9	[18.7–23.3%]	0.2103	3.50	NC_002505
<i>Vibrio celticus</i> Rd 8.15 <sup>T</sup>	38.5	[36.0–41.0%]	0.1047	0.48	NZ_MVJF01000005.1

In addition, phylogenetic analysis based on a multilocus sequence analysis tree using eight housekeeping genes (*ftsZ*, *gapA*, *gyrB*, *mreB*, *pyrH*, *recA*, *rpoA* and *tpoA*) confirmed that strain K08M4<sup>T</sup> forms a monophyletic group within the *Splendidus* clade (Fig. S1, available in the online version of this article).

Furthermore, we compared the genomic relatedness based on the average nucleotide identity (ANI) between strain K08M4<sup>T</sup> and all closed *Vibrio* sequences as reference available on NCBI (date of access 17.11.2022) using FastANI (<https://github.com/ParBLISS/FastANI> [37]). All closed *Vibrio* genomes were downloaded using the NCBI Datasets command line tools and the following command line 'datasets download genome taxon 662 --complete'. Firstly, the ANI values between the complete genome sequence of strain K08M4<sup>T</sup> were computed separately for chromosomes 1 and 2, as well as for the concatenated sequences of chromosomes 1 and 2 of all deduplicated closed *Vibrio* genomes ( $n=532$ ). ANI between the K08M4<sup>T</sup> comparisons and reference strains that were above 90 % from the concatenated sequences were selected for the purpose of visualization (Fig. 3) using PyANI [38] and the ANIb option (<https://github.com/widowquinn/pyani>). Most ANI values in any combination were below the threshold of 95%, the repeatedly established species demarcation cut-off value [37]. One exception is nucleotide identity between strain K08M4<sup>T</sup> and *V. tasmaniensis* LMG 20012 (ANI chromosome 1, 95.5%; chromosome 2, 93.96%; both chromosomes together, 95.03%).

Phylogenomic treeing and taxonomy assignment of K08M4<sup>T</sup> was additionally performed based on 120 bacterial marker genes with GTDB-Tk version 2.1.0 [39, 40] (database release R207) using the 'classify\_wf' function and default parameters. Genomes are assigned at the species level if the ANI to the closest GTDB species representative genome was  $\geq 95\%$  and the Alignment Fraction (AF) was  $\geq 30\%$ . Accordingly, K08M4<sup>T</sup> was characterized as *V. tasmaniensis* based on GTDB taxonomy. However, taking all sequence-based results (ANI of K08M4<sup>T</sup> and *V. tasmaniensis* LMG 20012 is 95.03%, directly at the



**Fig. 3.** Average nucleotide identity (ANI) analysis of strain K08M4<sup>T</sup> sequenced strain with available closed *Vibrio* genomes (ANI >90%) as references. ANI analysis based on BLAST alignment of the genome sequences was performed and visualized using PyANI.

boundary of the species demarcation cut-off) together with experimental confirmation, we can confidently confirm that K08M4<sup>T</sup> represents a novel *Vibrio* species.

## PATHOGENICITY

We performed a controlled infection experiment to test the virulence of strain K08M4<sup>T</sup> on pipefish larvae (a detailed description of the methods and the statistical analysis can be found in [19]). Briefly, we exposed 10–11 juvenile pipefish in triplicates to either 10<sup>9</sup> c.f.u. ml<sup>-1</sup> of strain K08M4<sup>T</sup>, 10<sup>9</sup> c.f.u. ml<sup>-1</sup> of a non-virulent *V. alginolyticus* K10K4, or seawater as control. Fish mortality was recorded daily. On day three, we sampled one fish per experimental unit and analysed the expression of 32 immune genes relative to two housekeeping genes using a Fluidigm BioMark as described in [41]. This included 17 target genes assigned to the innate immune system, two target genes assigned to the complement component system and 13 target genes assigned to the adaptive immune system (Table S5). Details about function of genes, sequences and primer design can be found in [41].

Strain K08M4<sup>T</sup> causes high rates of mortality (Fig. 4a) in pipefish larvae and elicits a different expression profile of selected immune genes compared to the non-virulent *V. alginolyticus* strain K10K4 or pipefish exposed to seawater (Fig. 4b).


Several genetically encoded virulence factors such as iron transport systems, flagellum/ motility, hemolysins, proteases, lipopolysaccharides, exopolysaccharides, repeats in toxins (RTX), outer membrane proteins and a type IV pilus could be identified (Table S6). Compared to other *Vibrio* species isolated from the Kiel Fjord, K08M4<sup>T</sup> has a unique virulence profile [42].

Based on the results of comparative genomic analysis and its phenotypic description, strain K08M4<sup>T</sup> represents a novel species of the genus *Vibrio*, with an increased virulence profile for juvenile pipefish, for which we propose the name *Vibrio syngnathi* sp. nov.

## DESCRIPTION OF *VIBRIO SYNGNATHI* SP. NOV.

*Vibrio syngnathi* (syn.gna'thi. N.L. gen. n. *syngnathi*, pertaining to the broad-nosed pipefish *Syngnathus typhle* from which the strain was originally isolated).

Cells are slightly curved rods, Gram-stain-negative, curved rod-shaped and motile by means of a single polar flagellum. Colonies are circular, yellow and 2–4 mm in size after 24 h growth on TCBS agar at 25 °C. Cells can grow at 9–40 °C and pH 4–10.5. Optimal growth is observed at 25–35 °C. Growth occurs in the presence of 0.4–12% (w/v) NaCl. Biochemical characteristics of K08M4<sup>T</sup> include positive results for oxidase, catalase, chitinase, indole production and O1/129 sensitivity. Acid is produced from glycerol, ribose, glucose, fructose, mannose, *N*-acetylglucosamine, aesculin, salicin, cellobiose, maltose, sucrose, trehalose, starch, glycogen and gluconate. K08M4<sup>T</sup> is negative for β-galactosidase, arginine dihydrolase, lysine decarboxylase, ornithine decarboxylase, citrate utilization, H<sub>2</sub>S production, urease, tryptophan deaminase, Voges–Proskauer test and gelatin hydrolysis.



**Fig. 4.** Pipefish mortality in per cent (a) and ordination of differentially expressed immune genes between K08M4, K10K4 or seawater-injected pipefish (SW).

The type strain can grow on glycerol, starch and glycogen. The major fatty acids (>10%) are C<sub>16:1</sub> ω7c and C<sub>16:0</sub>. K08M4<sup>T</sup> causes mortality in juvenile pipefish, *S. typhle*.

The type strain, K08M4<sup>T</sup> (=DSM 109818<sup>T</sup>=CECT 30086<sup>T</sup>), was isolated from the whole intestines of an adult broad-nosed pipefish *Syngnathus typhle*, caught in the Kiel Fjord, Germany.

#### Funding information

This study was supported by the Deutsche Forschungsgemeinschaft (DFG) SPP 1819 on rapid adaptation (WE5822/1-1, WE5822/1-2, and OR4628/4-2 given to C.C.W. and O.R.) and a Postdoctoral fellowship from the Cluster of Excellence 'The Future Ocean' given to C.C.W.

#### Acknowledgements

We are grateful to Anja Frühling, Gesa Martens, Anika Wasner, Bettina Ehlert, Simone Schrader and Nicole Heyer for excellent technical assistance.

#### Author contributions

O.R. isolated the strain. C.C.W., O.R. and H.L. designed the study. C.C.W., R.H., H.L., B.B., C.S. and J.O. were involved in the preparation of bacterial DNA and genome sequencing. C.C.W. performed infection experiments. H.G., K.S.W., M.M. and M.N.S. performed microbiological lab work. C.C., H.L. and B.B. analysed the bacteria genomes. M.H. performed electron microscopy. C.C.W. coordinated the project. C.C.W. and C.C. wrote the first draft of the manuscript. All authors approved the final version of the manuscript.

#### Conflicts of interest

The authors declare that there are no conflicts of interest.

#### Ethical statement

Approval for using pipefish during infection experiments was given by the Ministerium für Landwirtschaft, Umwelt und ländliche Räume des Landes Schleswig-Holstein V312-7224.121-19 (65-5/13).

#### References

- Sawabe T, Kita-Tsukamoto K, Thompson FL. Inferring the evolutionary history of vibrios by means of multilocus sequence analysis. *J Bacteriol* 2007;189:7932–7936.
- Sawabe T, Ogura Y, Matsumura Y, Feng G, Amin AR, et al. Updating the *Vibrio* clades defined by multilocus sequence phylogeny: proposal of eight new clades, and the description of *Vibrio tritonius* sp. nov. *Front Microbiol* 2013;4:414.
- Pérez-Cataluña A, Lucena T, Tarazona E, Arahall DR, Macián MC, et al. An MLSA approach for the taxonomic update of the *Splendidus* clade, a lineage containing several fish and shellfish pathogenic *Vibrio* spp. *Syst Appl Microbiol* 2016;39:361–369.
- Egidius E. Vibriosis: pathogenicity and pathology. A review. *Aquaculture* 1987;67:15–28.
- Ina-Salwany MY, Al-Saari N, Mohamad A, Mursidi F-A, Mohd-Aris A, et al. Vibriosis in fish: a review on disease development and prevention. *J Aquat Anim Health* 2019;31:3–22.
- Sugumar G, Nakai T, Hirata Y, Matsubara D, Muroga K. *Vibrio splendidus* biovar II as the causative agent of bacillary necrosis of Japanese oyster *Crassostrea gigas* larvae. *Dis Aquat Organ* 1998;33:111–118.
- Lacoste A, Jalabert F, Malham S, Cueff A, Gélébart F, et al. A *Vibrio splendidus* strain is associated with summer mortality of juvenile oysters *Crassostrea gigas* in the Bay of Morlaix (North Brittany, France). *Dis Aquat Organ* 2001;46:139–145.
- Gómez-León J, Villamil L, Lemos ML, Novoa B, Figueras A. Isolation of *Vibrio alginolyticus* and *Vibrio splendidus* from aquacultured carpet shell clam (*Ruditapes decussatus*) larvae associated with mass mortalities. *Appl Environ Microbiol* 2005;71:98–104.
- Balcázar JL, Gallo-Bueno A, Planas M, Pintado J. Isolation of *Vibrio alginolyticus* and *Vibrio splendidus* from captive-bred seahorses with disease symptoms. *Antonie van Leeuwenhoek* 2010;97:207–210.
- Baffone W, Pianetti A, Bruscolini F, Barbieri E, Citterio B. Occurrence and expression of virulence-related properties of *Vibrio* species isolated from widely consumed seafood products. *Int J Food Microbiol* 2000;54:9–18.
- Su YC, Liu C. *Vibrio parahaemolyticus*: a concern of seafood safety. *Food Microbiol* 2007;24:549–558.
- Weinbauer MG, Brettar I, Höfle MG. Lysogeny and virus-induced mortality of bacterioplankton in surface, deep, and anoxic marine waters. *Limnol Oceanogr* 2003;48:1457–1465.
- O'Shea YA, Boyd EF. Mobilization of the *Vibrio* pathogenicity island between *Vibrio cholerae* isolates mediated by CP-T1 generalized transduction. *FEMS Microbiol Lett* 2002;214:153–157.
- Wendling CC, Batista FM, Wegner KM. Persistence, seasonal dynamics and pathogenic potential of *Vibrio* communities from Pacific oyster hemolymph. *PLoS One* 2014;9:e94256.
- Poirier M, Listmann L, Roth O. Selection by higher-order effects of salinity and bacteria on early life-stages of Western Baltic spring-spawning herring. *Evol Appl* 2017;10:603–615.
- Nasfi H, Travers MA, de Lorgeril J, Habib C, Sannie T, et al. A European epidemiological survey of *Vibrio splendidus* clade shows unexplored diversity and massive exchange of virulence factors. *World J Microbiol Biotechnol* 2015;31:461–475.
- Chun J, Oren A, Ventosa A, Christensen H, Arahall DR, et al. Proposed minimal standards for the use of genome data for the taxonomy of prokaryotes. *Int J Syst Evol Microbiol* 2018;68:461–466.
- Roth O, Keller I, Landis SH, Salzburger W, Reusch TBH. Hosts are ahead in a marine host-parasite coevolutionary arms race: innate immune system adaptation in pipefish *Syngnathus typhle* against *Vibrio* phylotypes. *Evolution* 2012;66:2528–2539.
- Wendling CC, Pieczyk A, Refardt D, Chibani C, Hertel R, et al. Tripartite species interaction: eukaryotic hosts suffer more from phage susceptible than from phage resistant bacteria. *BMC Evol Biol* 2017;17:98.
- Chibani CM, Hertel R, Hoppert M, Liesegang H, Wendling CC. Closely related *Vibrio alginolyticus* strains encode an identical repertoire of caudovirales-like regions and filamentous phages. *Viruses* 2020;12:1359.
- Cerny G. Method for the distinction of gramnegative from grampositive bacteria. *European J Appl Microbiol* 1976;3:223–225.
- Kovacs N. Identification of *Pseudomonas pyocyanea* by the oxidase reaction. *Nature* 1956;178:703–.
- Sasser M. *Technical Note 101: Identification of Bacteria by Gas Chromatography of Cellular Fatty Acids*. Newark, DE: MIDI; 1990.
- Vieira S, Huber KJ, Neumann-Schaal M, Geppert A, Luckner M, et al. *Usitatibacter rugosus* gen. nov., sp. nov. and *Usitatibacter palustris* sp. nov., novel members of *Usitatibacteraceae* fam. nov. within the order Nitrosomonadales isolated from soil. *Int J Syst Evol Microbiol* 2021;71.



25. Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: assessing the quality of microbial genomes recovered from isolates, single cells, and metagenomes. *Genome Res* 2015;25:1043–1055.
26. Bowers RM, Kyrpides NC, Stepanauskas R, Harmon-Smith M, Doud D, et al. Minimum information about a single amplified genome (MISAG) and a metagenome-assembled genome (MIMAG) of bacteria and archaea. *Nat Biotechnol* 2017;35:725–731.
27. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* 2014;30:2068–2069.
28. Blum M, Chang H-Y, Chuguransky S, Grego T, Kandasamy S, et al. The InterPro protein families and domains database: 20 years on. *Nucleic Acids Res* 2021;49:D344–D354.
29. Galperin MY, Wolf YI, Makarova KS, Vera Alvarez R, Landsman D, et al. COG database update: focus on microbial diversity, model organisms, and widespread pathogens. *Nucleic Acids Res* 2021;49:D274–D281.
30. Arndt D, Grant JR, Marcu A, Sajed T, Pon A, et al. PHASTER: a better, faster version of the PAST phage search tool. *Nucleic Acids Res* 2016;44:W16–21.
31. Meier-Kolthoff JP, Göker M. TYGS is an automated high-throughput platform for state-of-the-art genome-based taxonomy. *Nat Commun* 2019;10:2182.
32. Meier-Kolthoff JP, Carbasse JS, Peinado-Olarte RL, Göker M. TYGS and LPSN: a database tandem for fast and reliable genome-based classification and nomenclature of prokaryotes. *Nucleic Acids Res* 2022;50:D801–D807.
33. Meier-Kolthoff JP, Auch AF, Klenk H-P, Göker M. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 2013;14:60.
34. Lefort V, Desper R, Gascuel O. FastME 2.0: a comprehensive, accurate, and fast distance-based phylogeny inference program. *Mol Biol Evol* 2015;32:2798–2800.
35. Farris JS. Estimating phylogenetic trees from distance matrices. *Am Nat* 1972;106:645–668.
36. Meier-Kolthoff JP, Hahnke RL, Petersen J, Scheuner C, Michael V, et al. Complete genome sequence of DSM 30083(T), the type strain (U5/41(T)) of *Escherichia coli*, and a proposal for delineating subspecies in microbial taxonomy. *Stand Genomic Sci* 2014;9:2.
37. Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI analysis of 90K prokaryotic genomes reveals clear species boundaries. *Nat Commun* 2018;9:5114.
38. Pritchard L, Glover RH, Humphris S, Elphinstone JG, Toth IK. Genomics and taxonomy in diagnostics for food security: soft-rotting enterobacterial plant pathogens. *Anal Methods* 2016;8:12–24.
39. Chaumeil PA, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk v2: memory friendly classification with the genome taxonomy database. *Bioinformatics* 2022;38:5315–5316.
40. Parks DH, Chuvochina M, Rinke C, Mussig AJ, Chaumeil P-A, et al. GTDB: an ongoing census of bacterial and archaeal diversity through a phylogenetically consistent, rank normalized and complete genome-based taxonomy. *Nucleic Acids Res* 2022;50:D785–D794.
41. Beemelmanns A, Roth O. Biparental immune priming in the pipefish *Syngnathus typhle*. *Zoology* 2016;119:262–272.
42. Chibani CM, Roth O, Liesegang H, Wendling CC. Genomic variation among closely related *Vibrio alginolyticus* strains is located on mobile genetic elements. *BMC Genomics* 2020;21:354.
43. Thompson FL, Thompson CC, Li Y, Gomez-Gil B, Vandenberghe J, et al. *Vibrio kanaloae* sp. nov., *Vibrio pomeroyi* sp. nov. and *Vibrio chagasii* sp. nov., from sea water and marine animals. *Int J Syst Evol Microbiol* 2003;53:753–759.
44. Beaz-Hidalgo R, Doce A, Pascual J, Toranzo AE, Romalde JL. *Vibrio gallaecicus* sp. nov. isolated from cultured clams in north-western Spain. *Syst Appl Microbiol* 2009;32:111–117.
45. Wang H, Liu J, Wang Y, Zhang XH. *Vibrio marisflavi* sp. nov., isolated from seawater. *Int J Syst Evol Microbiol* 2011;61:568–573.
46. Colwell RR. Polyphasic taxonomy of the genus *Vibrio*: numerical taxonomy of *Vibrio cholerae*, *Vibrio parahaemolyticus*, and related *Vibrio* species. *J Bacteriol* 1970;104:410–433.

### Five reasons to publish your next article with a Microbiology Society journal

1. When you submit to our journals, you are supporting Society activities for your community.
2. Experience a fair, transparent process and critical, constructive review.
3. If you are at a Publish and Read institution, you'll enjoy the benefits of Open Access across our journal portfolio.
4. Author feedback says our Editors are 'thorough and fair' and 'patient and caring'.
5. Increase your reach and impact and share your research more widely.

Find out more and submit your article at [microbiologyresearch.org](https://microbiologyresearch.org).