



# Investigating the effect of silicate- and calcium-based ocean alkalinity enhancement on diatom silicification

Aaron Ferderer<sup>1,2</sup>, Kai G. Schulz<sup>3</sup>, Ulf Riebesell<sup>4</sup>, Kirralee G. Baker<sup>1,5</sup>, Zanna Chase<sup>1</sup>, and Lennart T. Bach<sup>1</sup>

<sup>1</sup>Institute for Marine and Antarctic Studies, Ecology & Biodiversity, University of Tasmania, Hobart, TAS, Australia

<sup>2</sup>National Collections and Marine Infrastructure, Commonwealth Scientific and Industrial Research Organisation, Hobart, TAS, Australia

<sup>3</sup>Faculty of Science and Engineering, Southern Cross University, Lismore, NSW, Australia

<sup>4</sup>Marine Biogeochemistry, Biological Oceanography, GEOMAR Helmholtz Centre for Ocean Research Kiel, Kiel, Germany

<sup>5</sup>The Australian Centre for Excellence in Antarctic Science (ACEAS), University of Tasmania, Hobart, TAS, Australia

**Correspondence:** Aaron Ferderer (aaron.ferderer@utas.edu.au)

Received: 29 August 2023 – Discussion started: 12 September 2023

Revised: 26 April 2024 – Accepted: 2 May 2024 – Published: 12 June 2024

**Abstract.** Gigatonne-scale atmospheric carbon dioxide removal (CDR) will almost certainly be needed to supplement the emission reductions required to keep global warming between 1.5–2 °C. Ocean alkalinity enhancement (OAE) is an emerging marine CDR method with the addition of pulverised minerals to the surface ocean being one widely considered approach. A concern of this approach is the potential for dissolution products released from minerals to impact phytoplankton communities. We conducted an experiment with 10 pelagic mesocosms (M1–M10) in Raunefjorden, Bergen, Norway, to assess the implications of simulated silicate- and calcium-based mineral OAE on a coastal plankton community. Five mesocosms (M1, M3, M5, M7, and M9) were enriched with silicate ( $\sim 75 \mu\text{mol L}^{-1} \text{Na}_2\text{SiO}_3$ ), alkalinity along a gradient from 0 to  $\sim 600 \mu\text{mol kg}^{-1}$ , and magnesium in proportion to alkalinity additions. The other five mesocosms (M2, M4, M6, M8, M10) were enriched with alkalinity along the same gradient and calcium in proportion to alkalinity additions. The experiment explored many components of the plankton community, from microbes to fish larvae, and here we report on the influence of simulated mineral based OAE on diatom silicification. Macronutrients (nitrate and phosphate) limited silicification at the onset of the experiment until nutrient additions on day 26. Silicification was significantly greater in the silicate-based mineral treatment, with all genera except *Cylindrotheca* displaying an increase in silicification as a result of the increased concentration

of dissolved silicate. In contrast to the effect of differences in dissolved silicate concentrations between the two mineral treatments, increases in alkalinity only influenced the silicification of two genera, *Pseudo-nitzschia* and *Nitzschia*. The four other genera (*Arcocellulus*, *Cylindrotheca*, *Skeletonema*, and *Thalassiosira*) investigated here displayed no significant changes in silicification as a result of alkalinity increases between 0 and  $600 \mu\text{mol kg}^{-1}$  above natural levels. In summary, our findings illustrate that the enhancement of alkalinity via simulated silicate- and calcium-based methods has limited genus-specific impacts on the silicification of diatoms. This research underscores the importance of understanding the full breadth of different OAE approaches, their risks, co-benefits, and potential for interactive effects.

## 1 Introduction

Limiting global average surface temperature rise to 1.5–2 °C above pre-industrial levels necessitates rapid reductions in global CO<sub>2</sub> (carbon dioxide) emissions as well as sustained atmospheric carbon dioxide removal (CDR) (IPCC, 2021; Van Vuuren et al., 2018). However, prior to considering the implementation of large-scale CDR methods it is critical to assess the potential ecological impacts of these methods (Bach et al., 2019; Fuss et al., 2018; Renforth and Henderson, 2017).

Ocean alkalinity enhancement (OAE) is considered to be a promising marine CDR method due to its potential to remove CO<sub>2</sub> at a gigatonne scale (Burt et al., 2021; Feng et al., 2017; He and Tyka, 2023; Ilyina et al., 2013; Keller et al., 2014; Paquay and Zeebe, 2013). There are various approaches to implementing OAE, each with techno-economic and environmental advantages and disadvantages (Renforth and Henderson, 2017). Irrespective of the method, all approaches aim to increase the capacity of the ocean to store atmospheric CO<sub>2</sub> by increasing the alkalinity of seawater through the addition of substances that increase alkalinity, the removal of acid, and/or neutralisation of protons in seawater, all of which will increase seawater pH (Eisaman et al., 2023). One widely discussed method involves the addition of pulverised alkaline minerals such as magnesium silicates or calcium hydroxides to the surface ocean (Bach et al., 2019; Kheshgi, 1995; Renforth and Henderson, 2017).

In order for minerals to be suitable they must be alkaline; have relatively rapid dissolution rates; and ideally inexpensive, readily available, and containing minimal potential contaminants (Hartmann et al., 2013; Renforth and Henderson, 2017). Minerals fitting some or most of these criteria include olivine, a silicate-based naturally occurring mineral, as well as quick and/or hydrated lime, which are anthropogenic calcium-based minerals (Renforth and Henderson, 2017). However, the effects of dissolution products derived from these minerals on marine communities is yet to be fully assessed (Bach et al., 2019). Indeed, hotspots of dissolution products from mineral-based OAE will inevitably occur at sites of mineral additions resulting in high concentrations of, e.g. Mg<sup>+</sup>, Si(OH)<sub>4</sub>, Ca<sup>+</sup>, increased pH, and trace metals (Hartmann et al., 2013). Dissolution products may act to fertilise some organisms while inhibiting others, potentially leading to shifts in plankton communities (Bach et al., 2019; Guo et al., 2022; Hutchins et al., 2023). For example, calcium-based minerals are hypothesised to benefit pelagic and benthic calcifiers, with some studies supporting this (Albright et al., 2016; Bach et al., 2015; Gore et al., 2019), while others found neutral responses (Gately et al., 2023). In contrast, the dissolution of silicate-based minerals is expected to benefit silicifying plankton species including diatoms (Bach et al., 2019; Egge and Aksnes, 1992; Hauck et al., 2016). Thus, we expect mineral-based OAE to have some impact on marine communities with these impacts being highly dependent on the source minerals used. However, it is important that the environmental impacts and/or co-benefits resulting from OAE are evaluated against the potential climatic benefits.

This study aims to specifically assess and compare the potential implications of increased alkalinity, silicate, and magnesium concentrations associated with silicate-based mineral OAE and increased alkalinity and calcium concentrations associated with calcium-based mineral OAE on a coastal plankton community. In order to capture the potential maximum and minimum acceptable levels of alkalinity enhance-


ment, we increased concentrations of alkalinity in steps of 150 μmol kg<sup>-1</sup> by 0 to ~ 600 μmol kg<sup>-1</sup>. Such an increase in alkalinity is expected to influence the phytoplankton community as concentrations of CO<sub>2</sub> decrease below previously observed thresholds limiting growth (Chen and Durbin, 1994; Hinga, 2002; Paul and Bach, 2020; Riebesell et al., 1993).

Previous work has identified that increases in alkalinity of 500 μmol kg<sup>-1</sup> resulted in a significant decrease in silicate uptake and biogenic silica production (Ferderer et al., 2022). Furthermore, it is well known that silicate has a fertilising effect on diatoms, with concentrations above 2 μmol kg<sup>-1</sup> often resulting in their dominance within plankton communities (Egge and Aksnes, 1992; Escaravage and Prins, 2002). The dissolution of silicate-based minerals such as olivine to enhance alkalinity is predicted to significantly increase silicate concentrations at sites of addition and projected to induce diatom blooms (Hauck et al., 2016). The influence of varying silicate concentrations on diatoms is well known; however, the interaction between OAE and enhanced silicate concentrations as a result of silicate-based mineral dissolution is yet to be fully explored. Thus, in this study we focus on assessing the influence of mineral-based OAE, along an increasing gradient, on the incorporation of silica into the frustules of diatoms (silicification). Our primary goal is to elucidate the potential risks and or co-benefits of mineral-based alkalinity enhancement on diatoms.

## 2 Methods

### 2.1 Mesocosm deployment and maintenance

On the 7 May 2022, a total of 10 Kiel Off-Shore Mesocosms for Ocean Simulations (KOSMOS, M1–M10; Riebesell et al., 2013) devices were deployed from RV *ALKOR* in Raunefjorden, Bergen, Norway, ~ 1.5 km from the Espregrend marine research field station (Fig. 1). Mesocosms consisted of a cylindrical polyurethane bag 20 m in length, (2 m in diameter, ~ 60.01 ± 0.01 m<sup>3</sup> volume). Mesocosm bags were fitted within 8 m tall floating frames during deployment which were slowly lowered into the fjord, allowing the bags to gently fill while minimising disturbance to the plankton community. Once deployed the mesocosm bags remained open at their base (~ 20 m) and top (~ 1 m below the sea surface) allowing water exchange between the mesocosms and fjord. Mesocosms were closed off to the fjord on the 13 May when divers attached a 2 m long funnel-shaped sediment trap to each mesocosm, and the top of each mesocosm bag was raised ~ 1 m above the surface (Fig. 1). A ring with the same diameter as the mesocosms fitted with a 1 mm mesh was passed through each mesocosm after closing to remove any large nekton or plankton from the mesocosms. The sealing off of the mesocosms from the fjord marked the beginning of the experiment (Day 0). The volume of each mesocosm was determined on day 2 of the experiment via the addi-



**Figure 1.** Infographic depicting the relevant information pertaining to the mesocosm design and mooring site approximately 1.5 km from the Espesrend Marine research field station in Bergen, Norway (maps produced in RStudio using © Google Maps data).

tion of a NaCl brine solution. Water inside mesocosms was first homogenised by bubbling compressed air up through the mesocosms. Following homogenisation, 50 L of NaCl brine was evenly added to the mesocosm via a bespoke distribution device called “the spider” (Riebesell et al., 2013). The precise addition of NaCl enabled us to calculate the volume of each mesocosm following (Czerny et al., 2013). Mesocosm bags were cleaned approximately every week from the inside and outside to minimise any potential biofouling, which may impact the results of the experiment. External cleaning of the mesocosms was conducted by divers and/or surface attendants in small boats using brushes. Internal cleaning of the mesocosms was conducted using a large ring with rubber blades that was the same diameter as the mesocosms. This ring was sunk inside mesocosms with a 30 kg weight attached to its base, removing any growth from the inner walls of the mesocosm bags down to 1 m above the sediment trap.

## 2.2 Setup of OAE treatments and nutrient fertilisation

Mesocosms were split into two treatment groups: a calcium-based (Ca-OAE) treatment ( $N = 5$ ) and a silicate-based (Si-OAE) treatment ( $N = 5$ ) with one mesocosm in each group serving as a control. Alkalinity was enhanced along a gradient in each mineral-based treatment ranging from 0–600  $\mu\text{mol kg}^{-1}$  using varying amounts of NaOH (Merck), dissolved in 20 L of Milli-Q<sup>®</sup>. Simulated differences in the type of OAE were established via the addition of  $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$  in the Ca-OAE treatments and  $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$  and  $\text{Na}_2\text{SiO}_3 \cdot 5\text{H}_2\text{O}$  in the Si-OAE treatments, all dissolved in 20 L of Milli-Q<sup>®</sup>. The simulated enhancements of  $\text{Mg}^{2+}$  and  $\text{Ca}^{2+}$  were proportional to the addition of NaOH, i.e. increases by half the alkalinity enhancement. In contrast  $\text{Na}_2\text{SiO}_3$  was increased by equal concentrations (target of 75  $\mu\text{mol L}^{-1}$ ) in all mesocosms within the Si-OAE treatment group (including the control), instead of a gradient from 0–150  $\mu\text{mol L}^{-1}$ , which would be the corresponding concentrations for olivine dissolution. This approach

was adopted due to metasilicate solubility restrictions (data not shown), the potential for colloid formation to occur at high concentrations and enable clear distinctions to be made between silicate and alkalinity effects. Finally, the increase in alkalinity from silicate additions in a 2 : 1 ratio was taken into account by reducing respective NaOH additions and the addition of HCl in the silicate-based control ( $\Delta\text{alkalinity} = 0 \mu\text{mol kg}^{-1}$ ).

At the time of closure, all mesocosms had low concentrations of macronutrients ( $0.10 \pm 0.019 \mu\text{mol L}^{-1} \text{NO}_3^-$ ,  $0.03 \pm 0.005 \mu\text{mol L}^{-1} \text{PO}_4^{3-}$ , and  $0.16 \pm 0.048 \mu\text{mol L}^{-1} \text{Si(OH)}_4$ ). After observing communities in a prolonged phase of oligotrophic conditions, macronutrients were added to the mesocosms on day 26 (final range across mesocosms =  $3.59\text{--}3.8 \mu\text{mol L}^{-1} \text{NO}_3^-$ ,  $0.19\text{--}0.24 \mu\text{mol L}^{-1} \text{PO}_4^{3-}$ , and  $0.39\text{--}1.03 \mu\text{mol L}^{-1} \text{Si(OH)}_4$ ) to stimulate the phytoplankton community. Macronutrients were added to the mesocosms as two separate 20 L solutions with one consisting of  $\text{NaNO}_3$  (Merck, > 99.5 %) and  $\text{Na}_2\text{HPO}_4 \cdot \text{H}_2\text{O}$  (Merck, > 99.5 %) and the other consisting of  $\text{Na}_2\text{SiO}_3 \cdot 5\text{H}_2\text{O}$  (Roth > 95 %). Inorganic nutrient concentrations were measured the day before nutrient additions and  $\sim 2$  h after the addition of nutrients to quantify the additions and ensure appropriate stoichiometry within mesocosms. After the addition of nutrients, it was noted that the stoichiometry of macronutrients was not even across mesocosms, later identified to have been the result of a mistake during solution preparation. As such, a second addition of nitrate was completed on day 28 for those mesocosms below target concentrations. All solutions were added homogeneously to mesocosms using the spider distribution device.

Given the additions of alkalinity on day 7 and macronutrients on day 26 and 28, the experiment was divided into three distinct phases: phase 0 representing the period prior to alkalinity enhancement (day 0–6), phase I representing conditions prior to the addition of macronutrients (day 7–28), and phase II representing the period after nutrient additions (day 29–54).

### 2.3 Sampling methods

Sampling of the mesocosms was conducted every second day from small boats with sediment sampling first (08:00–10:00, here and in the following times are provided in GMT + 2) followed by particulate and dissolved substance sampling (09:00–13:00), zooplankton sampling (10:00–13:00), and finally conductivity–temperature–depth (CTD) FastOcean Ambient plus Dark (APD)/fluoroprobe (14:00–16:00). With the exception of particulate and dissolved substance sampling, which was carried out at random, mesocosms were sampled in order from M1 through to M10 with fjord samples taken directly next to M5. Sample containers were stored in boxes to avoid excess light and heat exposure during sampling and upon return to the research station (directly after

each round of sampling) were transferred to a room at ambient water temperature ( $8.7\text{--}15.4^\circ\text{C}$ ) until further processing. The sampling schedule remained consistent with the exception of day 15 where only sediment sampling was undertaken due to unsafe weather conditions. Additional samples for the determination of dissolved inorganic nutrients were taken on day 26 and day 28 to assess stoichiometry post nutrient additions performed earlier the same day.

Sinking particles were collected from the sediment traps of each mesocosm via a silicon tube attached to the base of the sediment trap at one end and a manual vacuum pump at the surface (Boxhammer et al., 2016). Suspended particulate matter and dissolved substances were collected using 5 L integrated water samplers (IWSs; Hydro-Bios, Kiel). IWSs were equipped with pressure sensors enabling an even collection of water within a specified depth, from the surface to the top of the sediment trap (0–20 m). Four IWSs were taken within each mesocosm and fjord, which were transferred into 10 L polyethylene carboys. Samples for quantification of changes in carbonate chemistry were collected from the first IWS taken within each mesocosm and filled directly into 500 mL glass bottles following protocols outlined in SOP 1 from Dickson et al. (2007).

### 2.4 Carbonate chemistry and dissolved inorganic nutrients

Samples for total alkalinity (TA), pH, and dissolved inorganic nutrients were sterile filtered using a peristaltic pump and 25 mm, 0.2  $\mu\text{m}$  pore size, polyethersulfone (PES) membrane, syringe filters to minimise biological processes and remove particles which can influence respective analyses. Dissolved inorganic nutrient concentrations  $\text{NO}_3^-$ ,  $\text{NO}_2^-$ ,  $\text{PO}_4^{3-}$ , and  $\text{Si(OH)}_4$  were determined spectrophotometrically following methods outlined by Hansen and Koroleff (1999). Dissolved inorganic nutrient samples were measured in triplicate to control for technical variability between measurements across the experiment. TA was determined using a two-step open cell titration following SOP3b outlined by Dickson et al. (2007). TA samples were measured in duplicate on a Metrohm 826 Compact Titrosampler coupled with an Aquatrod Plus with PT1000 temperature sensor and calibrated against certified reference material (CRM batch 193) supplied by Andrew Dickson's laboratory. The pH was determined in duplicate via spectrophotometric methods outlined in Dickson et al. (2007) (not shown here).

### 2.5 Particulate matter analysis

Sediment trap samples were processed immediately upon return of the sampling boat to the research station. Sample weight was first determined gravimetrically before resuspension of particles and homogenisation of the sample for subsampling (e.g. dissolution assays, particle sinking velocity). The remaining sample was enriched with 3 M  $\text{FeCl}_3$  fol-

lowed by 3 M NaOH to enhance flocculation, coagulation, and subsequent sedimentation of particles while maintaining pH (Boxhammer et al., 2016). Approximately 1 h after settling the supernatant was removed and samples centrifuged in two steps: first for 10 min at 5200 g in a 6–16KS centrifuge (Sigma) and then again for 10 min at 5000 g in a 3K12 centrifuge (Sigma). Following each step any supernatant was removed and the remaining pellet freeze dried to remove residual moisture. Finally, the dried samples were pulverised into a homogenous powder using a cell mill and were transported to GEOMAR, Kiel, Germany, for further analysis.

Subsamples of the powder used to determine concentrations of biogenic silica (BSi) were placed in 60 mL Nalgene™ polypropylene bottles, filled with 25 mL of 0.1 M NaOH solution, and then placed in a shaking water bath at 85 °C. After 135 min the bottles were removed and cooled before the addition (25 mL) of 0.05 M H<sub>2</sub>SO<sub>4</sub> to stop the leaching processes. The concentration of dissolved silicate was then measured spectrophotometrically following Hansen and Koroleff (1999). BSi concentrations of the measured subsamples were then scaled to represent the total sample from sediment traps normalised to mesocosm volume and are reported as the accumulation of BSi in the sediments over the experimental period.

Analysis of major elemental pools and phytoplankton pigments in the water column subsamples (pre-filtered through a 200 µm screen) of 0.5–1 L were taken from carboys after gentle mixing to homogenise samples. Subsamples for BSi were filtered onto cellulose acetate filters (pore size 0.45 µm) using a vacuum filtration system at ≤ 200 mbar and stored in plastic vials at –20 °C until analysis the following day. Filters were then placed in 60 mL Nalgene™ bottles, digested, and analysed following the same methods for BSi in the sediments described above. Chlorophyll *a* was filtered onto glass fibre filters (GFFs, nominal pore size = 0.7 µm) while minimising light exposure. Immediately after filtration filters were stored in plastic vials at –80 °C until analysis the following day. Samples were extracted with 90 % acetone and homogenised using glass beads in a cell mill. After homogenisation samples were centrifuged (10 min 800 g, 4 °C), then the supernatant was removed and analysed on a fluorometer (Turner 10-AU) to determine Chl *a* concentrations (Welschmeyer, 1994).

## 2.6 PDMPO labelling and determination of silicification rates via fluorescent microscopy

To investigate differences in diatom silicification, 350 mL samples were collected by means of IWS from each mesocosm every two to four days (variation in sampling schedule occurred due to unforeseen circumstances such as extreme weather events and COVID-19 infections). Due to the low abundance of diatoms within mesocosms (as observed through microscopy and BSi concentrations), samples were gravimetrically concentrated from 350 to 70 mL using a

47 mm filtering apparatus and polycarbonate filters (3 µm pore size). Organisms were resuspended by gentle stirring and inversion of the filtration system before the sample was transferred to three 17 mL polycarbonate tubes. The fluorescent dye 2-(4-pyridyl)-5-((4-(2-dimethylaminoethyl-aminocarbonyl)methoxy)phenyl)oxazole (PDMPO; LysoSensor yellow/blue DND-160 from ThermoFisher Scientific) was added at a final concentration of 0.125 µM before samples were incubated in a flow-through incubator placed outside of the Espeyend Marine research field station for temperature control (inlet location was ~ 1.5 km from the mesocosms so that temperature in the incubator was similar to mesocosms). The incubator was screened with shade cloth to ~ 30 % incident irradiance and incubations lasted for ~ 24 h.

At the conclusion of the incubation, 15 mL of the sample was filtered under gentle vacuum pressure onto 25 mm, 0.8 µm black polycarbonate membrane filters. Filters were mounted onto microscope slides with a drop of Prolong gold antifade followed by a glass coverslip and sealed with clear nail varnish. Prepared slides were then stored in the dark at –20 °C for later analysis via fluorescent microscopy at the University of Tasmania, Australia (within 6 months).

Prepared microscope slides were imaged using the software NIS elements and a Nikon eclipse Ci microscope equipped with a UV-1A (long-pass) filter cube, Nikon DS-Ri2 camera and Mercury lamp (Nikon C-SHG1). Prior to analysing slides, a yellow fluorescence slide (Thorlabs FSK3) was imaged 10 times, and the average fluorescence subtracted from all images taken that day to account for variation in the mercury lamp across imaging days. The entirety of each filter was systematically scanned at ×200 magnification; when a cell was located, it was imaged at ×400 magnification with all cells that were appropriate for measurements (e.g. not overlapping or partially destroyed cells) included in analysis. This gave final counts for each genus ranging from 1–176 cells per mesocosm on any given day. Cells were, when possible, identified to genus level as bright-field imaging was not possible on the black polycarbonate filters and fluorescent images did not provide enough detail for accurate identification beyond this level. However, in instances where differentiation between genera was difficult or impossible to complete with high confidence, cells were classified based on significant differences in the shape, size, and/or details of the frustule of cells. Each genus or group is therefore comprised of similar cells that show distinct differences in characteristics that influence the fluorescence of cells (Table S1). Images were later analysed by quantifying single-cell fluorescence following a custom-made procedure in ImageJ on the original TIFF images. For each image, cells were selected so that minimal background area was included before the fluorescence of the selected cell was recorded. Background fluorescence was measured at four locations directly surrounding the cell where no other cells were present and the average background fluorescence subtracted from cell fluores-

cence. Total cell fluorescence, corrected for background fluorescence, was then normalised to cell area to give mean cell fluorescence. Due to low abundances of diatoms and thus insufficient counts for meaningful analyses, only days 7, 11, and 17 were analysed prior to the addition of nutrients on day 26 and 28. After the addition of nutrients all filters were analysed; however, due to an outbreak of COVID-19, samples for the determination of silicification could not be taken for the final 8 d of the experiment.

Due to technical issues and the unavailability of specific instrumentation, we were unable to measure total community fluorescence during the experiment and as such convert fluorescence values to BSi incorporation. However, the aim of this experiment was to identify any relative differences in taxa-specific rates of silicification, something which is not achievable through the measurement of BSi. This was achieved with the use of the fluorescent dye PDMPO, which is incorporated at a rate proportional to BSi incorporation in diatoms (Leblanc and Hutchins, 2005; McNair et al., 2015; Znachor and Nedoma, 2008). PDMPO uptake and subsequent fluorescence within diatom cells therefore provides an appropriate proxy for the incorporation of silica into newly formed frustules irrespective of the units. As such, the PDMPO fluorescence of cells measured here as the fluorescence of a given cell normalised to cell area is referred to as silicification throughout this text.

## 2.7 Statistical analysis

To explore the effect of the alkalinity source mineral and alkalinity enhancement across mesocosms we first visualised the dataset using non-metric multidimensional scaling (NMDS) plots. Three separate plots were produced to explore the effects of (a) the treatments over the total extent of the experiment, (b) prior to the addition of nutrients, and (c) post nutrient addition on silicification.

A linear mixed-effect model was used to quantify the influence of the treatments (total alkalinity and alkalinity source mineral) on diatom silicification. The model was run with alkalinity source mineral, total alkalinity, experimental phase, and diatom genus as fixed effects, and silicification (square root transformed) as the dependent variable. To account for temporal pseudo-replication in the models, mesocosm ( $N = 10$ ) was nested within sampling occasion (day) and fitted as a random effect in all models. Several linear mixed-effect models were fit, with non-significant interactions removed and the Akaike information criterion (AIC) used to determine the best model. All statistical analyses were conducted in RStudio v 2023.6.1.524 (R Core Team, 2023). Estimated marginal means were calculated using the package *emmeans* (Lenth et al., 2023) to determine the significance of two- and three-way interactions within the model. Estimated marginal means of linear trends were calculated using *emtrends* from the package *emmeans* to assess the effect of total alkalinity


within significant interactions in the linear model (Lenth et al., 2023).

Finally, linear models were used to assess the influence of total alkalinity and alkalinity source mineral on the concentration of BSi in the water column and accumulation of BSi in the sediments. Linear models were run for each phase of the experiment with mean water column BSi and mean accumulated sediment BSi over each phase fitted as dependent variables and mean total alkalinity and alkalinity source mineral as fixed effects. An additional linear model was run to assess total accumulated BSi in the sediment trap with the accumulated sediment BSi up until day 53 fitted as the dependent variable and total alkalinity and alkalinity source mineral fitted as fixed effects. All statistical analyses including NMDS plots were conducted in RStudio v 2023.6.1.524 (R Core Team, 2023).

## 3 Results

Concentrations of  $\text{NO}_3^-$  were below detection limit, thereby constraining phytoplankton growth during phase I of the experiment (mean  $\text{NO}_3^-$  day 7–25 =  $0.004 \pm 0.035 \mu\text{mol L}^{-1}$ ) (Fig. 2a). In contrast, there was residual  $\text{PO}_4^{3-}$  ( $0.021 \pm 0.022 \mu\text{mol L}^{-1}$ ) and  $\text{Si}(\text{OH})_4$  (Ca-OAE treatment =  $0.202 \pm 0.99 \mu\text{mol L}^{-1}$ , Si-OAE treatment =  $67.929 \pm 1.04 \mu\text{mol L}^{-1}$ ), which likely supported the phytoplankton community in utilising remineralised nitrogen until the addition of nutrients on day 26 and 28 (Fig. 2b and c). Although  $\sim 75 \mu\text{mol L}^{-1}$  of  $\text{Na}_2\text{SiO}_3$  was added to the Si-OAE treatment, there was no discernible depletion of  $\text{Si}(\text{OH})_4$  during phase I (Fig. 2c). The addition of macronutrients ( $\text{NO}_3^-$ ,  $\text{PO}_4^{3-}$ , and  $\text{Si}(\text{OH})_4$ ) can be seen on day 26, with a secondary addition on day 28 to correct for unwanted differences in the stoichiometry between mesocosms (Fig. 2). Chlorophyll *a* concentrations were relatively low at the beginning of the experiment with  $1.01 \pm 0.17 \mu\text{g L}^{-1}$  (mean  $\pm$  SD) on day 3 (Fig. 2d). During phase II, nutrients steadily declined until the majority was depleted between days 39 and 49 (Fig. 2). Chlorophyll *a* remained low until day 33, after which it increased across all mesocosms at rates between  $0.03$ – $0.68 \mu\text{g L}^{-1} \text{d}^{-1}$  (Fig. 2d). The delayed or slow increase in chlorophyll *a* was likely due to the prolonged nutrient deficit within the mesocosms and subsequent small seed population. There was no discernible relationship between total alkalinity and chlorophyll *a*,  $\text{NO}_3^-$ , or  $\text{PO}_4^{3-}$  observed across the extent of the experimental period or in a particular phase (Fig. 2). However, in the Si-OAE treatments initial concentrations of  $\text{Si}(\text{OH})_4$  were lowest in the high-alkalinity mesocosm, with a difference of  $2.45 \mu\text{mol L}^{-1}$  between the  $\Delta 0$  and  $\Delta 600 \mu\text{mol kg}^{-1}$  alkalinity treatments (Fig. 2c). This trend appeared directly after the addition of the treatments but disappeared once nutrient uptake began (Fig. 2c).






**Figure 2.** Temporal variation in dissolved inorganic nutrients (a) nitrate ( $\text{NO}_3^-$ ), (b) phosphate ( $\text{PO}_4^{3-}$ ), (c) silicate ( $\text{Si}(\text{OH})_4$ ), (d) chlorophyll *a* (Chla), and (e) total alkalinity. Dissolved inorganic nutrient measurements commenced on day 0 while chlorophyll *a* measurements commenced on day 3. Vertical dashed lines represent the respective phases of the experiment: phase 0 (pre-alkalinity enhancement), phase I (pre-nutrient addition), phase II (post-nutrient addition).

Non-metric multidimensional scaling (NMDS) (Fig. 3) revealed distinct distances among treatments, including different alkalinity source minerals and total alkalinity, in relation to silicification of the various diatom genera. The distances between polygons, representing the Si- and Ca-based mineral treatments, indicate differences in silicification between the Si- and Ca-based mineral treatments over the total experimental period and after the addition of nutrients (Fig. 3a and c). In contrast, during phase I (pre-nutrient addition) polygons representing the Si- and Ca-based mineral treatments are overlapping, suggesting a weak relationship between alkalinity source mineral (Ca or Si) and silicification

prior to the addition of nutrients (Fig. 3b). In all plots, symbols representing the differing levels of total alkalinity within the Si-based treatment are relatively close when compared to the Ca-based treatment, which exhibits greater spread between differing levels of total alkalinity.

### 3.1 Results of linear mixed-effect model

Analysis of diatom silicification supported the distances observed within the NMDS plots with alkalinity source mineral (Ca or Si) having a significant influence on the silicification of diatom cells (Table 1, Fig. 4). Cells exposed to the Si-OAE treatment were more heavily silicified irrespective of changes



**Figure 3.** Non-metric multidimensional scaling ordination exploring the mean silicification of diatom genera across (a) the complete extent of the mesocosm experiment (stress = 0.0502); (b) pre-nutrient addition, phase I (stress =  $3.62 \times 10^{-5}$ ); and (c) post nutrient addition, phase II (stress = 0.0612). Due to the low stress values obtained ( $< 0.20$ ), it is assumed that all configurations accurately represent distinct dissimilarities in silicification among diatom genera.

**Table 1.** Statistical results of the linear mixed-effect model assessing the influence of mineral-based OAE on diatom silicification.


Source of variation	df	<i>f</i> value	<i>p</i> value
Alkalinity source mineral (Ca or Si)	1.64	146.95	$< 0.001^*$
Genus	5.11330	1305.96	$< 0.001^*$
Total alkalinity	1.64	1.59	0.21
Phase	1.6	5.26	0.06
Alkalinity source mineral · genus	5.11330	82.07	$< 0.001^*$
Alkalinity source mineral · total alkalinity	1.64	0.04	0.84
Total alkalinity · genus	5.11330	57.65	$< 0.001^*$
Alkalinity source mineral · phase	1.64	1.29	0.26
Total alkalinity · phase	1.64	4.64	0.03*
Alkalinity source mineral · genus · total alkalinity	5.11330	5.93	$< 0.001^*$
Alkalinity source mineral · total alkalinity · phase	1.64	4.02	0.05*

\*  $P < 0.05$ .

in total alkalinity (Fig. 4). However, the significant interaction between alkalinity source mineral and genus indicates that the effect of the alkalinity source mineral on silicification varies between genera (Table 1). All genera displayed significant differences between the two alkalinity source mineral treatments with the exception of *Cylindrotheca* which showed no significant difference in silicification between the Si- or Ca-based mineral treatments (Table 2). Total alkalinity

had a significant effect on silicification between the two experimental phases ( $t$  ratio =  $-2.16$ ,  $p = 0.03$ ), with silicification increasing as a function of total alkalinity during phase I of the experiment (emtrend = 0.24,  $t$  ratio = 2.57,  $p = 0.01$ ) (Table 1, Fig. 4). However, investigation of the three-way interaction revealed the significant effect of total alkalinity on diatom silicification only to be present in phase I of the experiment in the Si-based OAE treatment ( $t$  ratio = 3.22,





**Figure 4.** Single-cell silicification of the diatom community depicted as mean fluorescence (PDMPO) normalised to cell surface area and reported in fluorescence units (FU). Data visualised as box plots, with colours representing the different mineral sources (Ca-OAE; blue and Si-OAE; green) and shading indicating the total alkalinity gradient with darker colours indicating higher total alkalinity. Nutrient addition on day 26 and 28 is represented by the dashed line dividing the experiment into phase I (pre-nutrient addition) and phase II (post-nutrient addition).

**Table 2.** Results of post hoc tests (*emmeans*) comparing the influence of silicate- and calcium-based mineral treatments on genera-specific silicification.

Genus	Contrast	Estimate	Standard error	<i>t</i> ratio	<i>p</i> value
<i>Arcocellulus</i>	Ca – Si	−0.69	0.08	−8.64	< 0.001*
<i>Cylindrotheca</i>	Ca – Si	0.07	0.11	−0.67	0.51
<i>Nitzschia</i>	Ca – Si	−0.60	0.10	−6.02	< 0.001*
<i>Pseudo-nitzschia</i>	Ca – Si	−0.41	0.08	−5.04	< 0.001*
<i>Skeletonema</i>	Ca – Si	−1.24	0.09	−14.34	< 0.001*
<i>Thalassiosira</i>	Ca – Si	−0.91	0.09	−10.68	< 0.001*

\*  $P < 0.05$ .

$p = 0.002$ ) (Fig. 4, Table 1). Furthermore, the significant interaction between genus and total alkalinity suggests the influence of total alkalinity on silicification varies between genera.

Finally, the significant three-way interaction between alkalinity source mineral, genus, and total alkalinity suggests that the effect of alkalinity on silicification in given genera was dependent on the alkalinity mineral source (Table 1). Exploration of this interaction revealed the silicification of

cells in the genus *Pseudo-nitzschia* ( $N = 3510$ ) to be significantly influenced by alkalinity in both the Ca- and Si-based treatments, with silicification increasing with increasing alkalinity (Table 3). In contrast, the genus *Nitzschia* ( $N = 677$ ) displayed a significant increase in silicification in the Si-based treatment only while no other genera displayed significant differences in silicification between total alkalinity levels (Table 3). There were also significant differences in silicification between genera; however, this was reduced from

**Table 3.** Results of post hoc tests (*emrends*) assessing the influence of total alkalinity on genera-specific silicification across the two alkalinity source mineral treatments.

Genus	Mineral	Total alkalinity mean trend	Standard error	<i>t</i> ratio	<i>p</i> 'value
<i>Arcocellulus</i>	Ca	0.004	0.07	0.06	0.96
<i>Cylindrotheca</i>	Ca	0.10	0.09	1.13	0.26
<i>Nitzschia</i>	Ca	0.08	0.08	1.07	0.29
<i>Pseudo-nitzschia</i>	Ca	0.33	0.07	4.52	< 0.001*
<i>Skeletonema</i>	Ca	0.06	0.08	0.79	0.43
<i>Thalassiosira</i>	Ca	−0.06	0.08	−0.76	0.45
<i>Arcocellulus</i>	Si	0.00	0.07	0.01	0.99
<i>Cylindrotheca</i>	Si	0.06	0.09	0.64	0.52
<i>Nitzschia</i>	Si	0.40	0.08	5.02	< 0.001*
<i>Pseudo-nitzschia</i>	Si	0.37	0.07	5.31	< 0.001*
<i>Skeletonema</i>	Si	0.03	0.07	0.50	0.62
<i>Thalassiosira</i>	Si	−0.04	0.07	−0.61	0.55


\* *P* < 0.05.

eight significant differences in the Si-based mineral treatment to five in the Ca-based mineral treatment (Table S2).

Concentrations of BSi in the water column can be seen decreasing from day 0 (mesocosm closure) and remaining low until day ~ 33 (Fig. 6a). Directly after the addition of Na<sub>2</sub>SiO<sub>3</sub> (day 7), BSi in the water column spiked for 1 d in the silicate-based treatments, while no increase was observed in the calcium-based treatment (Fig. 6a). BSi began to increase in all mesocosms between days 33–35; however, concentrations in the Ca-OAE treatments remained relatively low (< 2 μmol L<sup>−1</sup>) for the extent of the experiment (Fig. 6a). We observed no significant relationship between BSi and alkalinity in either of the alkalinity source mineral treatments (Ca or Si) or any phase of the experiment (Table 3). During phase II, concentrations of BSi in the water column were significantly higher in the Si-OAE treatment when compared to the Ca-OAE treatment (Table 4, Fig. 6a). Additionally, we observed significant differences in the accumulation of BSi in the sediments between the two alkalinity source mineral types (Table 4, Fig. 6b). Similar to the water column, there was an initial increase in sedimented BSi in the Si-OAE treatment when compared to the Ca-OAE treatment (Fig. 6b). During phase I there was a 0.47 μmol L<sup>−1</sup> difference in the amount of BSi accumulated in the sediment trap between the highest (Δ600 μmol kg<sup>−1</sup>) and lowest (Δ0 μmol kg<sup>−1</sup>) alkalinity levels in the Si-OAE treatment. However, there was less variability between the Δ150, Δ300, and Δ450 alkalinity treatments, contributing to the non-significant effect of alkalinity on BSi accumulation in the sediment trap (Table 3). Furthermore, irrespective of experimental phase, there was no significant relationship between alkalinity and BSi accumulated in the sediment.

#### 4 Discussion

Understanding the potential environmental implications of CDR methods such as OAE is a crucial step before decisions are made upon their implementation at large scales. The aim of this mesocosm study was to form part of this research by assessing the potential effects of calcium- and silicate-based mineral OAE on a coastal plankton community. Here, we specifically discuss the influence of simulated mineral based OAE on diatom-community- and genus-specific silicification. Our results revealed silicate fertilisation associated with silicate-based OAE to significantly increase silicification of the diatom community and all genera with the exception of *Cylindrotheca*. It is important to note that low initial concentrations of dissolved silicate likely facilitated this observed increase in silicification. Under conditions where dissolved silicate is already replete, further increases due to silicate-based OAE may not result in similar increases in additional silicification. This increase in silicification was primarily a result of the difference in the dissolved silicate concentrations (Δ75 μmol kg<sup>−1</sup>) between the silicate and calcium-based OAE treatments rather than an increase in alkalinity. However, in addition to the influence of silicate fertilisation, *Pseudo-nitzschia* and *Nitzschia* were both significantly affected by changes in alkalinity. *Pseudo-nitzschia* exhibited a significant increase in silicification with increasing alkalinity across both OAE treatments (Si and Ca), while *Nitzschia* only displayed an increase with silicification in the Si-OAE treatment. Overall trends observed in the silicification of the diatom community were confirmed by increased concentrations of BSi in the water column and accumulated in the sediment trap of mesocosms in the Si-OAE treatment. The increase of seawater alkalinity by 0–600 μmol kg<sup>−1</sup> had no effect on the concentration of BSi in the water column or accumulated in the sediment or overall diatom community silicification. In conjunction with published OAE research,



**Figure 5.** Boxplots depicting single-cell silicification of different genera. Silicification is shown as mean cell fluorescence (PDMPO) normalised to cell area and reported in fluorescence units (FU). Colours represent the different mineral sources (Ca-OAE, blue; Si-OAE, green), and shading indicates the total alkalinity gradient, with darker colours indicating higher total alkalinity. Nutrient addition on day 26 and 28 is represented by the dashed line dividing the experiment into phase I (pre-nutrient addition) and phase II (post-nutrient addition), with genera names in the top right of each plot.

our findings highlight the need for research to cover a broad range of environmental conditions, approaches to OAE, and marine communities.

#### 4.1 Temporal dynamics of biogenic silica in the water column and sediment

We observed no significant relationship between alkalinity and BSi in the water column or accumulated in the sediments (Fig. 6a and b). In contrast, concentrations of BSi in the water column and accumulated in the sediments were

significantly greater in the Si-OAE treatment before and after macronutrient additions. These trends support observations for community- and genera-specific silicification, with diatoms in the Si-OAE being more heavily silicified.

Notably, BSi accumulation in the sediments of the Si-based OAE treatment group was greatest in the highest alkalinity mesocosm ( $\Delta 600 \mu\text{mol kg}^{-1}$ ) and lowest in the control mesocosm ( $\Delta 0 \mu\text{mol kg}^{-1}$ ) during phase I (Fig. 6b). This difference emerged immediately after the addition of  $\text{Na}_2\text{SiO}_3$  to the Si-based OAE treatment. However, no build-up of

**Table 4.** Results of linear models exploring the effect of alkalinity source mineral and total alkalinity on average concentrations of BSi in the water column or sediment trap for a given phase.

Phase	Parameter	Source of variation	df	<i>f</i> value	<i>p</i> value
Phase 0	BSi water column	Mineral	1	0.70	0.43
		TA	1	0.70	0.43
Phase 0	BSi sediment	Mineral	1	0.20	0.67
		Total alkalinity	1	0.35	0.57
Phase I	BSi water column	Mineral	1	70.88	< 0.001*
		TA	1	1.76	0.23
Phase I	BSi sediment	Mineral	1	46.44	< 0.001*
		Total alkalinity	1	4.80	0.06
Phase II	BSi water column	Mineral	1	47.15	< 0.001*
		TA	1	2.02	0.20
Phase II	BSi sediment	Mineral	1	82.6109	< 0.001*
		Total alkalinity	1	0.64	0.45
Day 53	BSi sediment	Mineral	1	45.57	< 0.001*
		Total alkalinity	1	0.0005	0.98


\*  $P < 0.05$ .

chlorophyll *a* or significant build-up of BSi in the water column was observed in the days prior to the emergence of this trend, suggesting the sedimented BSi is an artefact of the  $\text{Na}_2\text{SiO}_3$  addition to the silicate-based treatments. Our interpretation of these findings are that (1) there were residual precipitates in the  $\text{Na}_2\text{SiO}_3$  solution added to the mesocosms, resulting in increased BSi in the water column, and/or (2) that there was pH-dependent, inorganic precipitation of amorphous silicate, with these precipitates sinking out into the sediment trap (Goto, 1956; Okamoto et al., 1957; Owen, 1975). The latter is supported by recent work conducted by Gately et al. (2023), whose abiotic experiments revealed a decrease in dissolved silicate as alkalinity increased, with scanning electron microscopy revealing mineral precipitates formed in high-alkalinity seawater to be primarily composed of silicon and oxygen. Precipitation of silica within mesocosms may have been supported by ionic interactions between magnesium (or trace metals, e.g. iron, aluminium) and silicate, pressure, and relatively low temperatures at depth (Ehlert et al., 2016; Goto, 1956; Spinthaki et al., 2018).

#### 4.2 Effect of enhanced silicate concentration on silicification rates

Ca- and Si-based OAE appeared to have a notable influence on silicification of the diatom community. However, this influence is attributed to the mineral treatment type, either silicate- or calcium-based, with a significant relationship between silicification and alkalinity observed for the *Pseudonitzschia* and *Nitzschia* (Si-OAE treatment only). Diatoms in the Si-OAE treatment incorporated considerably more sili-

cate over the 24 h incubation period resulting in increased silicification. This outcome was expected, especially considering the consistently low (likely limiting) concentrations of  $\text{Si}(\text{OH})_4$  observed in the Ca-OAE treatment throughout the majority of the experiment.  $\text{Si}(\text{OH})_4$  is the key nutrient in the construction of the silicate-based frustule of diatoms, with low concentrations often becoming a limiting factor for growth (Martin-Jézéquel et al., 2000). It has been shown that before silicate concentrations become growth limiting, diatoms first respond by thinning their frustules (McNair et al., 2018; Paasche, 1975). Whilst this phenomenon has been observed in several studies (McNair et al., 2018; Rocha et al., 2010; Shimada et al., 2009), our mechanistic understanding of how diatoms adjust their silicon quotas is unclear (Milligan et al., 2004). Interestingly, an initial thinning followed by subsequent thickening of diatom frustules has been observed at non-growth-limiting silicate concentrations (McNair et al., 2018). This may be an adaptive trade-off by which diatoms respond to lower silicate concentrations by decreasing their silicon quotas in favour of maintaining similar growth rates; a strategy that allows cells to respond to dynamic changes in silicate concentrations while maintaining a similar population size. Such rapid responses have been observed in both culture- and field-based experiments with diatoms responding to increases in silicate within several hours, while responses to nitrate additions, after prolonged nitrate stress, took over 30 h (McNair et al., 2018; Rocha et al., 2010). As such, it is possible that the low concentrations of  $\text{Si}(\text{OH})_4$  observed in the Ca-OAE treatment resulted in diatoms prioritising growth over silicate incorporation, leading to significantly less silicification when compared to the Si-OAE



**Figure 6.** Temporal variations of (a) biogenic silica (BSi) in the water column and (b) accumulation of BSi in the sediment trap across alkalinity source minerals and total alkalinity treatments during the extent of the experimental period. Nutrient addition on day 26 and 28 is represented by the dashed line dividing the experiment into phase I (pre-nutrient addition) and phase II (post-nutrient addition).

treatment. Detailed measurements of diatom growth (not assessed here) would be required to confirm this hypothesis. We recommend future experiments consider this and assess diatom growth alongside measures of silicification to enable the exploration of potential trade-offs between growth and silicification.

#### 4.3 Effect of carbonate chemistry manipulations on diatom silicification

We detected no clear relationship between total alkalinity and silicification in this mesocosm study, with the exception of *Pseudo-nitzschia*, which was more heavily silicified in higher-alkalinity treatments. *Nitzschia* also displayed increases in silicification in higher-alkalinity treatments; however, this was only observed in the Si-OAE treatment. In addition, we observed a significant effect of total alkalinity on silicification during phase I of the experiment. This result was primarily due to the Si-OAE-based treatment, illustrated by the significant three-way interaction between total alkalinity, alkalinity source mineral, and experimental phase in

the linear mixed-effect model. Previous physiological studies have found tight links between diatom silicification and components of the marine carbonate chemistry system ( $\text{CO}_2$  and pH) (Gao et al., 2014; Hervé et al., 2012; Li et al., 2019; Petrou et al., 2019; Zepernick et al., 2021). However, current research presents some inconsistencies in the relationship between silicification and carbonate chemistry. Petrou et al. (2019) found that silicification decreased at increased  $\text{pCO}_2$  and low pH expected as a result of ocean acidification. In contrast, research conducted by Li et al. (2019) and Zepernick et al. (2021) found the opposite with silicification decreasing at increasing pH or alkaline conditions. It is important to note that both Li et al. (2019) and Petrou et al. (2019) simulated ocean acidification, increasing  $\text{pCO}_2$  while total alkalinity remained constant. In contrast, Hervé et al. (2012) and Zepernick et al. (2021) altered the carbonate chemistry of their respective media via additions of NaOH and HCl thereby manipulating the concentrations of carbon species but not altering total DIC values. This allows for a more direct comparison to be made with the results presented here as OAE in its unequilibrated form results in changes in carbon species concentrations without significant differences in DIC. Hervé et al. (2012) found silica incorporation rates to decrease from pH 6.4 to 8.2 and increase from 8.2 to 8.5. Visual inspection of the data presented by Hervé et al. (2012) showed that the incorporation of silicate into a cell at pH 8.5 was marginally less than that at lower pH values (no statistics were provided for this measurement in the cited article). In support of this, Zepernick et al. (2021) found no significant difference in the silicification of freshwater diatoms at pH 7.7 and 8.6, only between pH 7.7 and 9.2 was a significant decline in silicification observed. In our study, total alkalinity values corresponded to a pH range from approximately 8.0 to 8.75 (total scale). Thus, it is possible that the enhancement of alkalinity in our study and corresponding changes in carbonate chemistry were not extreme enough to result in a significant change in the silicification of the diatom community or specific genera.

The pennate *Pseudo-nitzschia* was the only genus to exhibit a significant increase in silicification with increasing alkalinity irrespective of the mineral treatment type. In contrast, *Nitzschia* displayed a similar relationship between silicification and alkalinity; however, this was only in the Si-OAE based treatment. This relationship is similar to that observed by Petrou et al. (2019), who found several pennate species to exhibit a close relationship between silicification and carbonate chemistry conditions. Such a finding suggests that changes to silicification as a result of OAE is likely to be genus or species-specific supporting current knowledge surrounding the large variation in silicification between species (Martin-Jézéquel et al., 2000; Rousseau et al., 2002; Timmermans et al., 2004).

To conclude this section, we would like to highlight that other factors, which were not controlled for in this experiment, may have also contributed to the lack of a signifi-

cant difference in silicification observed here. Silicification is influenced by a range of environmental parameters, such as macronutrient concentrations (primarily silicate) (Claquin et al., 2002; Shimada et al., 2009), light intensity (Su et al., 2018; Taylor, 1985), and predation (Liu et al., 2016; Pondaven et al., 2007), which may have masked potential effects of total alkalinity on silicification. It is possible that OAE effects on silicification may be identifiable in other experimental settings with different boundary conditions and environmental controls. This is supported by the specific scenarios in which silicification was significantly influenced by total alkalinity here (i.e. during phase I of the experiment in the Si-OAE-based treatment). Our previous OAE study assessing the influence of a  $\sim 500 \mu\text{mol kg}^{-1}$  alkalinity increase on coastal Tasmanian plankton communities found significant effects of OAE on silicate dynamics, suggesting changes in diatom community silicification (Ferderer et al., 2022). These differences suggest that boundary conditions are important and that many studies assessing the effects of OAE on diatom communities will be needed to extract more robust response patterns across a range of conditions, consistent with conclusions drawn from a synthesis on diatoms in the context of ocean acidification (Bach and Taucher, 2019).

## 5 Implications for the implementation of OAE and outlook

In conclusion, our study underlines that the use of silicate-based minerals for OAE has the potential to significantly affect silicification of the diatom community and specific genera. This result was expected and consistent with our current understanding of dissolved silicate effects on diatom communities (Baines et al., 2010; Egge and Aksnes, 1992; Hauck et al., 2016; Tréguer et al., 2021). The significant influence of  $\text{Si}(\text{OH})_4$  on diatom silicification is not surprising, as it can be a limiting nutrient for diatoms, with previous studies having shown the benefits of increased  $\text{Si}(\text{OH})_4$  concentrations as a result of olivine-based mineral dissolution (Baines et al., 2010; Hutchins et al., 2023; Martin-Jézéquel et al., 2000; Wischmeyer et al., 2003). It is important to note that this experiment was designed to specifically assess the potential interactive effects of OAE and magnesium silicate fertilisation associated with silicate based minerals proposed for use in mineral based OAE. Such an interaction was explored as there are several potential minerals proposed for use in mineral-based OAE, all of which vary in their respective dissolution products. As such we did not control for or assess other ecologically important dissolution products found in silicate-based minerals, e.g. iron and nickel, which have been shown to have significant ecological implications for plankton communities (Xin et al., 2023; Hutchins et al., 2023; Guo et al., 2022; Boyd et al., 2007). Further exploration of potential dissolution products and their effects on plankton com-

munities in the presence and absence of changing carbonate chemistry conditions are encouraged in future research.

In contrast to the clear effects of dissolved silicate fertilisation on silicification, our findings provide limited evidence to suggest that the enhancement of seawater alkalinity by  $0\text{--}600 \mu\text{mol kg}^{-1}$  affects genera-specific silicification. The lack of a clear alkalinity effect on silicification was unexpected, especially in the higher-alkalinity treatments, which corresponded to a substantial change in carbonate chemistry conditions. It is important to note that increases in alkalinity above  $\Delta 400 \mu\text{mol kg}^{-1}$  are considered to be relatively extreme levels of OAE, yet only two genera, *Pseudo-nitzschia* and *Nitzschia*, exhibited significant changes in silicification as a result of the gradient in carbonate chemistry employed here (Schulz et al., 2023). Real-world applications are predicted to employ significantly less extreme perturbations of the marine carbonate chemistry system (apart from sites of direct alkalinity addition). As such, one might hypothesise that impacts may be less significant. Furthermore, such perturbations would be relatively short-lived in real-world applications since dilution of the perturbed waterbodies occurs, unlike the sustained changes observed within mesocosms presented here (He and Tyka, 2023; Wang et al., 2023). Irrespective of this, there is substantial empirical evidence suggesting that changes in carbonate chemistry through ocean acidification will influence diatom communities, their growth, various aspects of silicification (e.g. rate, degree) and subsequently silicate cycling in the ocean (Bach and Taucher, 2019; Gao et al., 2014; Li et al., 2019; Milligan et al., 2004; Petrou et al., 2019; Taucher et al., 2022). Additionally, our previous work has shown that an increase in total alkalinity of  $\sim 500 \mu\text{mol kg}^{-1}$  has a significant influence on the uptake of dissolved silicate and production of BSi in a coastal phytoplankton community (Ferderer et al., 2022). The mixed outcomes observed here and in the limited OAE studies so far suggest that the responses of diatoms will differ and be dependent on the community and environmental boundary conditions. More community studies, ideally with closely aligned experimental setups, will be needed to discern whether the response of diatoms to OAE forms any robust patterns. Such ecological observations subsequently need mechanistic underpinning, potentially achievable through the intelligent design of physiological experiments (Collins et al., 2022). Ultimately, the goal should be to provide predictive understanding of the role of diatoms and eventually all major functional plankton groups under the differing strategies of OAE.

*Data availability.* Data are available from the Institute for Marine and Antarctic Studies (IMAS) data catalogue, University of Tasmania (UTAS) <https://doi.org/10.25959/G3FN-HE45> (Ferderer, 2023).



*Supplement.* The supplement related to this article is available online at: <https://doi.org/10.5194/bg-21-2777-2024-supplement>.

*Author contributions.* UR designed the mesocosm experiment, and AF designed the experiment assessing silicification rates, with input from LTB and KGB. AF was responsible for the investigation, data curation, formal analysis, and writing. AF wrote the manuscript with contributions from LTB, KGS, UR, KGB, and ZC.

*Competing interests.* The contact author has declared that none of the authors has any competing interests.

*Disclaimer.* Publisher's note: Copernicus Publications remains neutral with regard to jurisdictional claims made in the text, published maps, institutional affiliations, or any other geographical representation in this paper. While Copernicus Publications makes every effort to include appropriate place names, the final responsibility lies with the authors.

*Special issue statement.* This article is part of the special issue “Environmental impacts of ocean alkalinity enhancement”. It is not associated with a conference.

*Acknowledgements.* We would like to thank all participants of the campaign who assisted in sampling and general scientific discussion throughout. We would also like to acknowledge Juliane Tammen and Peter Fritzsche for the measurement of dissolved inorganic nutrients, Leila Kittu, Anna Groen, Lucas Krause, Jule Ploschke, and Kira Lange for their measurements of particulate matter; Jana Meyer for the measurement of sedimented material; Julieta Schneider for the measurement of carbonate chemistry; Andrea Ludwig for her organisation and management of the project; and the staff members at the research station for their assistance in daily tasks.

*Financial support.* This study was funded by the OceanNETS project (“Ocean-based Negative Emissions Technologies – analysing the feasibility, risks and co-benefits of ocean-based negative emission technologies for stabilizing the climate”, EU Horizon 2020 Research and Innovation Programme grant agreement no. 869357), and the Helmholtz European Partnering project Ocean-CDR (“Ocean-based carbon dioxide removal strategies”, project no. PIE-0021) with additional support from the AQUACOSM-plus project (EU H2020-INFRAIA project no. 871081, “AQUACOSM-plus: Network of Leading European AQUATIC MesoCOSM Facilities Connecting Rivers, Lakes, Estuaries and Oceans in Europe and beyond”). Additional funding was supplied by a Future Fellowship (FT200100846) awarded to Lennart T. Bach by the Australian Research Council. This research was also conducted while Aaron Ferderer was in receipt of an Australian Government Research Training Program (RTP) scholarship.

*Review statement.* This paper was edited by Lydia Kapsenberg and reviewed by two anonymous referees.

## References

- Albright, R., Caldeira, L., Hosfelt, J., Kwiatkowski, L., Maclaren, J. K., Mason, B. M., Nebuchina, Y., Ninokawa, A., Pongratz, J., Ricke, K. L., Rivlin, T., Schneider, K., Sesboüé, M., Shamberger, K., Silverman, J., Wolfe, K., Zhu, K., and Caldeira, K.: Reversal of ocean acidification enhances net coral reef calcification, *Nature*, 531, 362–365, <https://doi.org/10.1038/nature17155>, 2016.
- Bach, L. T. and Taucher, J.: CO<sub>2</sub> effects on diatoms: a synthesis of more than a decade of ocean acidification experiments with natural communities, *Ocean Sci.*, 15, 1159–1175, <https://doi.org/10.5194/os-15-1159-2019>, 2019.
- Bach, L. T., Riebesell, U., Gutowska, M. A., Federwisch, L., and Schulz, K. G.: A unifying concept of coccolithophore sensitivity to changing carbonate chemistry embedded in an ecological framework, *Prog. Oceanogr.*, 135, 125–138, <https://doi.org/10.1016/j.pocean.2015.04.012>, 2015.
- Bach, L. T., Gill, S. J., Rickaby, R. E. M., Gore, S., and Renforth, P.: CO<sub>2</sub> Removal With Enhanced Weathering and Ocean Alkalinity Enhancement: Potential Risks and Co-benefits for Marine Pelagic Ecosystems, *Front. Clim.*, 1, 7, <https://doi.org/10.3389/FCLIM.2019.00007>, 2019.
- Baines, S. B., Twining, B. S., Brzezinski, M. A., Nelson, D. M., and Fisher, N. S.: Causes and biogeochemical implications of regional differences in silicification of marine diatoms, *Global Biogeochem. Cy.*, 24, GB4031, <https://doi.org/10.1029/2010GB003856>, 2010.
- Boxhammer, T., Bach, L. T., Czerny, J., and Riebesell, U.: Technical note: Sampling and processing of mesocosm sediment trap material for quantitative biogeochemical analysis, *Biogeosciences*, 13, 2849–2858, <https://doi.org/10.5194/bg-13-2849-2016>, 2016.
- Boyd, P. W., Jickells, T., Law, C. S., Blain, S., Boyle, E. A., Buesseler, K. O., Coale, K. H., Cullen, J. J., de Baar, H. J. W., Follows, M., Harvey, M., Lancelot, C., Levasseur, M., Owens, N. P. J., Pollard, R., Rivkin, R. B., Sarmiento, J., Schoemann, V., Smetacek, V., Takeda, S., Tsuda, A., Turner, S., and Watson, A. J.: Mesoscale iron enrichment experiments 1993–2005: synthesis and future directions, *Science*, 315, 612–617, <https://doi.org/10.1126/science.1131669>, 2007.
- Burt, D. J., Fröb, F., and Ilyina, T.: The Sensitivity of the Marine Carbonate System to Regional Ocean Alkalinity Enhancement, *Front. Clim.*, 3, 1–15, <https://doi.org/10.3389/fclim.2021.624075>, 2021.
- Chen, C. Y. and Durbin, E. G.: Effects of pH on the growth and carbon uptake of marine phytoplankton, *Mar. Ecol. Prog. Ser.*, 109, 83–94, <https://doi.org/10.3354/meps109083>, 1994.
- Claquin, P., Martin-Jézéquel, V., Kromkamp, J. C., Veldhuis, M. J. W., and Kraay, G. W.: Uncoupling of Silicon Compared with Carbon and Nitrogen Metabolisms and the Role of the Cell Cycle in Continuous Cultures of *Thalassiosira Pseudonana* (bacillariophyceae) Under Light, Nitrogen, and Phosphorus Control, *J. Phycol.*, 38, 922–930, <https://doi.org/10.1046/j.1529-8817.2002.t01-1-01220.x>, 2002.
- Collins, S., Whittaker, H., and Thomas, M. K.: The need for unrealistic experiments in global change biology, *Curr. Opin. Micro-*

- biol., 68, 102151, <https://doi.org/10.1016/J.MIB.2022.102151>, 2022.
- Czerny, J., Schulz, K. G., Krug, S. A., Ludwig, A., and Riebesell, U.: Technical Note: The determination of enclosed water volume in large flexible-wall mesocosms “KOSMOS”, *Biogeosciences*, 10, 1937–1941, <https://doi.org/10.5194/bg-10-1937-2013>, 2013.
- Dickson, A. G., Sabine, C. L., and Christian, J. R.: Guide to best practices for ocean CO<sub>2</sub> measurements, PICES Special Publication, 3, 0–199, 2007.
- EGGE, J. and AKSNES, D.: Silicate as regulating nutrient in phytoplankton competition, *Mar. Ecol. Prog. Ser.*, 83, 281–289, <https://doi.org/10.3354/meps083281>, 1992.
- Ehlert, C., Reckhardt, A., Greskowiak, J., Liguori, B. T. P., Böning, P., Paffrath, R., Brumsack, H.-J., and Pahnke, K.: Transformation of silicon in a sandy beach ecosystem: Insights from stable silicon isotopes from fresh and saline groundwaters, *Chem. Geol.*, 440, 207–218, <https://doi.org/10.1016/j.chemgeo.2016.07.015>, 2016.
- Eisaman, M. D., Geilert, S., Renforth, P., Bastianini, L., Campbell, J., Dale, A. W., Foteinis, S., Grasse, P., Hawrot, O., Löscher, C. R., Rau, G. H., and Rønning, J.: Assessing the technical aspects of ocean-alkalinity-enhancement approaches, in: Guide to Best Practices in Ocean Alkalinity Enhancement Research, edited by: Oschlies, A., Stevenson, A., Bach, L. T., Fennel, K., Rickaby, R. E. M., Satterfield, T., Webb, R., and Gattuso, J.-P., Copernicus Publications, State Planet, 2-oea2023, 3, <https://doi.org/10.5194/sp-2-oea2023-3-2023>, 2023.
- Escaravage, V. and Prins, T. C.: Silicate availability, vertical mixing and grazing control of phytoplankton blooms in mesocosms, in: Sustainable Increase of Marine Harvesting: Fundamental Mechanisms and New Concepts, edited by: Vadstein, O. and Olsen, Y., Springer Netherlands, Dordrecht, 33–48, [https://doi.org/10.1007/978-94-017-3190-4\\_4](https://doi.org/10.1007/978-94-017-3190-4_4), 2002.
- Feng, E. Y., Koeve, W., Keller, D. P., and Oschlies, A.: Model-Based Assessment of the CO<sub>2</sub> Sequestration Potential of Coastal Ocean Alkalinization, *Earth’s Future*, 5, 1252–1266, <https://doi.org/10.1002/2017EF000659>, 2017.
- Ferderer, A.: Investigating the effect of silicate- and calcium-based ocean alkalinity enhancement on diatom silicification – data, Institute for Marine and Antarctic Studies (IMAS), University of Tasmania (UTAS) [data set], <https://doi.org/10.25959/G3FN-HE45>, 2023.
- Ferderer, A., Chase, Z., Kennedy, F., Schulz, K. G., and Bach, L. T.: Assessing the influence of ocean alkalinity enhancement on a coastal phytoplankton community, *Biogeosciences*, 19, 5375–5399, <https://doi.org/10.5194/bg-19-5375-2022>, 2022.
- Fuss, S., Lamb, W. F., Callaghan, M. W., Hilaire, J., Creutzig, F., Amann, T., Beringer, T., De Oliveira Garcia, W., Hartmann, J., Khanna, T., Luderer, G., Nemet, G. F., Rogelj, J., Smith, P., Vicente, J. V., Wilcox, J., Del Mar Zamora Dominguez, M., and Minx, J. C.: Negative emissions—Part 2: Costs, potentials and side effects, *Environ. Res. Lett.*, 13, 063002, <https://doi.org/10.1088/1748-9326/AABF9F>, 2018.
- Gao, K., Campbell, D. A., Gao, K., and Campbell, D. A.: Physiological responses of marine diatoms to elevated CO<sub>2</sub> and decreased pH: a review, *Funct. Plant Biol.*, 41, 449–459, <https://doi.org/10.1071/FP13247>, 2014.
- Gately, J. A., Kim, S. M., Jin, B., Brzezinski, M. A., and Iglesias-Rodriguez, M. D.: Coccolithophores and diatoms resilient to ocean alkalinity enhancement: A glimpse of hope?, *Science Advances*, 9, eadg6066, <https://doi.org/10.1126/sciadv.adg6066>, 2023.
- Gore, S., Renforth, P., and Perkins, R.: The potential environmental response to increasing ocean alkalinity for negative emissions, *Mitig. Adapt. Strat. Gl.*, 24, 1191–1211, <https://doi.org/10.1007/s11027-018-9830-z>, 2019.
- Goto, K.: Effect of pH on Polymerization of Silicic Acid, *J. Phys. Chem.*, 60, 1007–1008, <https://doi.org/10.1021/j150541a046>, 1956.
- Guo, J. A., Strzepek, R., Willis, A., Ferderer, A., and Bach, L. T.: Investigating the effect of nickel concentration on phytoplankton growth to assess potential side-effects of ocean alkalinity enhancement, *Biogeosciences*, 19, 3683–3697, <https://doi.org/10.5194/bg-19-3683-2022>, 2022.
- Hansen, H. P. and Koroleff, F.: Determination of nutrients, in: Methods of Seawater Analysis, John Wiley & Sons, Ltd, 159–228, <https://doi.org/10.1002/9783527613984.ch10>, 1999.
- Hartmann, J., West, A. J., Renforth, P., Köhler, P., Rocha, C. L. D. L., Wolf-Gladrow, D. A., Dürr, H. H., and Scheffran, J.: Enhanced chemical weathering as a geoengineering strategy to reduce atmospheric carbon dioxide, supply nutrients, and mitigate ocean acidification, *Rev. Geophys.*, 51, 113–149, <https://doi.org/10.1002/ROG.20004>, 2013.
- Hauck, J., Köhler, P., Wolf-Gladrow, D., and Völker, C.: Iron fertilisation and century-scale effects of open ocean dissolution of olivine in a simulated CO<sub>2</sub> removal experiment, *Environ. Res. Lett.*, 11, 024007, <https://doi.org/10.1088/1748-9326/11/2/024007>, 2016.
- He, J. and Tyka, M. D.: Limits and CO<sub>2</sub> equilibration of near-coast alkalinity enhancement, *Biogeosciences*, 20, 27–43, <https://doi.org/10.5194/bg-20-27-2023>, 2023.
- Hervé, V., Derr, J., Douady, S., Quinet, M., Moisan, L., and Lopez, P. J.: Multiparametric Analyses Reveal the pH-Dependence of Silicon Biomineralization in Diatoms, *PLOS ONE*, 7, e46722, <https://doi.org/10.1371/JOURNAL.PONE.0046722>, 2012.
- Hinga, K. R.: Effects of pH on coastal marine phytoplankton, *Mar. Ecol. Prog. Ser.*, 238, 281–300, <https://doi.org/10.3354/meps238281>, 2002.
- Hutchins, D. A., Fu, F.-X., Yang, S.-C., John, S. G., Romaniello, S. J., Andrews, M. G., and Walworth, N. G.: Responses of keystone phytoplankton groups to olivine dissolution products and implications for carbon dioxide removal via ocean alkalinity enhancement, *bioRxiv*, 2023.04.08.536121, <https://doi.org/10.1101/2023.04.08.536121>, 2023.
- Ilyina, T., Wolf-Gladrow, D., Munhoven, G., and Heinze, C.: Assessing the potential of calcium-based artificial ocean alkalinization to mitigate rising atmospheric CO<sub>2</sub> and ocean acidification, *Geophys. Res. Lett.*, 40, 5909–5914, <https://doi.org/10.1002/2013GL057981>, 2013.
- IPCC: Summary for Policymakers, in: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change, edited by: Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S. L., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M. I., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J. B. R., Maycock, T. K., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., Cambridge University Press,

- Cambridge, United Kingdom and New York, NY, USA, 3–32, <https://doi.org/10.1017/9781009157896.001>, 2021.
- Keller, D. P., Feng, E. Y., and Oschlies, A.: Potential climate engineering effectiveness and side effects during a high carbon dioxide-emission scenario, *Nat. Commun.*, 5, 3304, <https://doi.org/10.1038/ncomms4304>, 2014.
- Kheshgi, H. S.: Sequestering atmospheric carbon dioxide by increasing ocean alkalinity, *Energy*, 20, 915–922, [https://doi.org/10.1016/0360-5442\(95\)00035-F](https://doi.org/10.1016/0360-5442(95)00035-F), 1995.
- Leblanc, K. and Hutchins, D. A.: New applications of a biogenic silica deposition fluorophore in the study of oceanic diatoms, *Limnol. Oceanogr.-Meth.*, 3, 462–476, <https://doi.org/10.4319/LOM.2005.3.462>, 2005.
- Lenth, R., Buerkner, P., Herve, M., Love, J., Riebl, H., and Singmann, H.: emmeans: estimated marginal means, aka Least-Squares Means, R package version 1.7.0, <https://CRAN.R-project.org/package=emmeans> (last access: 11 June 2024), 2023.
- Li, F., Fan, J., Hu, L., Beardall, J., Xu, J., and Fields, D.: Physiological and biochemical responses of *Thalassiosira weissflogii* (diatom) to seawater acidification and alkalization, *ICES J. Mar. Sci.*, 76, 1850–1859, <https://doi.org/10.1093/ICESJMS/FSZ028>, 2019.
- Liu, H., Chen, M., Zhu, F., and Harrison, P. J.: Effect of Diatom Silica Content on Copepod Grazing, Growth and Reproduction, *Frontiers in Marine Science*, 3, 89, <https://doi.org/10.3389/fmars.2016.00089>, 2016.
- Martin-Jézéquel, V., Hildebrand, M., and Brzezinski, M. A.: Silicon metabolism in diatoms: Implications for growth, *J. Phycol.*, 36, 821–840, <https://doi.org/10.1046/J.1529-8817.2000.00019.X>, 2000.
- McNair, H. M., Brzezinski, M. A., and Krause, J. W.: Quantifying diatom silicification with the fluorescent dye, PDMPPO, *Limnol. Oceanogr.-Meth.*, 13, 587–599, <https://doi.org/10.1002/LOM3.10049>, 2015.
- McNair, H. M., Brzezinski, M. A., and Krause, J. W.: Diatom populations in an upwelling environment decrease silica content to avoid growth limitation, *Environ. Microbiol.*, 20, 4184–4193, <https://doi.org/10.1111/1462-2920.14431>, 2018.
- Milligan, A. J., Varela, D. E., Brzezinski, M. A., and Morel, F. M. M.: Dynamics of silicon metabolism and silicon isotopic discrimination in a marine diatom as a function of pCO<sub>2</sub>, *Limnol. Oceanogr.*, 49, 322–329, <https://doi.org/10.4319/LO.2004.49.2.0322>, 2004.
- Okamoto, G., Okura, T., and Goto, K.: Properties of silica in water, *Geochim. Cosmochim. Ac.*, 12, 123–132, [https://doi.org/10.1016/0016-7037\(57\)90023-6](https://doi.org/10.1016/0016-7037(57)90023-6), 1957.
- Owen, L.: Precipitation of amorphous silica from high-temperature hypersaline geothermal brines, California Univ., Livermore, USA, Lawrence Livermore Lab., 1975.
- Paasche, E.: Growth of the plankton diatom *Thalassiosira nordenskiöldii* Cleve at low silicate concentrations, *J. Exp. Mar. Biol. Ecol.*, 18, 173–183, [https://doi.org/10.1016/0022-0981\(75\)90072-6](https://doi.org/10.1016/0022-0981(75)90072-6), 1975.
- Paquay, F. S. and Zeebe, R. E.: Assessing possible consequences of ocean liming on ocean pH, atmospheric CO<sub>2</sub> concentration and associated costs, *Int. J. Greenh. Gas Con.*, 17, 183–188, <https://doi.org/10.1016/j.ijggc.2013.05.005>, 2013.
- Paul, A. J. and Bach, L. T.: Universal response pattern of phytoplankton growth rates to increasing CO<sub>2</sub>, *New Phytol.*, 228, 1710–1716, <https://doi.org/10.1111/NPH.16806>, 2020.
- Petrou, K., Baker, K. G., Nielsen, D. A., Hancock, A. M., Schulz, K. G., and Davidson, A. T.: Acidification diminishes diatom silica production in the Southern Ocean, *Nat. Climate Change*, 9, 781–786, <https://doi.org/10.1038/S41558-019-0557-Y>, 2019.
- Pondaven, P., Gallinari, M., Chollet, S., Bucciarelli, E., Sarthou, G., Schultes, S., and Jean, F.: Grazing-induced Changes in Cell Wall Silicification in a Marine Diatom, *Protist*, 158, 21–28, <https://doi.org/10.1016/J.PROTIS.2006.09.002>, 2007.
- R Core Team: R: A Language and Environment for Statistical Computing, R Foundation for Statistical Computing, Vienna, Austria, <https://www.R-project.org> (last access: 11 June 2024), 2023.
- Renforth, P. and Henderson, G.: Assessing ocean alkalinity for carbon sequestration, *Rev. Geophys.*, 55, 636–674, <https://doi.org/10.1002/2016RG000533>, 2017.
- Riebesell, U., Wolf-Gladrow, D. A., and Smetacek, V.: Carbon dioxide limitation of marine phytoplankton growth rates, *Nature*, 361, 249–251, <https://doi.org/10.1038/361249a0>, 1993.
- Riebesell, U., Czerny, J., von Bröckel, K., Boxhammer, T., Büdenbender, J., Deckelnick, M., Fischer, M., Hoffmann, D., Krug, S. A., Lentz, U., Ludwig, A., Mucche, R., and Schulz, K. G.: Technical Note: A mobile sea-going mesocosm system – new opportunities for ocean change research, *Biogeosciences*, 10, 1835–1847, <https://doi.org/10.5194/bg-10-1835-2013>, 2013.
- Rocha, C. L. D. L., Terbrüggen, A., Völker, C., and Hohn, S.: Response to and recovery from nitrogen and silicon starvation in *Thalassiosira weissflogii*: growth rates, nutrient uptake and C, Si and N content per cell, *Mar. Ecol. Prog. Ser.*, 412, 57–68, <https://doi.org/10.3354/meps08701>, 2010.
- Rousseau, V., Leynaert, A., Daoud, N., and Lancelot, C.: Diatom succession, silicification and silicic acid availability in Belgian coastal waters (Southern North Sea), *Mar. Ecol. Prog. Ser.*, 236, 61–73, <https://doi.org/10.3354/MEPS236061>, 2002.
- Schulz, K. G., Bach, L. T., and Dickson, A. G.: Seawater carbonate chemistry considerations for ocean alkalinity enhancement research: theory, measurements, and calculations, in: *Guide to Best Practices in Ocean Alkalinity Enhancement Research*, edited by: Oschlies, A., Stevenson, A., Bach, L. T., Fennel, K., Rickaby, R. E. M., Satterfield, T., Webb, R., and Gattuso, J.-P., Copernicus Publications, State Planet, 2-oae2023, 2, <https://doi.org/10.5194/sp-2-oae2023-2-2023>, 2023.
- Shimada, C., Nakamachi, M., Tanaka, Y., Yamasaki, M., and Kuwata, A.: Effects of nutrients on diatom skeletal silicification: Evidence from *Neodenticula seminiae* culture experiments and morphometric analysis, *Mar. Micropaleontol.*, 73, 164–177, <https://doi.org/10.1016/J.MARMICRO.2009.09.001>, 2009.
- Spinthaki, A., Petratos, G., Matheis, J., Hater, W., and Demadis, K. D.: The precipitation of “magnesium silicate” under geothermal stresses. Formation and characterization, *Geothermics*, 74, 172–180, <https://doi.org/10.1016/j.geothermics.2018.03.001>, 2018.
- Su, Y., Lundholm, N., and Ellegaard, M.: The effect of different light regimes on diatom frustule silicon concentration, *Algal Res.*, 29, 36–40, <https://doi.org/10.1016/j.algal.2017.11.014>, 2018.
- Taucher, J., Bach, L. T., Prowe, A. E. F., Boxhammer, T., Kvale, K., and Riebesell, U.: Enhanced silica export in a future

- ocean triggers global diatom decline, *Nature*, 605, 696–700, <https://doi.org/10.1038/s41586-022-04687-0>, 2022.
- Taylor, N. J.: Silica incorporation in the diatom *Coscinodiscus granii* as affected by light intensity, *Brit. Phycol. J.*, 20, 365–374, <https://doi.org/10.1080/00071618500650371>, 1985.
- Timmermans, K. R., Van Der Wagt, B., and De Baar, H. J. W.: Growth rates, half-saturation constants, and silicate, nitrate, and phosphate depletion in relation to iron availability of four large, open-ocean diatoms from the Southern Ocean, *Limnol. Oceanogr.*, 49, 2141–2151, <https://doi.org/10.4319/LO.2004.49.6.2141>, 2004.
- Tréguer, P. J., Sutton, J. N., Brzezinski, M., Charette, M. A., Devries, T., Dutkiewicz, S., Ehlert, C., Hawkings, J., Leynaert, A., Liu, S. M., Monferrer, N. L., López-Acosta, M., Maldonado, M., Rahman, S., Ran, L., and Rouxel, O.: Reviews and syntheses: The biogeochemical cycle of silicon in the modern ocean, *Biogeosciences*, 18, 1269–1289, <https://doi.org/10.5194/bg-18-1269-2021>, 2021.
- Van Vuuren, D. P., Stehfest, E., Gernaat, D. E. H. J., Van Den Berg, M., Bijl, D. L., De Boer, H. S., Daioglou, V., Doelman, J. C., Edelenbosch, O. Y., Harmsen, M., Hof, A. F., and Van Sluisveld, M. A. E.: Alternative pathways to the 1.5 °C target reduce the need for negative emission technologies, *Nat. Clim. Change*, 8, 391–397, <https://doi.org/10.1038/S41558-018-0119-8>, 2018.
- Wang, F., Lu, Z., Wang, Y., Yan, R., and Chen, N.: Porewater exchange drives the dissolved silicate export across the wetland-estuarine continuum, *Frontiers in Marine Science*, 10, 812, <https://doi.org/10.3389/FMARS.2023.1206776>, 2023.
- Welschmeyer, N. A.: Fluorometric analysis of chlorophyll *a* in the presence of chlorophyll *b* and pheopigments, *Limnol. Oceanogr.*, 39, 1985–1992, <https://doi.org/10.4319/lo.1994.39.8.1985>, 1994.
- Wischmeyer, A. G., Del Amo, Y., Brzezinski, M., and Wolf-Gladrow, D. A.: Theoretical constraints on the uptake of silicic acid species by marine diatoms, *Mar. Chem.*, 82, 13–29, [https://doi.org/10.1016/S0304-4203\(03\)00033-1](https://doi.org/10.1016/S0304-4203(03)00033-1), 2003.
- Xin, X., Faucher, G., and Riebesell, U.: Phytoplankton response to increased nickel in the context of ocean alkalinity enhancement, *Biogeosciences*, 21, 761–772, <https://doi.org/10.5194/bg-21-761-2024>, 2024.
- Zepernick, B. N., Gann, E. R., Martin, R. M., Pound, H. L., Krausfeldt, L. E., Chaffin, J. D., and Wilhelm, S. W.: Elevated pH Conditions Associated With *Microcystis* spp. Blooms Decrease Viability of the Cultured Diatom *Fragilaria crotonensis* and Natural Diatoms in Lake Erie, *Front. Microbiol.*, 12, 598736, <https://doi.org/10.3389/fmicb.2021.598736>, 2021.
- Znachor, P. and Nedoma, J.: Application of the pdmpo technique in studying silica deposition in natural populations of *fragilaria crotonensis* (bacillariophyceae) at different depths in a eutrophic reservoir1, *J. Phycol.*, 44, 518–525, <https://doi.org/10.1111/J.1529-8817.2008.00470.X>, 2008.