

## Nitrous oxide emissions from the upwelling area off Mauritania (NW Africa)

Franziska Wittke,<sup>1</sup> Annette Kock,<sup>1</sup> and Hermann W. Bange<sup>1</sup>

Received 8 January 2010; revised 23 April 2010; accepted 3 May 2010; published 16 June 2010.

[1] Nitrous oxide (N<sub>2</sub>O) flux densities across the ocean/atmosphere interface from the Mauritanian upwelling (16°–18.5°W, 16°–21°N) were computed with a simple upwelling model using N<sub>2</sub>O measurements from four cruises between 2006 and 2008 as well as wind data from the QuikSCAT satellite. The resulting N<sub>2</sub>O flux densities show a strong seasonality reflecting the wind-driven seasonality of the upwelling: N<sub>2</sub>O flux densities are highest in the northern part (19.5°–21°N) and show a decreasing trend towards the south. The summer periods with no upwelling (and thus associated with no or negligible N<sub>2</sub>O flux densities) are most pronounced in the southern part (16°–17°N). The mean seasonally and regionally weighted annual N<sub>2</sub>O emissions from the Mauritanian upwelling were estimated to 1.0 Gg N. This is low compared to other major upwelling areas (Arabian Sea, off Chile) indicating that N<sub>2</sub>O emissions from the Mauritanian upwelling are a minor source of atmospheric N<sub>2</sub>O. **Citation:** Wittke, F., A. Kock, and H. W. Bange (2010), Nitrous oxide emissions from the upwelling area off Mauritania (NW Africa), *Geophys. Res. Lett.*, 37, L12601, doi:10.1029/2010GL042442.

### 1. Introduction

[2] Nitrous oxide (N<sub>2</sub>O) is a gaseous trace component of the Earth's atmosphere and contributes significantly to both the greenhouse effect and the stratospheric ozone depletion [Forster *et al.*, 2007; Ravishankara *et al.*, 2009]. Main formation pathways of N<sub>2</sub>O are microbial nitrification and denitrification leading to an accumulation of N<sub>2</sub>O in oceanic subsurface layers [see, e.g., Bange, 2008]. The world's oceans, including their coastal zones, are a major source for N<sub>2</sub>O and contribute about 30% to the global atmospheric N<sub>2</sub>O budget [Bange, 2006; Denman *et al.*, 2007]. The major coastal upwelling areas such as found in the Arabian Sea and in the eastern tropical North and South Pacific are well-known as sites of enhanced N<sub>2</sub>O emissions to the atmosphere [Bange *et al.*, 2001; Cornejo *et al.*, 2006; Lueker *et al.*, 2003]. Coastal upwelling areas may contribute about 1–5% to the overall oceanic N<sub>2</sub>O emissions [Bange *et al.*, 1996a; Nevison *et al.*, 2004]. However, detailed studies of N<sub>2</sub>O emissions from major upwelling areas such as off NW Africa (Morocco/Mauritania) are missing. Based on the 'wind-only' upwelling model approach by Nevison *et al.* [2004] and N<sub>2</sub>O measurements from four cruises to the

coastal area off Mauritania between 2006 and 2008, we present a first regionally and seasonally weighted estimate of N<sub>2</sub>O emissions from the coastal upwelling off Mauritania (21°–16°N).

### 2. Study Site Description

[3] The eastern tropical North Atlantic Ocean is an area with strongly differing hydrographic and biological properties: On the one hand, the Canary and North Equatorial Currents form the eastern part of the North Atlantic Subtropical Gyre which is a zone of low nutrient-supply and thus low primary production. On the other hand nutrient-rich upwelled water masses feed zones of high biological productivity off the Northwest African coast. Upwelling events off Mauritania are characterized by a pronounced drop in sea surface temperatures (SST). SST of <17°C are typical for upwelled water masses along the coast [Mittelstaedt, 1986]. The coastal upwelling occurs within a narrow (10–20 km) band along the continental slope as a consequence of offshore Ekman transport due to the trade winds along the Mauritanian coastline. The upwelled water is coming from a water depth of about 100–200 m [Hagen, 2001; Mittelstaedt, 1986]. The occurrence of the upwelling shows a seasonality following the shifting of the Inter-Tropical Convergence Zone (ITCZ) throughout the year [Hagen, 2001]. In the region between Cap Vert (~15°N) and Cap Blanc (~21°N) seasonal upwelling takes place during winter/spring with the southernmost extension in February [Nykjer and Van Camp, 1994; Schemainda *et al.*, 1975; Wooster *et al.*, 1976].

### 3. Methods

[4] N<sub>2</sub>O flux densities (F<sub>N<sub>2</sub>O</sub>) across the ocean/atmosphere interface were calculated as

$$F_{N_2O} = k_w(C_w - C_a), \quad (1)$$

where  $k_w$  is the gas exchange coefficient and  $C_w$  is the N<sub>2</sub>O concentration in the ocean surface layer.  $C_a$  is the estimated N<sub>2</sub>O equilibrium concentration and was calculated as

$$C_a = \beta x' P, \quad (2)$$

where  $\beta$  is the Bunsen solubility of N<sub>2</sub>O in seawater [Weiss and Price, 1980].  $\beta$  was calculated with monthly surface temperatures and salinities from the World Ocean Atlas 2005 [Antonov *et al.*, 2006; Locarnini *et al.*, 2006].  $P$  is the ambient pressure (set to 1 atm) and  $x'$  is the dry mole fraction in the ambient air. For  $x'$  we used the monthly time series of atmospheric N<sub>2</sub>O from the Advanced Global Atmospheric Gases Experiment (AGAGE) monitoring station Ragged

<sup>1</sup>Forschungsbereich Marine Biogeochemie, Leibniz-Institut für Meereswissenschaften an der Universität Kiel (IFM-GEOMAR), Kiel, Germany.

**Table 1.** Concentrations of N<sub>2</sub>O in 100 m (C<sub>100m</sub>) From Four Cruises Between 2006 and 2008

Latitude (°N)	Longitude (°W)	Cruise	Date	C <sub>100m</sub> ± 1 SD <sup>a</sup> (nmol L <sup>-1</sup> )
18.0	17.0	M68/3	Jul 2006	22.7 ± 1.2
17.5	17.0	M68/3	Jul 2006	35.8 ± 0.3
18.0	16.5	M68/3	Jul 2006	35.7 ± 1.3
18.5	16.8	P347	Jan 2007	22.6 ± 2.5
17.5	16.5	P347	Jan 2007	31.7 ± 0.7
18.0	17.0	P348	Feb 2007	27.1 ± 0.6
19.0	17.0	P348	Feb 2007	32.5 ± 0.2
18.0	16.5	P348	Feb 2007	27.6 ± 0.1
20.0	18.0	ATA3	Feb 2008	28.0 ± 0.3
20.1	17.7	ATA3	Feb 2008	21.9 ± 0.8
16.2	17.1	ATA3	Feb 2008	28.3 ± 0.2
18.0	17.0	ATA3	Feb 2008	24.8 ± 0.7
17.0	17.0	ATA3	Feb 2008	31.2 ± 0.9
19.0	16.9	ATA3	Feb 2008	26.9 ± 0.4
17.0	16.8	ATA3	Feb 2008	33.5 ± 0.9
17.8	16.6	ATA3	Feb 2008	31.5 ± 0.9
17.8	16.6	ATA3	Feb 2008	30.8 ± 0.5

<sup>a</sup>SD stands for (estimated) standard deviation; for details of the calculation see *Walter et al.* [2006].

Point on Barbados (see <http://agage.eas.gatech.edu>) [*Prinn et al.*, 2000]. For the calculation of k<sub>w</sub> we used the approach of *Nightingale et al.* [2000]:

$$k_w = 0.01/3600(0.222u^2 + 0.333u)(Sc_{N_2O}/600)^{-n}, \quad (3)$$

where  $u$  is the wind speed in 10 m height,  $Sc_{N_2O}$  is the Schmidt number of N<sub>2</sub>O and  $n$  equals  $-0.66$  for  $0 < u < 3.6$  m s<sup>-1</sup> or  $n = -0.5$  for  $u > 3.6$  m s<sup>-1</sup>.  $Sc_{N_2O}$  was calculated as the ratio of the kinematic viscosity of seawater and the diffusion of N<sub>2</sub>O in water [see *Walter et al.*, 2004, and references therein].  $C_w$  was calculated following the ‘wind-only’ upwelling model approach described by *Nevison et al.* [2004]:

$$C_w = C_{100m}Q_wQ_t, \quad (4)$$

where  $C_{100m}$  is the N<sub>2</sub>O concentration in 100 m water depth which is assumed to be brought to the surface by upwelling.  $Q_w$  and  $Q_t$  are scaling functions depending on the upwelling velocity and time history of the upwelling (for details and parameters used see *Nevison et al.* [2004]).  $C_{100m}$  was taken from N<sub>2</sub>O profiles measured during four cruises between 2006 and 2008 (Table 1) conducted by the SOPRAN (Surface Ocean Processes in the Anthropocene: [www.sopran.pangaea.de](http://www.sopran.pangaea.de)) project which is a German contribution to international SOLAS (Surface Ocean – Lower Atmosphere Study: [www.solas-int.org](http://www.solas-int.org)). N<sub>2</sub>O was measured according to the method described in *Walter et al.* [2006]. The data presented in Table 1 are in agreement with  $C_{100m}$  (32.4 and 29.0 nmol L<sup>-1</sup>) at two stations (20.6°N, 18.2°W and 20.3°N, 17.8°W, respectively) measured off Mauritania in September 2003 [*Forster et al.*, 2009].

[5] Wind data were derived from an active microwave sensor (SeaWinds scatterometer) onboard of the National Aeronautics and Space Administration (NASA) QuikSCAT satellite (<http://podaac.jpl.nasa.gov>), which measures near-surface values of the scalar wind speed and the meridional and zonal components of wind velocity and direction under all weather and cloud conditions over the ice-free ocean


[*Liu*, 2002]. The QuikSCAT wind data used in our study have a spatial resolution of 25 km × 25 km and provide one measurement for the grid cells per day. A comparison of monthly mean QuikSCAT winds for 2001 with measured wind data from buoys deployed worldwide revealed a good correlation ( $r = 0.92$ ), a root-mean-square difference of 0.7 m s<sup>-1</sup> and a standard deviation for the QuikSCAT winds of ±1.7 m s<sup>-1</sup> [*Wallcraft et al.*, 2009]. Rain events and the existence of gaps decrease the accuracy of the data [*Liu*, 2002; *Wallcraft et al.*, 2009], however, this constraint is negligible for the area discussed here. For further calculations the  $v$  (i.e., the N–S) component of the wind speed, which is driving the Ekman upwelling, was used.

[6] Following *Nevison et al.* [2004] we assumed that upwelling takes place when the winds were blowing for more than three days with a wind speed  $u_{crit} > 4.6$  m s<sup>-1</sup>.  $u_{crit}$  was determined as


$$u_{crit} = (\omega_{crit} \rho_w f R_b / (\rho_a C_d))^{0.5}, \quad (5)$$

where  $\omega_{crit}$  is the velocity threshold for the onset of upwelling (10<sup>-5</sup> m s<sup>-1</sup>),  $\rho_w$  and  $\rho_a$  are the densities of seawater (1025 kg m<sup>-3</sup>) and atmosphere (1.3 kg m<sup>-3</sup>), respectively;  $f$  is the Coriolis force (4.55 s<sup>-1</sup>),  $R_b$  is the Rossby radius of deformation (35.2 km) [*Emery et al.*, 1984] and  $C_d$  is the dimensionless drag coefficient (0.0012).  $\omega_{crit}$ ,  $\rho_w$  and  $\rho_a$  were adapted from *Nevison et al.* [2004].

[7] In order to estimate the annual N<sub>2</sub>O emissions from the upwelling off Mauritania, we divided the upwelling region into 10 boxes with a latitudinal spacing of 0.5° (Figure 1). For each of the given QuikSCAT coordinates within each latitudinal box,  $F_{N_2O}$  was calculated for the calendar years 2006–2008 as described above. On the basis of the data in Table 1 we computed mean annual  $C_{100m}$  of 31 ± 7 nmol L<sup>-1</sup> (2006), 28 ± 4 nmol L<sup>-1</sup> (2007) and 29 ± 4 nmol L<sup>-1</sup> (2008). Please note that potential seasonal and spatial variabilities of  $C_{100m}$  as implied by the variability of the concentrations which range from 21.9 ± 0.8 to 35.8 ± 0.3 nmol L<sup>-1</sup> (see Table 1) are not accounted for in our approach. This introduces an uncertainty in our computa-



**Figure 1.** Schematic representation of the coastal region off Mauritania. The points represent the coordinates of available QuikSCAT data within the boxes #1–#10 (see text for details.)



**Figure 2.** Time series of N<sub>2</sub>O flux densities for the boxes #1–#10 (see Figure 1). Periods without data originate from times when  $u_{\text{crit}} < 4.6 \text{ m s}^{-1}$  and indicate periods of no upwelling.

tions which cannot be quantified reasonably because of the rather poor temporal and spatial data coverage of the Mauritania upwelling area. The mean annual  $F_{\text{N}_2\text{O}}$  for each box was then calculated as the average of the individual annual  $F_{\text{N}_2\text{O}}$  within the box. Multiplying the mean annual flux densities with the area of the box yielded the mean annual N<sub>2</sub>O emissions for each box which were then added to yield the overall mean annual N<sub>2</sub>O emissions for the years 2006–2008.

#### 4. Results and Discussion

[8] Figure 2 shows the time series of N<sub>2</sub>O flux densities for the boxes #1–#10 for the year 2008. (The time series for 2006 and 2007 are very similar to the one from 2008 and are, therefore, not shown.) Two trends are visible: 1) The data from boxes #1–#3 (21°–19.5°N) show flux densities throughout the year, whereas the data from boxes #4–#10 (19.5°–16.0°N) show periods during the summer where no flux densities were computed because the upwelling criterion was not matched (cf. equation (5)). The period with no upwelling reaches its maximum duration (July to October) in the southern most box (16.5°–16.0°N). 2) In general the flux densities are higher in the northern boxes (maximum  $> 2 \text{ nmol m}^{-2} \text{ s}^{-1}$ ) and decrease considerably towards the south (maximum  $< 1 \text{ nmol m}^{-2} \text{ s}^{-1}$ ). The N<sub>2</sub>O flux densities are obviously driven by the magnitude and seasonality of the wind speeds and, thus, reflect the seasonality of the upwelling off Mauritania (see study site description). However, the simple approach applied here does not take into account potential stratification of the water column and currents. Both effects are known to reduce the effective Ekman upwelling and, therefore, the N<sub>2</sub>O flux densities presented in Figure 2 are most likely overestimated. Calculations with a 3D high resolution model for the Atlantic Ocean (FLAME, Family of Linked Atlantic Model Experiments; <http://www.ifmgeomar.de/index.php?id=3239&L=1>) [Glessmer *et al.*, 2009] revealed that the vertical upwelling velocities of the Nevison *et al.*'s [2004] approach are well

correlated with the seasonality of the vertical velocity in 100 m, however, they were in general about 1.5 times higher. A correction factor of 0.68 was calculated based on the mean difference of the Ekman upwelling and the vertical velocity in 100 m averaged for the area 16°–18.5°W, 16°–21°N [Wittke, 2009]. The correction factor was applied to adjust the N<sub>2</sub>O flux densities: The uncorrected and corrected N<sub>2</sub>O emissions for the individual calendar years range from 0.9 to 1.1 Gg N yr<sup>-1</sup> and 0.5 to 0.7 Gg N yr<sup>-1</sup>, respectively. The mean total N<sub>2</sub>O emissions from the Mauritanian upwelling for 2006–2008 are 1.0 Gg N yr<sup>-1</sup> (corrected: 0.6 Gg N yr<sup>-1</sup>). Because the estimated error of the N<sub>2</sub>O emission for the individual calendar years is about  $\pm 2 \text{ Gg N yr}^{-1}$  [Wittke, 2009] it is impossible to detect any interannual variability or long-term trend.

[9] For comparison, Nevison *et al.*'s [2004] estimate for the upwelling off NW Africa ( $2.4 \cdot 10^5 \text{ km}^2$  between 5°–30°N) is 22 Gg N yr<sup>-1</sup>. Downscaling their estimate to the upwelling area considered in our study, which is considerably smaller ( $3.7 \cdot 10^4 \text{ km}^2$  between 16° and 21° N), yields 3.4 Gg N yr<sup>-1</sup>. Thus, Nevison *et al.*'s [2004] estimate appears to be considerably higher than our estimate. However, there are two caveats associated with their estimate: 1) For the computation of  $k_w$ , Nevison *et al.* [2004] applied the air-sea gas exchange approach of Wanninkhof [1992] which generally yields  $k_w$  significantly higher than those computed with Nightingale *et al.*'s [2000] approach used in our study. Using Nightingale *et al.*'s [2000] approach is in agreement with recent studies, which suggest that Wanninkhof's [1992] relationship is overestimating  $k_w$  by 33% [Wanninkhof *et al.*, 2009]. 2) Because of missing data, Nevison *et al.* [2004] had to estimate N<sub>2</sub>O in 100 m with an empirical N<sub>2</sub>O/O<sub>2</sub> relationship and they explicitly state that their estimate may be associated with an error of  $\pm 30\%$ . Thus, we conclude that Nevison *et al.*'s [2004] estimate is overestimated.

[10] Despite the fact that the Mauritanian upwelling area considered in our study represents about 2% of the global coastal upwelling ( $1.75 \cdot 10^6 \text{ km}^2$  as given by Nevison *et al.* [2004]), its N<sub>2</sub>O emissions contribute only about 0.5%

(corrected: 0.3%) of the global N<sub>2</sub>O upwelling emissions of  $200 \pm >140 \text{ Gg N yr}^{-1}$  [Nevison et al., 2004]. Therefore, we conclude that N<sub>2</sub>O emissions from the Mauritanian upwelling are only of minor importance for the atmospheric N<sub>2</sub>O budget and they do not represent a globally significant ‘hot spot’ of N<sub>2</sub>O emissions. Thus, in view of its N<sub>2</sub>O emissions, the Mauritanian upwelling is different from other major coastal upwelling areas, such as those found in the Arabian Sea or off Chile [Naqvi et al., 2005; Paulmier et al., 2008]. For example, N<sub>2</sub>O emissions along the coast of the Arabian Peninsula have been estimated to range from 47–90 Gg N during the upwelling period [Bange et al., 1996b]. One reason for the comparatively low total N<sub>2</sub>O emissions from off Mauritania is the fact that the subsurface oxygen minimum zone (OMZ) off Mauritania is not suboxic, i.e., the O<sub>2</sub> concentrations are always  $>40 \mu\text{mol L}^{-1}$  [Karstensen et al., 2008; Naqvi et al., 2009]. Suboxic conditions, however, would favor accumulation of N<sub>2</sub>O in subsurface layers via enhanced nitrification and/or denitrification [Bange, 2008]. This, in turn, can result in very high N<sub>2</sub>O emissions as have been shown only recently for the upwelling off Chile [Paulmier et al., 2008].

## 5. Summary

[11] N<sub>2</sub>O flux densities across the ocean/atmosphere interface from the Mauritanian upwelling (16°–21°N) were computed using the “wind-only” upwelling model approach of Nevison et al. [2004]. N<sub>2</sub>O measurements from four cruises between 2006 and 2008 as well as winds from the QuikSCAT satellite were used as input data. The resulting N<sub>2</sub>O flux densities showed a strong seasonality, reflecting the wind-driven seasonality of the upwelling off Mauritania: computed N<sub>2</sub>O flux densities are highest in the northern part (19.5°–21°N) and show a decreasing trend towards the south. The summer periods with no upwelling (and thus resulting in no or negligible N<sub>2</sub>O fluxes densities) are most pronounced in the southern part (16°–17°N). The mean total N<sub>2</sub>O emissions from the Mauritanian upwelling were estimated to  $1.0 \text{ Gg N yr}^{-1}$  indicating that N<sub>2</sub>O emissions from the Mauritanian upwelling are only of minor importance for the atmospheric N<sub>2</sub>O budget.

[12] **Acknowledgments.** We thank the authorities of Mauritania for the permission to work in their territorial waters. We acknowledge the support of the captains and crews of R/V Poseidon, R/V Meteor and R/V L’A-talante as well as the chief scientists Arne Körtzinger (M68/3, ATA3) and Marcus Dengler (P347). Moreover, we thank A. Freing, B. Fiedler, I. Grefe and T. Tanhua for their help with the N<sub>2</sub>O sampling. We thank Alina Freing, Mirjam Glessmer and Dietmar Dommengot for helpful discussions. Mirjam Glessmer kindly provided the FLAME model results. Moreover, we acknowledge the helpful comments of two anonymous reviewers. This work was funded by the German BMBF joint project SOPRAN (FKZ 03F0462A) and is a contribution to SOLAS.

## References

- Antonov, J. I., et al. (2006), *World Ocean Atlas 2005*, vol. 2, *Salinity*, NOAA Atlas NESDIS, vol. 62, edited by S. Levitus, 182 pp., NOAA, Silver Spring, Md.
- Bange, H. W. (2006), New directions: The importance of the oceanic nitrous oxide emissions, *Atmos. Environ.*, *40*, 198–199, doi:10.1016/j.atmosenv.2005.09.030.
- Bange, H. W. (2008), Gaseous nitrogen compounds (NO, N<sub>2</sub>O, N<sub>2</sub>, NH<sub>3</sub>) in the ocean, in *Nitrogen in the Marine Environment*, 2nd ed., edited by D. G. Capone et al., pp. 51–94, doi:10.1016/B978-0-12-372522-6.00002-5, Elsevier, Amsterdam.
- Bange, H. W., S. Rapsomanikis, and M. O. Andreae (1996a), Nitrous oxide in coastal waters, *Global Biogeochem. Cycles*, *10*, 197–207, doi:10.1029/95GB03834.
- Bange, H. W., S. Rapsomanikis, and M. O. Andreae (1996b), Nitrous oxide emissions from the Arabian Sea, *Geophys. Res. Lett.*, *23*, 3175–3178, doi:10.1029/96GL03072.
- Bange, H. W., et al. (2001), Nitrous oxide emissions from the Arabian Sea: A synthesis, *Atmos. Chem. Phys.*, *1*, 61–71, doi:10.5194/acp-1-61-2001.
- Cornejo, M., et al. (2006), Temporal variability in N<sub>2</sub>O water content and its air-sea exchange in an upwelling area off central Chile (36°S), *Mar. Chem.*, *101*, 85–94, doi:10.1016/j.marchem.2006.01.004.
- Denman, K. L., et al. (2007), Couplings between changes in the climate system and biogeochemistry, in *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by S. Solomon et al., pp. 499–587, Cambridge Univ. Press, Cambridge, U. K.
- Emery, W. J., et al. (1984), Geographic and seasonal distribution of Brunt-Vaisala frequency and Rossby radii in the North Pacific and North Atlantic, *J. Phys. Oceanogr.*, *14*, 294–317, doi:10.1175/1520-0485(1984)014<0294:GASDOB>2.0.CO;2.
- Forster, F. G., R. C. Upstill-Goddard, N. Gist, C. Robinson, G. Uher, and E. M. S. Woodward (2009), Nitrous oxide and methane in the Atlantic Ocean between 50°N and 52°S: Latitudinal distribution and sea-to-air flux, *Deep Sea Res., Part II*, *56*, 964–976, doi:10.1016/j.dsr2.2008.12.002.
- Forster, P., et al. (2007), Changes in atmospheric constituents and in radiative forcing, in *Climate Change 2007: The Physical Science Basis. Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*, edited by S. Solomon et al., pp. 129–234, Cambridge Univ. Press, Cambridge, U. K.
- Glessmer, M. S., et al. (2009), Contribution of oxygen minimum zone waters to the coastal upwelling off Mauritania, *Prog. Oceanogr.*, *83*, 143–150, doi:10.1016/j.pocan.2009.07.015.
- Hagen, E. (2001), Northwest African upwelling scenario, *Oceanol. Acta*, *24*, 113–127, doi:10.1016/S0399-1784(00)01110-5.
- Karstensen, J., et al. (2008), The oxygen minimum zones in the eastern tropical Atlantic and Pacific oceans, *Prog. Oceanogr.*, *77*, 331–350, doi:10.1016/j.pocan.2007.05.009.
- Liu, W. T. (2002), Progress in scatterometer application, *J. Oceanogr.*, *58*, 121–136, doi:10.1023/A:1015832919110.
- Locarnini, R. A., et al. (2006), *World Ocean Atlas 2005*, vol. 1, *Temperature*, NOAA Atlas NESDIS, vol. 61, edited by S. Levitus, 182 pp., NOAA, Silver Spring, Md.
- Lueker, T. J., et al. (2003), Coastal upwelling air-sea fluxes revealed in atmospheric observations of O<sub>2</sub>/N<sub>2</sub>, CO<sub>2</sub> and N<sub>2</sub>O, *Geophys. Res. Lett.*, *30*(6), 1292, doi:10.1029/2002GL016615.
- Mittelstaedt, E. (1986), Upwelling regions, in *Landoldt-Börnstein*, vol. 3, *Oceanography*, edited by J. Sündermann, pp. 135–166, Springer, Berlin.
- Naqvi, S. W. A., et al. (2005), Biogeochemical ocean-atmosphere transfers in the Arabian Sea, *Prog. Oceanogr.*, *65*, 116–144, doi:10.1016/j.pocan.2005.03.005.
- Naqvi, S. W. A., et al. (2009), Coastal hypoxia/anoxia as a source of CH<sub>4</sub> and N<sub>2</sub>O, *Biogeosci. Discuss.*, *6*, 9455–9523, doi:10.5194/bgd-6-9455-2009.
- Nevison, C. D., T. J. Lueker, and R. F. Weiss (2004), Quantifying the nitrous oxide source from coastal upwelling, *Global Biogeochem. Cycles*, *18*, GB1018, doi:10.1029/2003GB002110.
- Nightingale, P. D., G. Malin, C. S. Law, A. J. Watson, P. S. Liss, M. I. Liddicoat, J. Boutin, and R. C. Upstill-Goddard (2000), In situ evaluation of air-sea gas exchange parameterizations using novel conservative and volatile tracers, *Global Biogeochem. Cycles*, *14*, 373–387, doi:10.1029/1999GB900091.
- Nykjær, L., and L. Van Camp (1994), Seasonal and interannual variability of coastal upwelling along northwest Africa and Portugal from 1981 to 1991, *J. Geophys. Res.*, *99*, 14,197–14,207, doi:10.1029/94JC00814.
- Paulmier, A., et al. (2008), The oxygen minimum zone (OMZ) off Chile as intense source of CO<sub>2</sub> and N<sub>2</sub>O, *Cont. Shelf Res.*, *28*, 2746–2756, doi:10.1016/j.csr.2008.09.012.
- Prinn, R. G., et al. (2000), A history of chemically and radiatively important gases in air deduced from ALE/GAGE/AGAGE, *J. Geophys. Res.*, *105*, 17,751–17,792, doi:10.1029/2000JD900141.
- Ravishankara, A. R., et al. (2009), Nitrous oxide (N<sub>2</sub>O): The dominant ozone-depleting substance emitted in the 21st century, *Science*, *326*, 123–125, doi:10.1126/science.1176985.
- Schemainda, R., et al. (1975), Ozeanologische Untersuchungen zum Produktionspotential der nordwestafrikanischen Wasserauftriebsregionen 1970–1973, *Geod. Geophys. Veröffentlichungen*, *IV*, 1–88.
- Wallcraft, A. J., A. B. Kara, C. N. Barron, E. J. Metzger, R. L. Pauley, and M. A. Bourassa (2009), Comparisons of monthly mean 10 m wind speeds from satellites and NWP products over the global ocean, *J. Geophys. Res.*, *114*, D16109, doi:10.1029/2008JD011696.

- Walter, S., H. W. Bange, and D. W. R. Wallace (2004), Nitrous oxide in the surface layer of the tropical North Atlantic Ocean along a west to east transect, *Geophys. Res. Lett.*, *31*, L23S07, doi:10.1029/2004GL019937.
- Walter, S., et al. (2006), Nitrous oxide in the North Atlantic Ocean, *Biogeosciences*, *3*, 607–619, doi:10.5194/bg-3-607-2006.
- Wanninkhof, R. (1992), Relationship between wind speed and gas exchange over the ocean, *J. Geophys. Res.*, *97*, 7373–7382, doi:10.1029/92JC00188.
- Wanninkhof, R., et al. (2009), Advances in quantifying air-sea gas exchange and environmental forcing, *Annu. Rev. Mar. Sci.*, *1*, 213–244, doi:10.1146/annurev.marine.010908.163742.
- Weiss, R. F., and B. A. Price (1980), Nitrous oxide solubility in water and seawater, *Mar. Chem.*, *8*, 347–359, doi:10.1016/0304-4203(80)90024-9.
- Wittke, F. (2009), Ozean/Atmosphäre-Wechselwirkungen und N<sub>2</sub>O Emissionen in der Küstenauftriebsregion vor Mauretanien, Diploma thesis, 88 pp., Univ. of Kiel, Kiel, Germany.
- Wooster, W. S., et al. (1976), The seasonal upwelling cycle along the eastern boundary of the North Atlantic, *J. Mar. Res.*, *34*, 131–141.

---

H. W. Bange, A. Kock, and F. Wittke, Forschungsbereich Marine Biogeochemie, Leibniz-Institut für Meereswissenschaften an der Universität Kiel (IFM-GEOMAR), Düsternbrooker Weg 20, D-24105 Kiel, Germany. (hbange@ifm-geomar.de)